



HOSTED BY



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://ees.elsevier.com/ejbas/default.asp>

Full Length Article

Traveling wave solutions of generalized Zakharov–Kuznetsov–Benjamin–Bona–Mahony and simplified modified form of Camassa–Holm equation $\exp(-\phi(\eta))$ – Expansion method



Ayyaz Ali, Muhammad Asad Iqbal, Syed Tauseef Mohyud-Din *

Department of Mathematics, Faculty of Sciences, HITEC University Taxila, Pakistan

ARTICLE INFO

Article history:

Received 20 May 2015

Received in revised form 7 January 2016

Accepted 7 January 2016

Available online 16 March 2016

Keywords:

Exp(- $\phi(\eta)$)-expansion method
 Nonlinear evolution equation
 Generalized Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation
 Simplified Modified form of Camassa–Holm equation
 Traveling wave solutions

ABSTRACT

In this article, we established abundant traveling wave solutions for nonlinear evolution equations. The $\exp(-\phi(\eta))$ -expansion method is used to construct traveling wave solutions for the generalized Zakharov–Kuznetsov–Benjamin–Bona–Mahony equation and Simplified Modified form of Camassa–Holm equation. The traveling wave solutions are expressed in terms of the hyperbolic functions, the trigonometric functions and the rational functions. The proposed solutions are found to be important for the explanation of some practical physical problems in mathematical physics and engineering.

© 2016 Mansoura University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

It is well known that seeking exact solutions [1–53] for nonlinear evolution equations (NLEEs) plays an important role in mathematical physics. For instance, nonlinear evolution equations (NEEs) are widely used as models to describe complex physical phenomena in various fields of sciences, especially

in fluid mechanics, solid-state physics, plasma physics, plasma waves and biology. One of the basic physical problems for those models is to obtain their travelling wave solutions. In particular, various methods have been utilized to explore different kinds of solutions of physical models described by nonlinear partial differential equations (NPDEs). In the past few decades or so, many effective methods have been presented, which contain the inverse scattering transform method, the Backlund

* Corresponding author. Tel.: +92 3235577701.

E-mail address: syedtauseefs@hotmail.com (S.T. Mohyud-Din).

<http://dx.doi.org/10.1016/j.ejbas.2016.01.001>

2314-808X/© 2016 Mansoura University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

transformation [1], bilinear transformation, the tanh-sech method [2], the extended tanh method, the pseudo-spectral method [3,8-10,14,15], trial function and the sine-cosine method [4,5], Hirota method [6], tanh-coth method [2,7,11,13], the exponential function method [16-24], the (G/G)-expansion method [25-29], the homogeneous balance method [30,31], F-expansion method [33-35] and the Jacobi elliptic function expansion method [36-38] and so on. In a subsequent work, Ma et al. [39] developed the complexiton solutions for Toda lattice equation through the Casoratian formulation and hence obtained a set of coupled conditions which guaranteed Casorati determinants to be the solution of Toda Lattice which consequently produced complexiton solutions. Moreover, Ma and You [40] used variation of parameters for solving the involved non-homogeneous partial differential equations and obtained solution formulas helpful in constructing the existing solutions coupled with a number of other new solutions including rational solutions, solitons, positons, negatons, breathers, complexions and interaction solutions of the KdV equations. It is needed to be highlighted that the basic spirit of the exp-function method which is the conversion of nonlinear partial differential equations into integrable ordinary differential equations was explicitly presented and minutely analyzed in 1996 by Ma and Fuchssteiner [41]. In fact, the exp-function method is restricted to produce rational solutions in the form of transformed variables and such solutions can be obtained easily by making use of other techniques including Wronskian and Casoratian [41-43]. Recently, Ma, Wu and He [44] presented a much more general idea to yield exact solutions to nonlinear wave equations by searching for the so-called Frobenius transformations. Some recently developed methods, such as, the modified simple equation [45-49], the enhanced $\text{Exp}(-\varphi(\xi))$ -expansion method [50,51], the Enhanced (G'/G)-Expansion method [52,53], etc. which provide useful exact solutions to NLEEs have been discussed.

The objective of this article is to apply the $\text{exp}(-\varphi(\eta))$ -expansion method to construct the exact solutions for nonlinear evolution equations in mathematical physics via generalized Zakharov-Kuznetsov-Benjamin-Bona-Mahony equation and Simplified Modified form of Camassa-Holm equation. The subject matter of this method is that the traveling wave solutions of a nonlinear evolution equation can be expressed by a polynomial in $\text{exp}(-\varphi(\eta))$, where $\varphi(\eta)$ satisfies the ordinary differential equation (ODE):

$$(\varphi'(\eta)) = \text{exp}(-\varphi(\eta)) + \mu \text{exp}(-\varphi(\eta)) + \lambda \tag{1}$$

Where $\eta = x - Vt$.

2. Description of $\text{exp}(-\varphi(\eta))$ -expansion method

Now we explain the $\text{exp}(-\varphi(\eta))$ -expansion method for finding traveling wave solutions of nonlinear evolution equations. Let us consider the general nonlinear partial differential equation of the form.

$$P(u, u_t, u_x, u_{tt}, u_{xx}, u_{xxx}, \dots), \tag{2}$$

where $u = u(x, t)$ is an unknown function, P is a polynomial in $u(x, t)$ and its various partial derivatives, in which the highest order derivatives and nonlinear terms are involved. In order to solve Eq. (2) by using the $\text{exp}(-\varphi(\eta))$ -expansion method we have to follow the following steps.

Step 1. Combining the real variables x and t by a compound variable η we assume

$$u(x, t) = u(\eta), \quad \eta = x - Vt \tag{3}$$

where V is the speed of the traveling wave. Using the traveling wave variable (3), Eq. (2) is reduced to the following ODE for $u = u(\eta)$

$$Q(u, u', u'', u''', u''', \dots) = 0, \tag{4}$$

where Q is a function of $u(\eta)$ and its derivatives, prime denotes derivative with respect to η .

Step 2. Suppose the solution of (4) can be expressed by a polynomial in $\text{exp}(-\varphi(\eta))$ as follows

$$u(\eta) = a_n (\text{exp}(-\varphi(\eta)))^n + a_{n-1} (\text{exp}(-\varphi(\eta)))^{n-1} + \dots, \tag{5}$$

where a_n, a_{n-1}, \dots and V are constants to be determined later such that $a_n \neq 0$ and $\varphi(\eta)$ satisfies Eq. (1).

Step 3. By using the homogeneous principle, we can evaluate the value of positive integer n between the highest order linear terms and nonlinear terms of the highest order in Eq. (4). Our solutions now depend on the parameters involved in Eq. (1).

Case 1. $\lambda^2 - 4\mu > 0$ and $\mu \neq 0$,

$$\varphi(\eta) = \ln \left\{ \frac{1}{2\mu} \left(-\sqrt{\lambda^2 - 4\mu} \tanh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} (\eta + c_1) \right) - \lambda \right) \right\}, \tag{6}$$

where c_1 is a constant of integration.

Case 2. $\lambda^2 - 4\mu < 0$ and $\mu \neq 0$,

$$\varphi(\eta) = \ln \left\{ \frac{1}{2\mu} \left(-\lambda + \sqrt{-\lambda^2 + 4\mu} \tan \left(\frac{\sqrt{-\lambda^2 + 4\mu}}{2} (\eta + c_1) \right) \right) \right\} \tag{7}$$

Case 3. $\mu = 0$ and $\lambda \neq 0$,

$$\varphi(\eta) = -\ln \left\{ \frac{\lambda}{\text{exp}(\lambda(\eta + c_1)) - 1} \right\} \tag{8}$$

Case 4. $\lambda^2 - 4\mu = 0, \lambda \neq 0$, and $\mu \neq 0$,

$$\varphi(\eta) = \ln \left\{ \frac{2(\lambda(\eta + c_1) + 2)}{(\lambda^2(\eta + c_1))} \right\} \tag{9}$$

Case 5. $\lambda = 0$, and $\mu = 0$,

$$\varphi(\xi) = \ln(\eta + c_1) \tag{10}$$

Download English Version:

<https://daneshyari.com/en/article/560947>

Download Persian Version:

<https://daneshyari.com/article/560947>

[Daneshyari.com](https://daneshyari.com)