



Original Research Article (Experimental)

Involvement of tyrosine-specific protein kinase and protein kinase C in J774A.1 macrophage functions activated by *Tinospora cordifolia*



Priti More, Kalpana Pai*

Department of Zoology, S. P. Pune University, India

ARTICLE INFO

Article history:

Received 23 November 2015

Received in revised form

7 December 2016

Accepted 28 December 2016

Available online 7 June 2017

Keywords:

Biological response modifiers

Guduchi

Macrophage activation

Tyrosine-specific protein kinase

Protein kinase C

ABSTRACT

Background: Macrophages are the first line of defense and constitute important participant in the bi-directional interaction between innate and specific immunity. Macrophages are in a quiescent form and get activated when given a stimulus. In our previous studies we have reported that guduchi or LPS treatment of macrophages enhanced production of nitric oxide (NO) and increased tumoricidal activity against L929 fibroblast cells.

Objective: In the present study effect of *Tinospora cordifolia* commonly known as guduchi on macrophage activation and the mechanism of action i.e. involvement of protein kinase C inhibitor and tyrosine-specific protein kinase inhibitor was investigated.

Materials and Methods: The present study was undertaken to determine whether H-7 (inhibitor of protein kinase C) and/or genistein (inhibitor of tyrosine-specific protein kinase) could inhibit guduchi or LPS-induced macrophage NO and TNF- α production or reduce the cytolysis of L929 fibroblast cells.

Results: It was observed that *in vitro* incubation with H-7 and/or genistein completely inhibited guduchi or LPS-induced NO and TNF- α production by macrophages (J774A.1).

Conclusion: The inhibitory effects of H-7 and/or genistein, suggest that phosphorylation via these kinases may upregulate the NO synthase activity in macrophages.

© 2017 Transdisciplinary University, Bangalore and World Ayurveda Foundation. Publishing Services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Macrophages are quiescent cells, which get activated when stimulated. Different types of agents such as antibiotics, anti-metabolites and cytokines may exert an immunomodulating action that is expressed in the augmentation and/or inhibition of different immune responses [15]. One of the most promising recent alternatives to classical antibiotic treatment is the use of immunomodulators for enhancing host defense responses [23]. Number of natural products and synthetic immunopotentiators termed as Biological Response Modifiers (BRMs) are becoming increasingly popular for their potential in augmenting immune responses. Among the natural BRMs many herbs and medicinal plants have long been known

for their immunoaugmentary potential; however, only recently scientists have recognized them for their possible BRM actions. The herb guduchi isolated from botanical sources have attracted a great deal of attention in the biomedical arena because of its broad spectrum of therapeutic properties and relatively low toxicity. Our study is based on the investigation of the BRM *Tinospora cordifolia* (guduchi) as an activating agent of macrophages *in vitro* and the bacterial endotoxin LPS as a positive control.

In the present study two different inhibitors (H-7), an inhibitor of protein kinase C and/or (genistein), a tyrosine-specific protein kinase inhibitor were used to analyze the regulation of guduchi or LPS mediated macrophage activation. Protein kinase C (PKC) has been shown to be a signal transducer during tumorigenesis, tumor cell invasion, and metastasis. Recent studies have reported that the PKC inhibitor, 7-hydroxystaurosporine (H-7), inhibits tumor cell invasion [22]. Genistein, an isoflavone, has been shown to inhibit cell proliferation and enhance apoptosis in cancer cells. Accumulating bodies of evidence suggest that genistein is expected to synergistically promote the anti-proliferative effects of chemotherapeutic agents on neoplasia without toxicity [24]. In the present study involvement of these specific kinases, in

* Corresponding author. Department of Zoology, Centre of Advanced Studies, Savitribai Phule Pune University (Formerly University of Pune), Ganeshkhind, Pune, 411007, Maharashtra, India.

E-mail address: drkalpanapai@gmail.com (K. Pai).

Peer review under responsibility of Transdisciplinary University, Bangalore.

guduchi or LPS-mediated macrophage activation has been investigated.

Macrophage activation is known to occur through a series of stages ranging from a level equivalent to resident tissue macrophages and culminating at an activated state whereupon macrophages become competent to kill several pathogens and lyse tumor cells [1,14]. Although activated macrophages produce a number of physiologically active molecules with cytotoxic and/or cytostatic effects, only interleukin-1 (IL-1), tumor necrosis factor (TNF) and reactive nitrogen intermediates (RNI) have been clearly implicated in monocyte/macrophage mediated tumor cytotoxicity [2,5,8,9,13,21].

Nitric oxide (NO) is formed enzymatically from a terminal guanidine-nitrogen of L-arginine by the so called NO synthases (NOSs) that yield L-citrulline as a co-product [16,17]. Both in the NO producing cell and in specific NO target cells, NO functions as the first messenger of nitrinergic signal transduction, activating GC-S [GTP pyrophosphate-lyase (cyclizing)] [3] and thereby increasing the intracellular concentration of the second messenger molecule cGMP [4,18]. Whether the complete pathway operates in macrophages has not been investigated. In guduchi activated macrophages NO biosynthesis could also be the mediator of macrophage-mediated tumor cytotoxicity [7,9–11,21]. While our understanding of the mechanism of action of this BRM is still developing, it appears that the primary mechanism involves induction of the immune system. Our previous studies demonstrate that the basic mechanism of the immunostimulatory, antitumor, bactericidal and other therapeutic effects of guduchi is thought to occur via macrophage stimulation [10–12]. We have focused this study on the involvement of protein kinase C and tyrosine-specific protein kinase in the BRM (guduchi) or LPS-mediated macrophage functions. The purpose of the present study was to investigate whether H-7 (inhibitor of protein kinase C) and/or genistein (inhibitor of tyrosine-specific protein kinase) could decrease macrophage derived NO and TNF- α production in the setting of an *in vitro* herbal (guduchi) treatment or endotoxin (LPS) challenge.

2. Materials and methods

Reagents: Dulbecco's Modified Eagle Medium (DMEM) with L-glutamine and 25 mM HEPES buffer were purchased from (HiMedia Pvt. Ltd. India.) Fetal bovine serum was purchased from Hyclone (Logan, USA) and heat inactivated at 56 °C for 30 min. The whole plant extract of *T. cordifolia* was used for the study. The plant was obtained from medicinal plant nursery, Pune, Maharashtra. The plant was subjected to extraction with 200 ml methanol at 50 °C for 8 cycles by Soxhlet extraction process. The extract was then concentrated with rotator vacuum evaporator and used for further analysis. For enzymatic analysis, fresh crude extract with phosphate buffer (pH 7) was used. The guduchi extract prepared in incomplete DMEM were tested for endotoxin contamination by limulus amoebocyte lysate assay which showed insignificant levels [0.0007 ng/mg]. Necessary precautions were taken to avoid endotoxin contamination through out the investigation, by using endotoxin free buffers, reagents and sterile water. All other chemicals and solvents used in this study were obtained from Sigma Chemical Company (St. Louis, USA) and were of analytical grade or the highest grade available.

Cells: The macrophage J774A.1 and the fibroblast L929 cell lines were obtained from National Center for Cell Sciences (NCCS, Pune). J774A.1 cell line was used as source of macrophages, (Origin: BALB/c mouse; Nature: Mature) grown and maintained in the Dulbecco's Modified Eagle Medium (DMEM) (pH 7.5) enriched with 10% fetal bovine serum, at 37 °C and 5% CO₂.

2.1. Viability assay

Cell viability was determined by the trypan blue dye exclusion technique. Equal volumes of cell suspensions were mixed with 0.4% trypan blue in PBS, and the unstained viable cells were determined. These cells were further used for cytotoxicity assay in 2×10^6 densities per ml in the 96 well tissue culture plates.

Stimulation of macrophages: The macrophage cells (cell line J774A.1) from late log phase of growth (subconfluent) were seeded in 96 well flat bottom microtiter plates (Tarsons, India) in a volume of 100 μ l under adequate culture conditions. Guduchi (80 μ g/well) or LPS (10 μ g/well) were added in a volume of 100 μ l with and/or without inhibitors in triplicate. The cultures were incubated at 37 °C and 5% CO₂ environment. After 24 h and 48 h incubation percent viability was checked and culture supernatants were collected and assayed for nitric oxide and TNF- α activity.

2.2. Inhibitor treatment to the macrophages as second messengers

Along with guduchi or LPS macrophage cells were treated with two different inhibitors, H-7 (inhibitor of protein kinase C) and/or genistein (inhibitor of tyrosine-specific protein kinase), within a concentration of 10 μ M/ml, respectively. Supernatant collected from these cells were assayed for nitrite generation and for tumoricidal activity to check inhibition.

2.3. Nitrite assay

The concentration of stable nitrite, an end product of the nitric oxide present in the supernatant of treated or untreated J774A macrophage cell cultures (2×10^6 cells/ml), was measured by the method of Ding et al. [6] based on the Griess reaction [20]. Briefly, 50 μ l of supernatant was incubated with an equal volume of Griess reagent (1% sulphanilamide in 2.5% H₃PO₄ and 0.1% naphthyl-ethylene-diaminedihydrochloride in distilled water; both solutions mixed in a ratio of 1:1 at room temperature) for 10 min. The absorbance at 550 nm was then measured in a microtitre plate reader. The standard curve for nitrite was prepared by using 10–100 μ M sodium nitrite in distilled water.




Fig. 1. Nitrite levels in macrophages (J774A.1) after treatment with guduchi or LPS along with the inhibitors. Values represent nitrite content in μ M. Bars show mean \pm SD for triplicate cultures. Data are representative of three separate experiments (* $p < 0.05$; versus value of guduchi or LPS treated macrophages) [w/o: without inhibitor].

Download English Version:

<https://daneshyari.com/en/article/5635265>

Download Persian Version:

<https://daneshyari.com/article/5635265>

[Daneshyari.com](https://daneshyari.com)