



# Predictors of hepatitis C testing intention among African American Baby Boomers



Mohamed Rashrash<sup>a,\*</sup>, Mary Maneno<sup>b</sup>, Anthony Wutoh<sup>b</sup>, Earl Ettienne<sup>b</sup>, Monika Daftary<sup>b</sup>

<sup>a</sup> Chapman University School of Pharmacy, 9401 Jeronimo Road, Irvine, CA 92618, USA

<sup>b</sup> Department of Clinical Social Administrative Pharmacy, Howard University College of Pharmacy, 2300 4th Street, NW Washington, DC 20059, USA

## ARTICLE INFO

### Article history:

Received 9 September 2015

Received in revised form 10 December 2016

Accepted 15 December 2016

Available online 16 February 2017

### Keywords:

African-American  
Baby Boomers  
Health Belief Model  
Hepatitis C  
Intention

## ABSTRACT

Baby Boomers (BBs) are responsible for three-quarters of hepatitis C virus (HCV) infections in the United States; however, HCV testing is distinctly underused by them. A cross-sectional study was conducted to assess the prevalence of HCV testing and to evaluate predictors of HCV testing intention among African-American BBs. The study was guided by the Health Belief Model and theory of reasoned action frameworks. Of the 137 participants included in the study, 44.8% had at least a college education; 13.9% received prior to 1992 blood transfusion. Findings related to HCV testing showed that 32.1% of the participants intended to test for HCV within 6 months and 43.8% had received a previous HCV test. Significant predictors of HCV testing intention within 6 months included having a blood transfusion prior to 1992 [odds ratio (OR) = 8.25, 95% confidence interval (CI): 2.02–33.61], perceptions of benefits (OR = 1.57, 95% CI: 1.13–2.18), severity (OR = 1.39, 95% CI: 1.17–1.65), and subjective norms (OR = 1.42, 95% CI: 1.12–1.79). These predictors of HCV testing intention can be used to develop future HCV testing initiatives for African-American BBs.

© 2017 Ministry of Health, Saudi Arabia. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Hepatitis C virus (HCV) infection started as an epidemic between the 1960s and 1980s [1]. The World Health Organization estimates that >170 million people are living with HCV worldwide [2]. This infection is a primary public health concern among ~4 million people infected with HCV in the United States; particularly among people born between 1945 and 1965, for whom this is the main blood borne infection [1,3,4]. This population subgroup, commonly known as Baby Boomers (BBs), has five times the risk of HCV than other groups [5].

Encouraging testing and treatment for BBs is critical. However, these efforts have been challenging because BBs are mostly unaware of their HCV vulnerability and status [6]. Evidence suggests underuse of HCV testing services, despite the earlier promotion of testing by the United States Centers for Disease Control and Prevention (CDC) in 1998 [7]. Recently updated recommendations from the CDC published in August 2012 emphasize that all BBs should have a one-time HCV test to prevent adverse health

consequences [5]. Given the evidence of a low perception of susceptibility to HCV [6], it is important to examine whether this is a predictor of intention to test for it in this population.

A review of the literature review has shown a scarcity of theory-based research examining factors that predict or explain the willingness of BBs to have an HCV test. Specifically, the effects of variables related to the behavioral theories, such as knowledge perceptions, cues to action, and subjective norms [8,9], on intention to test for HCV have not been well investigated. Lack of such theory-driven HCV testing studies among African-American (AA) BBs is also evident in the literature. Given that sociodemographic and psychosocial factors determine health-seeking behaviors, it would be significant to understand which of these factors affect intention to test for HCV among AABBs.

The Health Belief Model (HBM) and the theory of reasoned action (TRA) include several psychosocial constructs that can be used to model preventive seeking behavior [10,11]; however, few studies have used them to evaluate HCV testing behavior. The purpose of this study was to assess the prevalence of HCV testing and to determine whether HBM/TRA constructs, prior HCV testing and other sociodemographic characteristics predict intention to testing for HCV among AABBs. We were particularly interested in testing whether the perception of susceptibility would affect intention to test for HCV.

Peer review under responsibility of Ministry of Health, Saudi Arabia.

\* Corresponding author at: 11450 Maple Drive, Fishers, IN 46038, USA.

E-mail address: [mrashrash@gmail.com](mailto:mrashrash@gmail.com) (M. Rashrash).

<http://dx.doi.org/10.1016/j.jegh.2016.12.005>

2210-6006/© 2017 Ministry of Health, Saudi Arabia. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 2. Participants and methods

### 2.1. Study design and population

The study had a cross-sectional design and included a convenience sample of 137 persons. The study started on February 1, 2014 and continued to March 30, 2014. Respondents recruited were AAs residing in Washington DC area, aged 44–69 years, and who were visiting or receiving service at Howard University Hospital and Ms. Bernice Elizabeth Fonteneau Senior Wellness Center in Washington DC. Respondents who were incapable of using the audio computer-assisted self-interview (ACASI) system were excluded from the study. A gift card for 10 US dollars was provided as an incentive. This study was approved by the Howard University Institutional Review Board.

### 2.2. Sample size calculation

A total of 137 participants were included in the study; 80 of them were recruited from Howard University Hospital and the remaining 57 were from Ms. Bernice Elizabeth Fonteneau Senior Wellness Center. This sample size was sufficient and had 80% power to test the primary hypothesis based on a multiple logistic regression analysis assuming  $\alpha = 0.05$ ; an assumed odds ratio and  $R^2$  of 2.96 and 0.20, respectively. These assumptions were based on previous studies that have utilized the HBM model.

### 2.3. Data collection

The interviews were conducted using ACASI with an instrument developed specifically for this study. The instrument used in this study was created from previously used questionnaires by Poss et al. [12], which provided items for assessment of HBM/TRA constructs and from Lindsay et al. [13] and Proeschold et al. [14], who offered items relevant for the evaluation of HCV knowledge. Items to collect sociodemographic and prior HCV testing history were also added. The survey items were modified to suit our study population and subject matter and retested for reliability in a separate pilot study before the start of this study. After informed consent had been provided, data collection was done in a designated area at the recruitment sites under the oversight of study personnel.

### 2.4. Statistical analysis

Descriptive statistics were estimated for the primary outcome HCV testing intention within 6 months, along with other study variables (i.e., demographics, prior testing history, HBM and TRA constructs, and HCV knowledge). Means and standard deviations were used for continuous variables while frequencies and proportions were used for categorical variables. A logistic regression analysis was conducted to examine factors that predict willingness to test for HCV within 6 months. Both simple and multiple logistic regressions were performed. Covariate selection into the final multivariable model was based on whether the variable met the criteria of  $p < 0.2$  in the simple logistic regression or whether it was of theoretical/clinical importance in the study. The variables assessed included HBM constructs, TRA construct of subjective norms, age, marital status, income, gender, education, previous HCV testing, health insurance, having a blood transfusion prior to 1992 and HCV knowledge. Adjusted and unadjusted odds ratios (ORs) with their 95% confidence intervals (CIs) were estimated and reported. Based on literature findings, interaction testing was also conducted to evaluate possible effect modification of the effect of perceived susceptibility (an HBM construct) and subjective norms (TRA

construct). All analyses were performed using SPSS version 22 at an  $\alpha$  value of 0.05.

## 3. Results

### 3.1. Patient characteristics

Of the 137 participants, 60.6% were female, and 44.8% had at least a college education. The mean age of respondents was 58.85 years. An estimated 43.8% of AABBs had a previous HCV test; 11 of whom were HCV positive while 32.1% of respondents had the intention to test for HCV within 6 months. About 68% did not consider themselves susceptible to HCV as AABBs and 72.3% as BBs. Other characteristics are summarized in Table 1.

### 3.2. Predictors of intention to test for HCV within 6 months

Table 2 shows results of the logistic regression analysis modeling intention to test for HCV within 6 months. Findings from the adjusted analysis showed that age, prior blood transfusion, subjective norms, perceived severity, and perceived benefits were statistically significant adjusted predictors of intention to test for HCV within 6 months. The model  $\chi^2$  was 54.24 (degrees of freedom = 10,  $p \leq 0.0001$ ), and the Hosmer–Lemeshow test was not significant ( $p = 0.347$ ) indicating good model fit. Nagelkerke  $R^2$  was 0.329, which means that 32.9% variability in intention to test for HCV within 6 months is explained by the variables included in the model.

Table 2 shows that age had an inverse relationship with intention to test for HCV within 6 months. A 1-year increase in age was associated with a decrease in the odds of having an HCV test

**Table 1**  
Sociodemographic and health-related characteristics of respondents.

Sociodemographic characteristics	Findings
Age	58.85 ± 6.38
HCV knowledge score	48.74 ± 26.24%
Gender	
Male	54 (39.40)
Female	83 (60.60)
Education	
No school/grades 1–11	16 (11.80)
High school	59 (43.40)
College degree	35 (25.70)
Graduate degree	26 (19.10)
Prior to 1992 received a blood transfusion or blood products	
Yes	19 (13.90)
No/unsure	118 (86.20)
Susceptible to HCV as African American	
Yes	44 (32.10)
No	93 (67.90)
Susceptible to HCV as Baby Boomer	
Yes	38 (27.70)
No	99 (72.30)
Prior test for hepatitis C	
Yes	60 (43.80)
No	77 (56.20)
Test result (n = 60)	
Negative	49 (81.70)
Male	20 (40.80)
Female	29 (59.20)
Positive	11 (18.30)
Male	5 (45.50)
Female	6 (54.50)
Intention to test for HCV within next 6 months	
Yes	44 (32.10)
No	93 (67.90)

Data are presented as mean ± standard deviation or n (%).  
HCV = hepatitis C virus.

Download English Version:

<https://daneshyari.com/en/article/5662961>

Download Persian Version:

<https://daneshyari.com/article/5662961>

[Daneshyari.com](https://daneshyari.com)