



Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

## Bilateral methachronous testicular germ cell tumor and testicular microlithiasis in a child: Genetic analysis and insights. A case report



N. Boudaoud<sup>a</sup>, G. Loron<sup>b</sup>, M. Pons<sup>a</sup>, E. Landais<sup>c</sup>, S. Kozal<sup>a</sup>, M. Doco-Fenzy<sup>c</sup>, M.L. Poli-Merol<sup>a,\*</sup>

<sup>a</sup> Department of Pediatric Surgery, American Memorial Hospital, CHU REIMS, France

<sup>b</sup> Department of Pediatrics, American Memorial Hospital, CHU REIMS, France

<sup>c</sup> Genetic Department, CHU-Reims, SFR-CAP Santé, UFR Médecine Reims, EA 3801, France

## ARTICLE INFO

## Article history:

Received 3 August 2017

Received in revised form

25 September 2017

Accepted 27 September 2017

## Keywords:

Testicular microlithiasis

Testicular germ cell tumor

Genetics

Children

CGH array

## ABSTRACT

**OBJECTIVES:** To report our experience with a case of a child with bilateral testicular micro-lithiasis (TML) who developed bilateral metachronous testicular germ cell tumor (TGCT) and determine the most appropriate follow-up and care management in children with testicular micro calcifications in regards to the theoretical risk of testicular cancer.

**CASE REPORT:** A 12 year-old boy was diagnosed with TGCT and TML. Ten years after complete remission, he presented with a recurrence on the contralateral testis. Genetic screening was performed on both resected and the patient's karyotype was analyzed.

**RESULTS:** Blood karyotype was normal. Aberrations were found in the tumor karyotype. CGH array showed alterations in chromosome arm 12p.

**DISCUSSION:** TML is frequently associated with testicular malignancy in adults: in 16.9% of cases the normal contralateral testicle develops TML in TGCT. Recent works of literature find no relationship between TML and cancer in general, but in patients with additional risks, the relationship becomes stronger. Some authors suggest that environmental components and genetics are determinant factors. This is highly suspected in our reported case. It would seem that TML is not a precancerous lesion per se, but rather a marker of an at-risk situation. Long term evolution is uncertain and regular self-palpation that starts before puberty is the only way to ensure proper screening and monitoring.

**CONCLUSION:** TML have been suspected to be a sign of testicular dysgenesis syndrome, which yields a risk of developing TGCT in case of noxious associations.

In patients with a history of TGCT contralateral TML is alarming and aggressive surgical management should be discussed. Therapeutic education of these patients on self-palpation is the best way to ensure proper follow-up.

© 2017 Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

The management of TML is still being discussed in the literature, especially in children. We report the case of a child who presented with TGCT associated with TML and a long term contralateral recurrence. Due to the rarity of the case, the association with TML, both tumor were explored genetically. As far as we know, this is the first case of a Yolk sac tumor reporting the results of karyotype and DNA oligo-array in tumors developed within an interval of 10 years, in both testis in a young adult presenting TML.

### 2. Case report

A 12 year-old boy presented to our clinic for testicular tumor discovered by the family physician during routine examination. The right testis was enlarged, firm, tender to the touch and irregular. The left testicle was normal. The boy presented with clinical symptoms of precocious puberty.

Ultrasound confirmed an enlarged testicle infiltrated by a vascularized, heterogeneous tumor (Fig. 1). TML were noticed both in the remaining right parenchyma and in the left testis. Preoperative laboratory tests showed increased Alpha-fetoprotein levels.  $\beta$ CGH and testosterone levels were also elevated due to peripheral secretion, responsible of the precocious puberty. The tumor was still localized and orchiectomy was performed, after initial control of testicular vessels, via an inguinal approach. Histopathology validated the ter-

\* Corresponding author at: Pediatric Surgery – AMH 47 Rue Cognac Jay, 51100 Reims, France.

E-mail address: [mpolimerol@chu-reims.fr](mailto:mpolimerol@chu-reims.fr) (M.L. Poli-Merol).




Fig. 1. Left testis US at 12 year-old.

atocarcinoma. Postoperative recovery was uneventful and biology lab results were totally normalized after 3 weeks of follow-up.

The child was then regularly assessed, every three months then twice a year, and finally once a year until he reached the age of 18. During this 6-year follow-up, ultrasound examinations of the remaining testis and empty scrotum were normal except for previously evidenced calcifications in the left testicle. Self-palpation was strongly recommended and the systematic follow-up ended.

Ten years later, during a routine examination again, his family physician noticed a change in his left testicle. The patient admitted he had noticed the change 3 months before. Additional examinations confirmed the recurrence with mediastinal and pulmonary metastases. Alpha-fetoprotein levels were increased. A left orchiectomy was performed after sperm storage. Adapted chemotherapy was done with a good response. Lung metastases were removed and the patient is now recurrence-free after a 62-month follow up.

Due to the rarity of the case, we retrospectively explored both tumors and a sample of the patient's blood via karyotype and CGH array. Patient consent was mandatory and obtained.

Tumour samples from both sides were enzymatically dissociated; cells were cultured on ECM coated flask surfaces and fed with an enriched culture medium. The cells did grow slowly. After about 20 days the monolayer cultures were trypsinised and cells were karyotyped after banding.

### 3. Array-CGH

Genomic DNA samples were extracted from peripheral blood using the QIAamp DNA Blood Midi kit (Qiagen, Valencia, CA, USA) and from paraffin embedded slides using the QIAamp DNA Tissue Midi kit (Qiagen). Array-CGH was performed in all samples using the Agilent 44k oligoarrays (Human Genome CGH Microarray Kit, Agilent Technologies, Santa Clara, CA, USA). Random primer labelling and hybridization were carried out with sex-matched reference DNA according to the manufacturer's recommendations. Images were acquired using an Agilent scanner. Data were processed with Cytogenomics software (Agilent Technologies), and results were analysed with the CGH Analytics software (v3.5.14; Agilent Technologies) using the ADM2 algorithm and a three-point filter.

Results are reported in Table 1.

#### 3.1. Interpretation

Unfortunately, the M-FISH could not be realised to better characterise the chromosomes breakpoints, but instead the array-CGH showed the global gains and losses of the chromosomes. With this analysis, less rearrangements were detected than expected consid-

ering the complex structural abnormalities on the Karyotype. The micro-array results confirmed and defined the size of the *i*(12p) observed on both tumours, it showed the quantitative rearrangements present in both tumours and not in the peripheral blood of the patient.

### 4. Discussion

TML have recently been increasingly documented in the literature especially regarding the potential relationship between TML and TGCT, and its impact on the follow-up. TML are usually asymptomatic and detected during routine ultrasound examination of the testicles [1–6]. Radiologists established TML definition by the presence of at least five small non-shadowing hyperechoic foci ranging from 1 to 3 mm in diameter. Three stages have been described depending on the number of calcifications, without any clinical validated significance regarding evolution toward regression or tumor development [4,6,7,8].

Different studies on TML prevalence reported that 2.4–5.6% of asymptomatic adults and 1.1–4.2% of asymptomatic children present with TML [5–11].

TML are present in 45% of malignant testicular tumors in adults and in 14–16.9% of the normal testicle in case of contralateral malignant tumor and some authors report a high testicular recurrence rate in cases of previous germ-cell tumor associated with ML in the contralateral testicle, with an odd-ratio ranging between 12 and 17 [12,13].

In 2010, a meta-analysis reported no causal link between TML and cancer in the absence of additional risk factors. Yet, in the presence of risk factors, TML was associated with a substantially elevated risk of TGCT [14].

TML are supposed to originate from Seminiferous tubules degeneration, with intratubular calcium deposits, showing concentrically laminations containing DNA, phospholipid and protein scraps [1,4,10,12]. Drut suggested the role of an early disruption in gonad development leading to Sertoli cells dysfunction [15]. From 2004, some authors brought up the hypothesis that TML was a feature of “testicular dysgenesis syndrome” (TDS) and that genetics plays a key role in the onset of TML [16–21].

Moreover, recent studies still underlines the statistically significant augmentation of TGCT in TML population, either in children or adults, even if they do not retain TML as a risk factor per se [22,23].

According to Skotheim, TGCT of adolescent and young adults develop from precursor lesions or carcinoma in situ (CIS) believed to originate from diploid primordial germ cells during foetal life [21]. The procedure starts with aneuploidisation and extensive chromosome instability [21,24,25]. The transformation into invasive TGCT is associated with increased copy number of chromosome arm 12p mostly *i*(12p), indeed 12p gain is the most consistent chromosomal abnormality in human testicular cell tumours [26,27]. TGCT karyotypes are characterized by hyperdiploidy and frequently in the triploid range. This is the case in our patient with the hypotriploid modal number of the chromosomes in the first culture. Interestingly when the cells were cultured during a few more days, the karyotype of the abnormal cells became triploid. We wonder if the development of the second tumour is dependent or independent of the first one. Do the cells of both testicular tumours originate from the same clone? We showed that the second tumour shares the 12p amplification with the first tumour but has less abnormal gain or losses. Is the acquisition of the 12p amplification (or *i*(12p)) in the second tumour due to a secondary clone deriving from the first abnormal embryonic germ cells or is it depending only on the predisposition factors or genes giving “life” to a second independent tumour clone?

Download English Version:

<https://daneshyari.com/en/article/5732467>

Download Persian Version:

<https://daneshyari.com/article/5732467>

[Daneshyari.com](https://daneshyari.com)