



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Long-term observation of a large keratocystic odontogenic tumour of the mandible treated by a single enucleation procedure: A case report and literature review

Tobias L. Pittl^a, Marius Meier^a, Paul Hakl^b, Walter Sutter^b, Dritan Turhani^{b,*}^a Centre of Oral and Maxillofacial Surgery, Faculty of Dentistry, Danube Private University, Steiner Landstraße 124, A-3500 Krems-Stein, Austria^b Centre of Oral and Maxillofacial Surgery, Danube Private University, Steiner Landstraße 124, A-3500 Krems-Stein, Austria

ARTICLE INFO

Article history:

Received 15 December 2016

Received in revised form 21 March 2017

Accepted 21 March 2017

Available online 23 March 2017

Keywords:

Enucleation

Keratocystic odontogenic tumour

Odontogenic keratocyst

Recurrence

Surgery

ABSTRACT

INTRODUCTION: Keratocystic odontogenic tumours (KCOTs) are benign lesions of the jaw that are characterised by expansive growth and high rates of recurrence. Herein, we present a novel minimally invasive method for the surgical treatment of KCOTs.

CASE PRESENTATION: We present a 49-year-old woman with a rare, large KCOT of the mandible extending from tooth 37 to tooth 47. A single enucleation procedure was selected as the surgical technique combined with a titanium plate. Teeth interfering with the cystic lumen were preserved. The bone surrounding the cyst was partially removed and rinsed with Carnoy's solution. Recurrence was observed 1 year later between teeth 43 and 45, and was treated by single enucleation with Carnoy's solution. After the second operation, the teeth interfering with the KCOT still elicited a positive response to dental pulp testing. No further signs of recurrence were observed after a total observation period of 7 years.

DISCUSSION: Various surgical interventions have been described for KCOTs. However, a general guideline for a specific surgical intervention has not yet been established owing to the heterogeneity of these tumours.

CONCLUSION: Immediate enucleation in combination with a titanium plate offers an effective surgical approach for the treatment of large KCOTs of the mandible without the need for radical tooth removal or resection of the afflicted side.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Keratocystic odontogenic tumours (KCOTs) were first described in 1956 as benign neoplasms of the jaw that have a high risk of recurrence [1]. Considering their aggressive infiltrative behaviour that often leads to penetration of the surrounding tissues, KCOTs exhibit the combined characteristics of cysts and benign tumours [1,2]. KCOTs are usually characterised by an 8–10 cell thick parakeratinised epithelium and represent 12–14% of all odontogenic cysts of the jaw [1,3]. Clinical symptoms of odontogenic keratocysts are rarely observed; they are often discovered incidentally through radiographic examination [4]. KCOTs generally present as

well-defined radiolucent lesions with a radiopaque cortical margin [5].

In most instances, when the KCOT grows to a large size, marsupialisation or decompression with subsequent enucleation is performed to reduce the size of the tumour and preserve adjacent structures [6].

Herein, we present a case involving a large KCOT of the mandible treated by enucleation with Carnoy's solution (CS), peripheral ostectomy, and a 2.0-mm 16-hole titanium plate. Teeth interfering with the cystic lesion were not extracted, but still elicited a positive response to dental pulp testing after the operation. This case report is compliant with the SCARE guidelines [7].

2. Case presentation

In May 2009, a 42-year-old woman developed a sudden, painless swelling in the posterior region of the right side of her mandible that continued for approximately 14 days. The patient had experienced neither pathological mobility of the teeth nor malocclusion. Her medical and family histories were unremarkable. A panoramic radiographic (PR) examination had not been previously performed. Subsequent PR imaging revealed a large, radiolucent, expanding

Abbreviations: CT, computed tomography; CS, Carnoy's solution; KCOT, keratocystic odontogenic tumour; PR, panoramic radiograph.

* Corresponding author.

E-mail addresses: tobias.pittl@gmail.com (T.L. Pittl), meier_marius.1@hotmail.com (M. Meier), paul.hakl@dp-uni.ac.at (P. Hakl), walter.sutter@dp-uni.ac.at (W. Sutter), dritan.turhani@dp-uni.ac.at, dritan.turhani@gmail.com (D. Turhani).

<http://dx.doi.org/10.1016/j.ijscr.2017.03.033>

2210-2612/© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Pre-surgical panoramic radiographic image (2009).




Fig. 2. Intra-oral view of the cystic lesion during surgery with perforation of the cortical bone (2009).

lesion involving the majority of the mandibular bone from the distal side of tooth 35 to the distal side of tooth 45 with no evidence of root resorption (Fig. 1). An axial computed tomography (CT) scan revealed a well-defined multilocular cystic lesion. On the right side, a perforation in the cortical region was examined (approximately 10 mm in diameter) with the overlying soft tissue. Biopsy findings were suggestive of a KCOT.

Surgery was performed under nasotracheal intubation. A mucoperiosteal flap extending from teeth 37–47 was prepared. Exceedingly thin layers of compact bone covering the cystic lumen were removed (Fig. 2). The cyst was enucleated by removing the surrounding bone. Owing to the large size of the KCOT, and to reduce the risk of a mandibular fracture, a 2.0-mm titanium plate (MatrixMANDIBLE Plating System[®]; Synthes CMF, West Chester, PA, USA), fixed with 3- × 10-mm screws on each side, was installed (Fig. 3). The cystic defect and the area surrounding the roots of the affected teeth were carefully swabbed with CS (60% ethanol, 30% chloroform, and 10% glacial acetic acid) as a curative agent for only 1 min to avoid damaging the nerves around the teeth, followed by irrigation with sodium chloride. The defect was then closed using a 5-0 PROLENE[®] polypropylene suture. Histological analysis of the specimen confirmed a final diagnosis of a parakeratinised KCOT.

After the initial surgery, the patient underwent clinical examinations every 3 months and PR and CT examinations every 6 months. In August 2010, a recurrence of the KCOT was suspected based on PR and CT findings. A CT scan revealed three radiolucent regions in the mandibular bone extending from the mesial side of tooth 43 to the distal side of tooth 45 (Fig. 4). A second operation was performed under nasotracheal intubation. Three cystic lesions were enucleated, the surrounding bone was partially removed, and CS was applied for 1 min followed by irrigation with sodium




Fig. 3. Intra-operative placement of the 2.0 mm 16-hole titanium plate (MatrixMANDIBLE Plating System[®]; Synthes CMF, West Chester, PA, USA) (2009).



Fig. 4. Pre-surgical computed tomography image showing three recurrent cystic lesions (2010).

chloride, as in the first operation. Histological analysis of the specimen confirmed a diagnosis of a recurrent KCOT. No complications occurred during surgery, and post-surgical healing was satisfactory.

Clinical follow-up examinations with PR and CT scans were performed for early detection of any recurrence every 6 months until March 2016; no signs of recurrence were observed (Fig. 5).

3. Discussion

KCOTs are benign lesions of the jaw that frequently expand into the medullary cavity in the anteroposterior direction without perforating the solid bone [8,9]. Rapid growth of KCOTs may occur because of an active epithelial lining with a high rate of proliferation [5]. In our patient, the KCOT was located in the posterior region of the mandible on each side, as well as in the symphysis and premolar regions. Such KCOTs were reported in only 1.8% of 107 cases in a previous study [10]. Our patient noticed only a slight swelling

Download English Version:

<https://daneshyari.com/en/article/5732841>

Download Persian Version:

<https://daneshyari.com/article/5732841>

[Daneshyari.com](https://daneshyari.com)