


Accepted Manuscript

Title: Mitigation of postnatal ethanol-induced neuroinflammation ameliorates trace fear memory deficits in juvenile rats

Authors: Molly J. Goodfellow, Youn Ju Shin, Derick H. Lindquist



PII: S0166-4328(17)31235-4
DOI: <https://doi.org/10.1016/j.bbr.2017.09.047>
Reference: BBR 11116

To appear in: *Behavioural Brain Research*

Received date: 27-7-2017
Revised date: 22-9-2017
Accepted date: 27-9-2017

Please cite this article as: Goodfellow Molly J, Shin Youn Ju, Lindquist Derick H. Mitigation of postnatal ethanol-induced neuroinflammation ameliorates trace fear memory deficits in juvenile rats. *Behavioural Brain Research* <https://doi.org/10.1016/j.bbr.2017.09.047>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mitigation of postnatal ethanol-induced neuroinflammation ameliorates trace fear memory deficits in juvenile rats

Molly J. Goodfellow^{a, 1}, Youn Ju Shin^b, and Derick H. Lindquist^{a, b}

Departments of Psychology^a and Neuroscience^b

The Ohio State University | 1835 Neil Ave. Columbus, OH 43210, USA

Emails: mgoodfellow@som.umaryland.edu, shin.467@osu.edu, lindquist.40@osu.edu

Pages: 10

Tables: 1

Figures: 1

Abstract words: 184

Manuscript words: 2574

Total Words (minus Title page): 3418

Corresponding Author:

Dr. Derick H. Lindquist

Departments of Psychology & Neuroscience

The Ohio State University

1835 Neil Avenue, Room 49

Columbus, OH 43210

Email: lindquist.40@osu.edu

Phone: 614-292-2236

Fax: 614-688-4733

Abstract

Impairments in behavior and cognition are common in individuals diagnosed with fetal alcohol spectrum disorders (FASD). In this study, FASD model rats were intragastrically intubated with ethanol (5 g/kg/day; 5E), sham-intubated (SI), or maintained as naïve controls (NC) over

¹ **Present address:** Department of Anesthesiology and the Center for Shock, Trauma, and Anesthesiology Research (S.T.A.R.) | University of Maryland School of Medicine | MSTF 5-54 685 W. Baltimore St. Baltimore, MD 21201, USA

Download English Version:

<https://daneshyari.com/en/article/5735175>

Download Persian Version:

<https://daneshyari.com/article/5735175>

[Daneshyari.com](https://daneshyari.com)