




Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa



# Harmonic mappings with hereditary starlikeness

Ngin-Tee Koh

School of Mathematics and Statistics, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

## ARTICLE INFO

*Article history:*  
 Received 6 February 2017  
 Available online xxxx  
 Submitted by R. Gornet

*Keywords:*  
 Annulus  
 Doubly-connected  
 Harmonic mappings  
 Hereditary property  
 Starlike

## ABSTRACT

We study a hereditary starlikeness property for planar harmonic mappings on a disk and on an annulus. While such a property is a common trait of conformal mappings, it may be absent in harmonic mappings. It turns out that a sufficient condition for a harmonic mapping  $f$  to possess this hereditary property is to have a harmonic argument — a striking feature of conformal mappings that does not extend to all harmonic mappings.

© 2017 Elsevier Inc. All rights reserved.

## 1. Introduction

Harmonic mappings, which are complex-valued sense-preserving one-to-one functions satisfying Laplace’s equation  $\Delta f = 0$  on their respective domains in  $\mathbb{C}$ , possess some interesting properties. For instance, while it follows from a sharp result of Heinz [4, Lemma] that Euclidean and hyperbolic distances are not necessarily shortened by harmonic mappings of hyperbolic regions (see, e.g., [3, page 77] or [11, page 91]), the Lebesgue area measure of concentric disks

$$\mathbb{D}_r = \{z \in \mathbb{C} : |z| \leq r < 1\}$$

is reduced by harmonic mappings preserving the unit disk  $\mathbb{D}$  [11, Theorem 1.1].

If the image of the unit disk under a conformal mapping is a starlike region  $\Omega$ , then the image of every disk in  $\mathbb{D}$  is also starlike (see, e.g., [2, proof of Theorem 2.10]). This hereditary starlikeness property need not hold for harmonic mappings. For example, the harmonic mapping

$$f(z) = \operatorname{Re} \frac{z}{1-z} + i \operatorname{Im} \frac{z}{(1-z)^2}$$

E-mail address: [ngin-tee.koh@canterbury.ac.nz](mailto:ngin-tee.koh@canterbury.ac.nz).

<http://dx.doi.org/10.1016/j.jmaa.2017.08.032>  
 0022-247X/© 2017 Elsevier Inc. All rights reserved.

maps  $\mathbb{D}$  onto the half plane  $\{w : w > -\frac{1}{2}\}$ , which is starlike, but  $f(\mathbb{D}_r)$  is not starlike for  $\sqrt{\frac{7\sqrt{7}-17}{2}} < r < 1$  [13, Example 1.1]. Nonetheless, we will provide sufficient conditions for harmonic mappings to possess this hereditary property. We will also establish analogous results for harmonic mappings between doubly-connected regions, which is the primary focus of this paper.

**2. Preliminaries and main results**

For  $0 < \rho \leq r < 1$ , let  $\mathbb{A}_\rho$  denote the annulus  $\{z \in \mathbb{C} : \rho < |z| < 1\}$ , let  $\overline{\mathbb{A}_\rho}$  be its closure, and let  $\mathbb{T}_r$  represent the circle  $\{z \in \mathbb{C} : |z| = r\}$ . We will use  $\mathbb{T}$  to denote the unit circle  $\partial\mathbb{D}$ . A *curve* is a continuous image of the interval  $[0, 1]$ .

*2.1. Starlikeness of simply and doubly connected regions*

Let  $f$  be a harmonic mapping of  $D$ , where  $D$  is either  $\mathbb{D}$  or  $\mathbb{A}_\rho$ . In the latter situation, we may assume without loss of generality that the inner and outer boundaries of  $D$  are mapped respectively to the inner and outer boundaries of  $f(D)$ , and it should be noted that the outer boundary of  $f(D)$  necessarily contains points in  $\mathbb{C}$  [7, Section 4.1]. Then the analytic characterization for  $f(\mathbb{T}_r)$ , where  $0 < \rho < r < 1$ , to enclose a starlike region  $S$  is the existence of a point  $a \in S$  such that

$$\frac{\partial}{\partial t} \arg[f(re^{it}) - a] \geq 0. \tag{1}$$

Geometrically,  $\arg[f(re^{it}) - a]$  increases as  $\mathbb{T}_r$  is traced counterclockwise, and any ray emanating from  $a$  intersects  $f(D)$  in a single, possibly infinite, line segment. It is standard terminology to say that  $S$  is starlike with respect to  $a$ . In particular,  $S$  is strictly starlike with respect to  $a$  if the inequality in (1) is strict. In geometrical terms, strict starlikeness with respect to  $a$  means that no tangent line to the boundary of  $S$  contains  $a$ . A normalization may be achieved by applying a translation to  $S$  (or  $f(D)$ ) so that  $a = 0$ , and we will define a curve to be (*strictly*) *starlike* if it forms the boundary of a region that is (strictly) starlike with respect to the origin. A *doubly-connected starlike region* is one whose intersection with any ray from the origin is either a single line segment or a single ray. If  $D = \mathbb{D}$ , then the representation  $f = h + \bar{g}$  for some holomorphic functions  $g$  and  $h$  allows one to rewrite (1) when  $a = 0$  as (see [1, page 139])

$$|h(z)|^2 \operatorname{Re} \frac{zh'(z)}{h(z)} \geq |g(z)|^2 \operatorname{Re} \frac{zg'(z)}{g(z)} + \operatorname{Re}[z(h(z)g'(z) - h'(z)g(z))].$$

Nevertheless, we will work directly with (1). Our first result is as follows.

**Theorem 2.1.** *Suppose  $f$  is a harmonic mapping of  $\mathbb{D}$  onto a starlike region  $\Omega_0 \subset \mathbb{C}$ . Assume that on  $\mathbb{A}_{\sqrt{2}-1}$ ,*

$$\Delta \operatorname{Im} \log f = 0, \tag{2}$$

*where  $\Delta$  represents the Laplace operator. Then  $f(\mathbb{D}_r)$  is a strictly starlike region for  $0 < r < 1$ .*

**Remark 2.2.** The boundary of a starlike region need not be a curve. For instance, the set  $\{(2 + \sin \frac{1}{t}) e^{it} : t \in (0, 2\pi]\} \cup [1, 3]$  is the boundary of a starlike region, but is not locally connected, and is therefore not a curve by the Hahn–Mazurkiewicz theorem (see, e.g., [14, page 89]).

Given a harmonic mapping  $f$  of  $\mathbb{D}$ , where  $f(\mathbb{D})$  is a starlike region with respect to the origin, it is known that  $f(\mathbb{D}_r)$  is starlike with respect to the origin for at least  $0 < r \leq \sqrt{2}-1$  (see, e.g., [13, Theorem 2.16(iii)]). This explains the restriction on (2) in Theorem 2.1 to the annulus  $\mathbb{A}_{\sqrt{2}-1}$ . It follows that Theorem 2.1 is a consequence of the more general result below.

Download English Version:

<https://daneshyari.com/en/article/5774663>

Download Persian Version:

<https://daneshyari.com/article/5774663>

[Daneshyari.com](https://daneshyari.com)