

Incidence of Sports-Related Concussion among Youth Football Players Aged 8-12 Years

Anthony P. Kontos, PhD¹, R. J. Elbin, PhD¹, Vanessa C. Fazio-Sumrock, PhD¹, Scott Burkhart, PsyD¹,
Hasani Swindell, BS², Joseph Maroon, MD³, and Michael W. Collins, PhD¹

Objective To determine the risk of concussion among youth football players (ages 8-12 years).

Study design Participants included 468 male youth football players in western Pennsylvania during the 2011 youth football season. Incidence rates (IRs) and incidence density ratios (IDRs) of concussion were calculated for games and practices and for age groups.

Results There was a total of 11 338 (8415 practice and 2923 game) athletic exposures (AEs) in the study period, during which 20 medically diagnosed concussions occurred. A majority of concussions were the result of head-to-head (45%) contact. The combined concussion IR for practices and games was 1.76 per 1000 AEs (95% CI 0.99-2.54). The concussion IR was 0.24 per 1000 AEs (95% CI 0.04-0.79) in practices and 6.16 per 1000 AEs (95% CI 3.76-9.54) in games. The IDR for concussions in games to practices was 25.91 (95% CI 6.01-111.70). The IDR for concussions for youth aged 11-12 years compared with youth aged 8-10 years was 2.72 (95% CI 0.66-4.78).

Conclusions The overall IR for concussion in youth football players aged 8-12 years was comparable with that reported previously for high school and collegiate samples. However, participation in games was associated with an increase in risk of concussion compared with practices, which was higher than rates previously reported for high school and collegiate athletes. Younger players were slightly less likely to incur a concussion than were older players. (*J Pediatr* 2013;163:717-20).

Sports-related concussion is referred to as an “epidemic” by the Centers for Disease Control and Prevention (CDC).¹ Emergency department visits for concussions increased 62% between 2001 and 2009,² and among 8- to 13-year-olds, the number of emergency department visits nearly doubled. Collision sports such as American football are purported to have high rates of concussion. Recent studies estimate that concussions comprise between 8.9%³ and 13.2%⁴ of all high school injuries and 7.9%⁵ of all collegiate sport injuries.

High school football players have the highest likelihood (0.47 per 1000 athletic exposures [AEs]) of sustaining a concussion in the High School Reporting Information Online database.³ There are, however, no published prospective studies on the prevalence and incidence rates (IRs) of concussion in youth football for children aged 8-12 years. Despite the lack of data, there is no shortage of claims that youth football at these ages is dangerous for children. These claims have begun to shape public opinion about whether children should be allowed to play youth football and, if so, how the game should be played or changed from its current form. In fact, Pop Warner, the largest national youth football organization in the US, recently limited youth football contact time to reduce the likelihood of concussion.⁶

In the US, there are approximately 3 million youth participating in tackle football compared with 2000 professional, 100 000 collegiate, and 1.3 million high school participants.⁷ Organized youth tackle football leagues such as Pop Warner report that approximately 425 000 athletes ranging from 5 to 16 years are involved in their tackle football programs.⁶ The IRs for concussion in high school football range from 0.47 to 0.60 concussion per 1000 AEs,^{3,8} whereas IRs for college football range from 0.43 to 0.61 concussion per 1000 AEs.³ Recent reports indicate that concussion IRs are on the rise for high school football. Lincoln et al⁸ reported an 8% increase for concussions in high school football over a 20-year period from 1997-1998 to 2007-2008.

An empirical investigation of concussion incidence in this younger population is warranted because younger athletes demonstrate worse outcomes following concussion^{9,10} and are at a high risk of catastrophic injury (ie, second impact syndrome) if they continue to play with a symptomatic concussion.¹⁰⁻¹² It has even been speculated that playing youth tackle football may lead to residual impairment; however, no empirical evidence supports this assertion. The purpose of the current study was to investigate concussion incidence in youth tackle football

From the ¹Department of Orthopaedic Surgery/ University of Pittsburgh Medical Center Sports Medicine Concussion Program, University of Pittsburgh, Pittsburgh, PA; ²Columbia University College of Physicians and Surgeons, New York, NY; and ³Department of Neurosurgery, University of Pittsburgh, Pittsburgh, PA

Funded by National Football League Charities Medical Research grant. The authors declare no conflicts of interest.

0022-3476/\$ - see front matter. Copyright © 2013 Mosby Inc. All rights reserved. <http://dx.doi.org/10.1016/j.jpeds.2013.04.011>

AE	Athletic exposure
CDC	Centers for Disease Control and Prevention
IDR	Incidence density ratio
IR	Incidence rate

participants aged 8-12 years. Specifically, we prospectively investigated the overall IR of sport-related concussion in youth tackle football games and practices and compared the rates of players aged 8-10 years with those of players aged 11-12 years. In addition, we compared the current study's IRs with published IRs for high school and college athletes.

Methods

A prospective cohort study was conducted between August 2011 and December 2011 on 468 of a total of 571 (82% participation rate) participants aged 8-12 years old (mean \pm SD 10.12 \pm 1.31 years) from 4 nonscholastic youth tackle football leagues composed of 18 teams. Inclusion criteria included age of 8-12 years, written informed consent (parent) and assent (child), players participating only in the sport of football (during the study period), and league and coach agreement to participate. Exclusion criteria included a history of any of the following: brain surgery, traumatic brain injury, or neurologic or psychiatric disorder and/or current concussion or a concussion in the past 6 months.

According to recent CDC guidelines¹ and the 3rd International Consensus Statement on Concussion in Sports,¹³ a concussion was defined as any mild closed head injury involving altered cognitive functioning (eg, confusion, memory loss, disorientation) or signs or symptoms (eg, headache, dizziness, balance problems, nausea) or brief loss of consciousness of no longer than 1 minute following a direct or indirect blow to the head. Head injuries involving any structural damage or abnormality (eg, skull fracture, subdural hematoma, other lesions) were excluded from the research study.

The University of Pittsburgh institutional review board approved the protocol and consent forms before data collection. Following a brief overview of the risks and benefits of the study, informed written parental consent was obtained from each participant's parent/guardian, and assent was obtained from each participant. Throughout the study period, a designated and trained representative for each team recorded practice and game exposures for their players. Player exposures were only recorded if a player was present and actively participated in a game or practice. A member of the research team was present on the sidelines during games to record injury mechanism data for any suspected concussion. Licensed medical professionals (eg, emergency medical technicians, physicians) assessed and managed athletes with suspected concussions during games. However, medical coverage was limited during practices. Consequently, we contacted coaches 2 or 3 times per week to determine if any suspected concussions occurred during practices. To assist in this process, the coaches in the study participated in preseason concussion education and data collection training sessions. In addition, players and parents were provided with concussion education training. For each suspected concussion, we gathered information by structured phone and in-person interviews from all participants (and their parents) and recorded additional information regarding the

mechanism of injury (ie, helmet-to-helmet contact, helmet-to-ground contact). Only participants with suspected concussions that were subsequently confirmed by a licensed medical professional, as per the definition, were recorded as concussed. Confirmation was in the form of a written physician's note, and players with a diagnosed concussion were removed from play as per their league's concussion policy.

Statistical Analyses

Player exposure data were tabulated for each participant and used to calculate overall IRs per 1000 AEs with 95% CIs for both practices and games. Incidence density ratios (IDRs) with 95% CIs comparing the rates of concussion in games with that practice and between the 8- to 10-year-old and the 11- to 12-year-old age groups were also calculated. A χ^2 analysis with OR was used to compare the odds of a concussion for 11- to 12-year-olds compared with 8- to 10-year-olds. A statistical significance level of $P \leq .05$ and 2-sided tests were used for all analyses in this study. SPSS version 20 (SPSS Inc, Chicago, Illinois) was used for all analyses.

Results

Four male youth tackle football leagues consisting of 22 teams were asked to participate in the study. A total of 468 of 571 possible participants (82.0%) aged 8-12 years old (mean \pm SD 10.12 \pm 1.31 years) comprising 18 teams participated in the study. Four of the 22 teams declined to participate in the study due to perceived time requirements or disinterest in the study. The remaining 18 teams had full participation from their eligible and consenting players. A summary of the demographic data is presented in **Table I**. Reasons for nonparticipation at the team level included testing/scheduling conflicts and coach refusal to participate. Reasons for nonparticipation at the individual level included meeting ≥ 1 exclusion criteria, refusal of parent to sign consent form, and inability to attend informational meetings or testing sessions.

Concussions in Practices and Games

There was a total of 11 338 (8415 practice and 2923 game) AEs in the study period during which 20 medically diagnosed concussions involving 20 different participants occurred. There were 2 concussions that occurred in practice and 18 concussions occurred during games. Identified mechanisms of concussions consisted of 45% head-to-head, 5% head-to-ground, and 5% head-to-body contact. The mechanisms

Table I. Summary of demographic information for 8- to 10-year-old and 11- to 12-year-old participants

Characteristic	Participants aged 8-10 y	Participants aged 11-12 y
Age, mean (SD) y	9.28 (0.74)	11.50 (0.71)
Height, mean (SD) cm	138.08 (9.24)	156.21 (12.57)
Weight, mean (SD) kg	36.04 (7.68)	55.79 (22.45)
No. of previous concussions, mean (SD)	0.06 (0.24)	0.50 (0.71)

Download English Version:

<https://daneshyari.com/en/article/6222718>

Download Persian Version:

<https://daneshyari.com/article/6222718>

[Daneshyari.com](https://daneshyari.com)