



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)



### Data Article

# Data on a delivery of biomolecules into *Nicotiana benthamiana* leaves using different nanoparticles

Antonida V. Makhotenko<sup>a,b</sup>, Ekaterina A. Snigir<sup>a</sup>,  
Natalia O. Kalinina<sup>a,b</sup>, Valentin V. Makarov<sup>a,b,\*</sup>,  
Michael E. Taliansky<sup>a,c</sup>

<sup>a</sup> DokaGene Ltd., Rogachevo, Moscow Region, Russia

<sup>b</sup> Lomonosov Moscow State University, Leninsky Gory, Moscow 119992, Russia

<sup>c</sup> James Hutton Institute, Invergowrie, DD2 5DA Dundee, UK

#### ARTICLE INFO

##### Article history:

Received 3 October 2017

Received in revised form

3 December 2017

Accepted 12 December 2017

Available online 21 December 2017

#### ABSTRACT

Nanoparticles (NPs) have a number of unique properties associated with their ultrasmall size and exhibit many advantages compared with existing plant biotechnology platforms for delivery of proteins, RNA and DNA of various sizes into the plant cells (Arruda et al., 2015; Silva et al., 2010; Martin-Ortigosa et al., 2014; Mitter et al., 2017) [1–4]. The data presented in this article demonstrate a delivery of biomolecules into *Nicotiana benthamiana* plant leaves using various types of NPs including gold, iron oxide and chitosan NPs and methods of biolistic bombardment and infiltration. The data demonstrate physical characteristics of NPs coated with fluorescently labeled protein and small RNA (size and zeta-potential) and visualization of nanocomplexes delivery into cells of *N. benthamiana* leaves by fluorescence microscopy.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

\* Corresponding author at: DokaGene Ltd., Rogachevo, Moscow Region, Russia.

E-mail address: [makarovvalentine@gmail.com](mailto:makarovvalentine@gmail.com) (V.V. Makarov).

## Specifications Table

Subject area	<i>Biotechnology</i>
More specific subject area	<i>Plant biotechnology</i>
Type of data	<i>Table and images</i>
How data was acquired	<i>Dynamic laser light scattering (DLS) and zeta-potential data on characterization of NPs were obtained on Malvern Zetasizer instrument - NanoZS, Malvern Instruments, UK. Fluorescence microscope Axiovert 200 M, Carl Zeiss, Germany was used for images of fluorescently labeled biomolecules in plant cells.</i>
Data format	<i>Analyzed data</i>
Experimental factors	<i>Fluorescence labeling of protein (bovine serum albumin with fluorescein isothiocyanate; BSA-FITC) and small RNA (tRNA with cyanine 3; tRNA-Cy3), assembly of nanoplatfoms consisting of (gold, iron oxide and chitosan) NPs and fluorescently labeled biomolecules, delivery of nanocomplexes into <i>N. benthamiana</i> leaves by biolistic bombardment (for gold and iron oxide NPs) and syringe infiltration (for chitosan NPs)</i>
Experimental features	<i>Characterization of NPs coated with biomolecules (size and zeta potential), visualization of labeled biomolecules delivered into cells of <i>N. benthamiana</i> leaves by fluorescence microscopy</i>
Data source location	<i>DokaGene Ltd., Rogachevo, Moscow Region, Russia, and Belozersky Institute of Physico-Chemical Biology, Lomonosov Moscow State University, Leninsky Gory, Moscow, 119992, Russia</i>
Data accessibility	<i>Data are provided with this article</i>

## Value of the data

- The data describe the assembly and characteristics of nanocomplexes consisting of different (gold, iron oxide and chitosan) NPs and fluorescently labeled protein (BSA-FITC) and small RNA (tRNA-Cy3).
- The data demonstrate delivery of the nanocomplexes into *N. benthamiana* leaves using two different techniques (biolistic bombardment and syringe infiltration) with subsequent visualization of fluorescently labeled biomolecules inside epidermis cells.
- The data may be used for development of easy and effective methods of biologically active component delivery to plant cells for different biotechnological applications.

## 1. Data

The data presented demonstrate loading capacity of the fluorescently labeled protein (BSA-FITC) and small RNA (tRNA-Cy3) on the surface of the gold, iron oxide and chitosan NPs (Fig. 1) and represent physical characteristics of assembled nanocomplexes built on gold and iron oxide NPs (Table 1, DLS and zeta-potential data). In Fig. 2, the images of plant cells were taken after delivery of nanocomplexes containing the fluorescently labeled BSA and tRNA into *N. benthamiana* leaves. *N. benthamiana* has been chosen as a model experimental plant highly convenient for plant biology and microscopy studies [5].

## 2. Experimental design, materials and methods

### 2.1. Nanocomplex assembly

We used commercial preparations of gold (1652262, Bio-Rad) and iron oxide (MP025-PEG-1ML, Nanocs) NPs. Chitosan NPs were prepared using tripolyphosphates (TPP), as described previously [6].

Download English Version:

<https://daneshyari.com/en/article/6597182>

Download Persian Version:

<https://daneshyari.com/article/6597182>

[Daneshyari.com](https://daneshyari.com)