


## Accepted Manuscript

Title: Psychosocial perspective and suicidal behaviors correlated with adolescent male smoking and illicit drug use

Authors: Hanan E. Badr, Konstantinos Francis

PII: S1876-2018(18)30010-8  
DOI: <https://doi.org/10.1016/j.ajp.2018.08.004>  
Reference: AJP 1509



To appear in:

Received date: 8-1-2018  
Revised date: 22-7-2018  
Accepted date: 4-8-2018

Please cite this article as: Badr HE, Francis K, Psychosocial perspective and suicidal behaviors correlated with adolescent male smoking and illicit drug use, *Asian Journal of Psychiatry* (2018), <https://doi.org/10.1016/j.ajp.2018.08.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Psychosocial perspective and suicidal behaviors correlated with  
adolescent male smoking and illicit drug use**

Hanan E. Badr, MD, MPH, Dr.PH & Konstantinos Francis, MD, PhD

**Corresponding author:**

Hanan E. Badr, MD, MPH, Dr. PH

Department of Community Medicine and Behavioural Sciences

Faculty of Medicine, Kuwait University

P.O. Box 24923 Safat

13110 Kuwait

Telephone number: +965 9956 5177

Fax number: +965 2533 8948

Email: [hanan@hsc.edu.kw](mailto:hanan@hsc.edu.kw) / [hnnbadr@gmail.com](mailto:hnnbadr@gmail.com)

Konstantinos Francis, MD, PhD

Consultant Child Psychiatrist - Kuwait Center for Mental Health

Email: [cfrancis@otenet.gr](mailto:cfrancis@otenet.gr)

**Highlights for Review**

The manuscript highlighted the following main points in the first study that acknowledge illicit drug use among teenagers in Kuwait among this age:

- Smoking and illicit drug use is highly prevalent among male adolescents in Kuwait.
- Smoking and illicit drug use are associated with suicide behaviors.
- Psychosocial factors contributed to smoking and illicit drug use habits.
- Using illicit drugs amplified the risk of suicide behaviors more than smoking only.

Download English Version:

<https://daneshyari.com/en/article/6787251>

Download Persian Version:

<https://daneshyari.com/article/6787251>

[Daneshyari.com](https://daneshyari.com)