



Enzymatic reduction of α -substituted ketones with concomitant dynamic kinetic resolution



Ronald L. Hanson, Zhiwei Guo, Francisco González-Bobes, Michaël D.B. Fenster, Animesh Goswami*

Chemical and Synthetic Development, Bristol-Myers Squibb, One Squibb Drive, New Brunswick, NJ 08903, USA

ARTICLE INFO

Article history:

Received 6 June 2016

Received in revised form 14 July 2016

Accepted 15 July 2016

Available online 18 July 2016

Keywords:

Enzymatic reduction

Dynamic kinetic resolution

α -Substituted ketones

Chiral alcohols

ABSTRACT

Racemic α -substituted ketones were converted to the corresponding chiral alcohols with high diastereo- and enantioselectivities using enzymatic reduction with concomitant dynamic kinetic resolution. Reductions of N-protected α -amino ketones by microorganisms and commercial enzymes provided N-protected α -amino alcohols. Choice of buffer was found to be a crucial factor for the successful reduction and simultaneous dynamic resolution of an α -methyl ketone to the corresponding chiral alcohol.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Ketoreductases are frequently used to reduce ketones to chiral alcohols [1]. With an enolizable group at the α -position the stereochemistry at the adjacent carbon can often be established due to keto-enol tautomerism and selection of only one of the ketone enantiomers for reduction by the enzyme leading to reduction with concomitant dynamic kinetic resolution (DKR) [2–5]. Although there are many examples of such ketone reduction with DKR, there are only a few reports on synthesis of α -alkyl or N-protected α -amino hydroxy esters from the corresponding ketones using enzymatic reduction of ketone and DKR [6–8].

Compound **1** (Fig. 1) is a positive allosteric modulator for the metabotropic glutamate receptor (mGluR5) considered for development as a possible treatment of neurological disorders [9]. The initial synthetic route to **1** used an intermediate *syn*-(1*R*,2*R*) N-protected amino alcohol **2** prepared by an aminohydroxylation catalyzed by OsO₄ which was further converted to a key intermediate **3** for the synthesis of **1** [10]. The aminohydroxylation reaction had only moderate regioselectivity and enantioselectivity, and required careful attention to downstream removal of osmium. Using alternative chemistry, both the *syn*-**2**, *anti*-**4** protected amino

alcohols were further converted to **3** the penultimate for the initial synthetic route.

The alcohol **8** (Fig. 2) is a key chiral intermediate for a GPR40 modulator drug candidate **10** for the treatment of diabetes [11,12]. Initial synthesis of **8** was carried out in seven steps with three isolations involving Evan's auxiliary to create chiral centers, and finally hydrogenation of benzyl amine **9** to **8** with an overall yield of 12% [13].

This report describes the preparation of both *syn*- and *anti* N-protected amino alcohols **2** and **4** from ketone **5** (Fig. 1), and *anti* alcohols **9** and **12** from ketone **11** (Fig. 2). In both cases, an α -substituted racemic ketone was enzymatically reduced to a chiral alcohol with the stereochemistry at the second chiral center (α -carbon) set via a dynamic resolution. Summaries of our work on enzymatic reductions and the detailed synthesis of drug candidates have been reported by us in separate publications [13,14].

2. Experimental

Starting materials and product markers were prepared as described [9–14]. Enzymes were purchased from commercial sources. Microorganisms were purchased from ATCC (ATCC#) or obtained from our own culture collection (SC#). Reaction conversion and purity of isolated product were determined from achiral HPLC area percentage (AP). Diastereomeric excess (de) and enantiomeric excess (ee) were determined from chiral HPLC.

* Corresponding author.

E-mail address: animesh.goswami@bms.com (A. Goswami).




Fig. 1. Reduction of ketone 5.




Fig. 2. Reduction of ketone 11.

2.1. Screening for reduction with commercial enzymes

Ketoreductases (1 mg) were screened in 1-mL reactions in 24-well plates containing 1 mg ketone dissolved in 0.02 mL DMSO, 1 mM NADP or NAD, 5 mg glucose, 0.05 mg glucose dehydrogenase (3.2 U, Amano), 1 mM dithiothreitol and 0.98 mL 0.1 M potassium phosphate buffer pH 7. After 20–87 h at 30 °C and 400 rpm, reactions were quenched with 1 mL methanol and analyzed by HPLC. For the Codexis screening kit and KRED panels the manufacturer's protocol was followed.

2.2. Screening for reduction with microorganisms

Microbial strains were screened in 24-well plates containing inoculum and 2 mL F7 medium (1% malt extract, 1% yeast extract, 0.1% peptone and 2% dextrose adjusted to pH 7) per well. Plates were incubated for 1 day at 28 °C, 600 rpm then 2 mg ketone in 0.04 mL DMSO was added and the incubation was continued for 2 days. The incubations were quenched by addition of 2 mL methanol per well and analyzed by HPLC.

دانلود مقاله



<http://daneshyari.com/article/69297>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات