

Urease-catalyzed synthesis of aminocyanopyridines from urea under fully green conditions



Fatemeh Tamaddon*, Somayeh Ghazi, Mohammad Reza Noorbala

Department of Chemistry, Yazd University, Yazd 89195-741, Iran

ARTICLE INFO

Article history:

Received 21 October 2015

Received in revised form 23 February 2016

Accepted 24 February 2016

Available online 26 February 2016

Keywords:

Urease

Enzymatic reactions

In situ generation of ammonia

Biocompatible

Pyridines

ABSTRACT

This is an original work on catalytic performance of urease in organic synthesis which in one-pot dissociation of urea and condensation of the *in situ* generated ammonia with aldehydes, acetophenones, and malononitrile occurs in water to give 2-amino-3-cyanopyridines. Comparative experiments with ammonium salts supported the enzymatic specificity of 0.01 g/mL (50 U/mg) of urease for bio-production of ammonia, while trace amount of heavy metal ions such as Pb^{2+} , Hg^{2+} , and Ag^+ inhibit these specific reactions. The scalability and promiscuity of urease facilitate the applicability of the process for biotechnological organic reactions based on ammonia.

© 2016 Published by Elsevier B.V.

1. Introduction

Replacement of hazardous procedures with eco-environmentally benign alternatives is one of the essential challenges of green chemistry [1,2], though the enzymatic water-based organic reactions are of the most attractive processes [3–7] due to the biocompatibility, non-toxicity, and easier workup. Hydrolases are known as substrate specific biocatalysts in organic reactions with high enzymatic promiscuity [8] that categorized as condition promiscuity, substrate promiscuity, and catalytic promiscuity. These promiscuities define as preservation of enzymatic activity under different conditions [9–11], broad range of substrate specificity [12,13], and the ability of the active site of a single enzyme to catalyze various chemical transformations [14]. While immobilization of enzymes improves their reusability, purity, stability, activity, specificity, selectivity, and inhibitions [15], hydrolase enzymes illustrated the catalytic promiscuity in water-based enzymatic organic synthesis [16–18]. Urease is a binuclear Ni-containing hydrolase enzyme with high substrate specificity for dissociation of urea to ammonia which in conjugation with urea can be considered as a bio source of nitrogen instead of the risky odorous ones.

Pyridine derivatives are among the interest heterocycles which constitute the back bone of many natural products and bioactive molecules [19] with anti-microbial, cardiotoxic, anti-parkinsonism, anti HIV, antitumor, and anti-inflammatory activities. Multi-functionalized pyridines with amino- and cyano-substituents (AmCyPys) are of these biologically active and fluorescent molecules [20] that extra conversion of the cyano and amino into the other functional groups [21] make them favored for synthesis of bioorganic compounds such as vitamin B₃ [22]. There are various malononitrile-based multi-component reactions (MCRs) to synthesis of these potent pyridines [19,23–25], although due to the benefits of the MCRs in water [26–30] and malononitrile-based synthesis of AmCyPys [31–35], their conjugation with advanced enzymatic reactions is highly desirable. To the best of our knowledge this is the first report on the application of urease in organic synthesis that due to the benefits of the water-based enzymatic reactions have been designed to improve the four-component synthesis of AmCyPys under fully biocompatible conditions (Scheme 1).


2. Material and methods

2.1. Materials and analytical methods

All materials and urease with art no from jack beans were purchased from Merck. All reagents were obtained from commercial suppliers and were used without further purification unless otherwise noted. The NMR spectra were recorded on a Bruker 500 MHz

* Corresponding author.

E-mail addresses: ftamaddon@yazd.ac.ir (F. Tamaddon), s.ghazi@stu.yazd.ac.ir (S. Ghazi), noorbala@yazd.ac.ir (M.R. Noorbala).



Scheme 1. Biocompatible synthesis of AmCyPys in water.

Table 1
Optimization of urease-catalyzed reaction.^a

Entry	Conditions				Yield ^b (%)
	Urea (mmol)	Temp. (°C)	Urease (g)	Time (h)	
1	1.2	60	0.01	8	50
2	1.2	70	0.005	8	40
3	1.2	70	0.01	4.5	68
4	1.2	70	0.02	4.5	70
5	1.2	80	0.01	6	50
6	2	70	0.01	3.5	75
7	3	70	0.01	2.5	80
8	4	60	0.01	3	60
9	4	65	0.01	2	70
10	4	70	0.005	5	60
11	4	70	0.01	1.5	85
12	4	70	0.015	1.5	85
13	4	70	0.02	1.4	87
14	5	70	0.02	1.5	85
15	4	80	0.01	3	60
16	4 ^c	70	0.01	12	–

^a Reaction was performed with benzaldehyde, acetophenone, malononitrile, and urea (in ratio of 1:1:1:1.2–4 mmol) in 0.5 mL of water and 0.005–0.02 g of urease.

^b Isolated yield based on 3.

^c Reaction was run with thiourea.

instrument using DMSO-*d*₆ or CDCl₃ as solvents. Chemical shifts (δ) were expressed in ppm with TMS as internal standard, and coupling constants (*J*) were reported in Hz.

2.2. General procedure for the synthesis of 2-amino-3-cyanopyridines

A mixture of an aldehyde (1 mmol), substituted acetophenone (1 mmol), malononitrile (1 mmol), urea (1.5 mmol) and urease (0.01 g (50 U/mg)) in 0.25 mL water was stirred at 70 °C for appropriate times (Table 1). The progress of the reaction was monitored by TLC (70:30, *n*-hexane/acetone). The reaction mixture was cooled after completion and 95% cold EtOH (2 mL) was added. The precipitate was filtered off, washed with cold ethanol, and dried to give the pure product.

3. Results and discussion

Initially, to optimize the reaction conditions for access to the maximum activity and specificity of enzyme, the catalytic performance of the commercially available urease was investigated in the reaction of benzaldehyde (1), acetophenone (2), malononitrile (3), and urea (4) screened at various temperatures and loading of ureases (Table 2).

As results show, the maximum yield of 2-amino-4,6-diphenylnicotinonitrile **5** was obtained in water using 0.01 g




Fig. 1. Effect of enzyme loading on the reaction time and yield.




Fig. 2. Effect of urea loading on the yield of **5**.




Fig. 3. Effect of temperature on the rate of urease-catalyzed reaction.

of commercially available urease after 1.5 h at 70 °C (Table 1, entry 13). A control experiment without urease under similar conditions did not give the product **5** even after 24 h and confirmed the catalytic role of urease for dissociation of urea to ammonia.

To distinct the catalytic role of urease for the *in situ* generation of ammonia or further transformations, the model reaction was run with ammonium acetate instead of urea in the presence and absence of 0.01 g of urease. The similar yields and reaction times of both of reactions support the substrate specificity of urease for the only *in situ* generation of ammonia from urea in the first step of the reaction. Another control experiment with thiourea supported the promiscuity of urease for exclusive dissociation of urea (entry 16).

To determine the details of this enzymatic reaction, the influences of enzyme loading (Fig. 1), molar ratio of starting materials (Fig. 2), and reaction temperature (Fig. 3) were finely investigated. In respect of the reaction time and yield, the optimal reaction temperature, enzyme loading, and molar ratios of components 1–4 were 70 °C, 0.01 g/mL of enzyme, and mole ratio of 1:1:1:4 were (1)–(4), respectively.

دانلود مقاله



<http://daneshyari.com/article/69438>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات