Accepted Manuscript

Short Communication

Response of humic acid formation to elevated nitrate during chicken manure composting

Mingzi Shi, Zimin Wei, Liqin Wang, Junqiu Wu, Duoying Zhang, Dan Wei, Yu Tang, Yue Zhao

PII: S0960-8524(18)30404-8

DOI: https://doi.org/10.1016/j.biortech.2018.03.056

Reference: BITE 19695

To appear in: Bioresource Technology

Received Date: 11 January 2018 Revised Date: 7 March 2018 Accepted Date: 11 March 2018



Please cite this article as: Shi, M., Wei, Z., Wang, L., Wu, J., Zhang, D., Wei, D., Tang, Y., Zhao, Y., Response of humic acid formation to elevated nitrate during chicken manure composting, *Bioresource Technology* (2018), doi: https://doi.org/10.1016/j.biortech.2018.03.056

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Response of humic acid formation to elevated nitrate during chicken manure composting

Mingzi Shi^a, Zimin Wei^a, Liqin Wang^a, Junqiu Wu^a, Duoying Zhang^b, Dan Wei^c, Yu Tang^a, Yue Zhao^{a,*}

^a College of Life Science, Northeast Agricultural University, Harbin 150030, China.

^b School of Civil Engineering, Heilongjiang University, Harbin 150080, China.

^c Soil and Environmental Resources Institute, Heilongjiang Academy of Agricultural

Science, Harbin, Heilongjiang 150086, China.

Corresponding author Address: College of Life Science, Northeast Agricultural

University, Harbin 150030, China.

Tel/Fax: +86-018745724658

E-mail address: zhaoyue@neau.edu.cn or zhao1970yue@163.com

Abstract: Nitrate can stimulate microbes to degrade aromatic compounds, whereas humic acid (HA) as a high molecular weight aromatic compound, its formation may be affected by elevated nitrate during composting. Therefore, this study is conducted to determine the effect of elevated nitrate on HA formation. Five tests were executed by adding different nitrate concentrations to chicken manure composting. Results demonstrate that the concentration of HA in treatment group is significantly decreased compared with control group (p < 0.05), especially in the highest nitrate concentration group. RDA indicates that the microbes associated with HA and environmental parameters are influenced by elevated nitrate. Furthermore, structural equation model

Download English Version:

https://daneshyari.com/en/article/7067658

Download Persian Version:

https://daneshyari.com/article/7067658

<u>Daneshyari.com</u>