ARTICLE IN PRESS

Journal of King Saud University - Science xxx (2018) xxx-xxx

Contents lists available at ScienceDirect



Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

Proposition of an appropriate technique to diagnose catheters fungal infectivities

Sidi Mohammed Lahbib Seddiki ^{a,b,*}, Zahia Boucherit-Otmani ^b, Yassine Moustafa Mahdad ^a, Bendahmane Ahmed Fouad ^c, Dennis Kunkel ^d

^a University Center of Naâma, Algeria

^b Laboratory: Antifungal Antibiotic, Physico-Chemical Synthesis and Biological Activity, University of Tlemcen, Algeria

^c Department of Hematology, University Hospital, University of Tlemcen, Algeria

^d Dennis Kunkel Microscopy Inc, Kailua, HI, USA

ARTICLE INFO

Article history: Received 11 November 2017 Accepted 10 April 2018 Available online xxxx

Keywords: Fungal infectivities Candida sp. Catheters Diagnostic Technical

ABSTRACT

Despite their great importance in the hospital environment, catheters are medical devices that can harm patients following their alterations by bacteria and/or fungal microorganisms. These alterations are classified into three types of infectivities, which may be simple contaminations/colonizations or serious infections. Bacterial infectivities were well studied using the technique of (Brun-Buisson et al., 1987); however, no technique was designed for yeast. Thus, we aimed to adapt the bacterial technique toward the assessment of the fungal infectivities. In order to check its reproducibility, both techniques were used in parallel. The results obtained were not consistent. That of Brun-Buisson et al. reflected more the catheters fungal contamination or colonization; nevertheless, the modified technique was better appropriate to the fungal infections of catheters considering the reduced time for its realization.

© 2018 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

The scourge of nosocomial infections is a real public health problem due to its epidemiological frequency and its socioeconomic cost (Fki et al., 2008). Moreover, there is considerable evidence of an increase in fungal infections acquired in hospitals, including candidiasis in immune-compromised patients with an implanted catheter (Martin et al., 2003; Patterson, 2007; Luzzati et al., 2005; Miceli et al., 2011). These infections can occur at any time when using these devices. Indeed, the strong use of catheters accentuates the exposure of patients to the risks of nosocomial candidiasis (Carrière and Marchandin, 2001) whatever is the type of their material or the pathology that required its implantation (Mermel et al., 2001; Kojic and Darouiche, 2004). On the other hand, it has been described that the diagnosis of catheter-related

* Corresponding author at: Laboratory: Antifungal Antibiotic, Physico-Chemical Synthesis and Biological Activity, University of Tlemcen, Algeria.

E-mail address: seddiki.med@gmail.com (S.M.L. Seddiki).

Peer review under responsibility of King Saud University.



candidemia is difficult to prove before the removal of the catheter, especially when Candida sp. forms a biofilm on its surface (Gallien et al., 2007). Furthermore, biofilms constitute a potential source of microorganisms endowed with properties of resistance to antifungal agents (Chandra et al., 2008; Espinasse et al., 2010). Thus, (Brun-Buisson, 1994) suggests that it is imperative to distinguish the infection of the catheter from its simple colonization or contamination. In the same context, we prescribed the term infectivity of catheters to reveal the degree of infectiveness of these devices (Seddiki et al., 2013), which may help clinicians in their decision to initiate antifungal therapy. The classification of the catheters' infectivities is determined by the presence or the absence of local and/or general clinical manifestations (local signs, infectious syndrome) and by their comparison with the qualitative results with regard to possible presence of microorganisms on the internal and/or external surface of the catheter (Carrière and Marchandin, 2001; Mimoz et al., 2007; Ryan et al., 1974). Whatever the clinical data is, the only way to assert catheter infection is to remove it and put it into culture. Thus, the positivity of the culture of the distal end of the catheter relative to the threshold of significance allows the establishment of this classification (Carrière and Marchandin, 2001; Merrer, 2005). It is nevertheless important to recall that the threshold of 15 colony-forming units (CFU) with the semi-quantitative technique of (Maki et al., 1977) was criticized

https://doi.org/10.1016/j.jksus.2018.04.012

1018-3647/© 2018 Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

by several authors (Snydman et al., 1982; Rello, et al., 1991; Moyer et al., 1983; Hilton et al., 1988). They reproach him with his lack of sensitivity knowing that this method explores only the external face of the catheter. On the other hand, this threshold value is 10^3 CFU/mL with the quantitative technique of (Brun-Buisson et al., 1987), which examines the internal and external faces of the catheter. However, these two techniques are usually recognized for the evaluation of bacterial infectivities. Owing to this, we aimed to assess the productivity of the yeast-adapted technique by examining the correlation between fungal infectivities results obtained using both techniques.

2. Material and methods

2.1. Sampling

Samples were collected for two months in three different hospitals in Tlemcen, Algeria. According to the recommendations of the literature (Gürcüoglu et al., 2010; Quinet, 2006), catheters implanted for 48 h and more were instantly collected after their removal from the patients. For this, their distal ends were cut using a sterile chisel and then placed in tubes containing 1 mL of sterile physiological solution. To avoid contamination, the tubes were opened nearby the flame of a torch and then quickly closed. Subsequently, the samples were transported to the laboratory in a cooled bag.

2.2. Identification of fungal strains

The fungal strains isolated from the catheters were purified and then identified by micro-culture series techniques and biochemical identification galleries (API 20 C AUX, Biomerieux, France)

2.3. Evaluation of different types of fungal infectivities

In order to assess the fungal infectivities of the catheters, the method of (Brun-Buisson et al., 1987) and the one modified by our team were used at the same time. The first one consists in carrying out a culture of the sample on agar, and the results were then evaluated by an enumeration of CFU/mL. The second was based on a direct enumeration of yeasts using a hematometer (Thoma blade) and then, the results were reported in cells/mL.

Briefly, the sampling tubes were first vortexing for one minute, this operation makes it possible to detach the yeasts adhered to the walls of the catheters and to examine both their internal and external faces (Brun-Buisson et al., 1987; Boucherit-Atmani et al., 2011). Then, 10 μ L were taken with a micropipette from each tube. Petri dishes previously prepared with a Sabouraud agar supplemented with 50 μ g/mL of chloramphenicol were inoculated by spreading with a rake. The addition of chloramphenicol (Bio-Rad, France) allows the inhibition of bacteria. The dishes were then incubated at 37 °C for 24 h up to 5 days and the count of the colonies appearing on the agar was reported in CFUs/mL. The experiment was performed twice for each sample.

In parallel, the counting of the yeast cells was carried out (three replicates) using the Thoma blades by the deposition of a volume of the specimen between the blade and the coverslip. The cells count was then carried out under an optical microscope (Magnification \times 400) and was reported in cells/mL. Only yeasts within the lines delineating the surface of the Thoma blade were counted (Gognies and Belarbi, 2010).

As previously described, the different types of fungal infectivities of the catheters (infection, colonization and contamination) were evaluated taking into account the threshold of significance $(10^3 \text{ CFU/mL or cells/mL})$ and the clinical data recorded during the sampling (clinical signs of infection). In order to situate the results in their epidemiological context, a statistical analysis of the variance was carried out via GenStat Discovery 3 software (VSN International). The threshold of significance: $\alpha = 0.01$.

This study was complemented by taking pictures of the view field of the optical microscope during the counting of the yeast cells. In addition, some catheter segments were examined using field emission scanning electron microscopy (Hitachi S-4800, Japan).

3. Results and discussion

During the study period, 399 catheters were taken from 374 inpatients. Some patients had two or more catheter types at the same time. The identification of isolated yeasts revealed 42 strains of Candida distributed among the following species: Candida albicans, C. glabrata, C. parapsilosis, C. guilliermondii, C. famata, C. tropicalis, C. krusei and C. kefyr. The involvement of each species in relation to catheter type, patient clinical data, admissions services and hospital structures was variable (unpublished data). Many studies have shown that C. albicans are the most involved in fungal infections related to catheters use (Boucherit-Atmani et al., 2011; Beck-Sague and Jarvis, 1993; Dekeyser et al., 2003; Phan et al., 2007; Pereira and Maisch, 2012). However, the isolation of nonalbicans Candida species from medical devices was constantly increasing; this indicates the non-negligible importance of these species in fungal pathology (Dekeyser et al., 2003; Dupont, 2007; Trofa et al., 2008; Miceli et al., 2011; Taieb et al., 2011; Seddiki et al., 2015). Overall, regardless of the species involved, contamination was the dominant type of infectivity, followed by colonization and then infections. It has been postponed that colonizations present an epidemiological interest but have no clinical relevance (Rijnders et al., 2002). Despite this, it may progress to a serious infection

According to our results, variability was observed according to the method used. 86.22% (344) of the samples did not reveal the presence of any colony or cell using either of the two methods cited above. On the contrary, when either method did not reveal any CFU or fungal cell (4.26% of all cases), the other method showed at least one CFU or cell and vice versa. On the other hand, in only 1% of cases, the count of CFUs according to the Brun-Buisson method was difficult to determine given their large number in the same Petri dish.

Moreover, all the other results were proved to be positive with both methods despite their divergence in many cases. In addition, according to the analysis of the variance, these results showed a



Fig. 1. Comparison of two methods of enumeration. Vertical bars indicate standard errors of average. Bars with the same letters (a) indicate a non-significant difference between the means according to the Duncan test of means of separation with a significance level of 0,01.

Download English Version:

https://daneshyari.com/en/article/7216585

Download Persian Version:

https://daneshyari.com/article/7216585

Daneshyari.com