



Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Search for dark matter effects on gravitational signals from neutron star mergers

John Ellis^{a,b,c}, Andi Hektor^a, Gert Hütsi^{a,d}, Kristjan Kannike^a, Luca Marzola^a,
Martti Raidal^a, Ville Vaskonen^a

^a NICPB, Rävåla pst. 10, 10143 Tallinn, Estonia

^b Theoretical Particle Physics and Cosmology Group, Physics Department, King's College London, London WC2R 2LS, United Kingdom

^c Theoretical Physics Department, CERN, Switzerland

^d Tartu Observatory, Tõravere 61602, Estonia

ARTICLE INFO

Article history:

Received 10 January 2018

Accepted 24 April 2018

Available online xxxx

Editor: M. Trodden

ABSTRACT

Motivated by the recent detection of the gravitational wave signal emitted by a binary neutron star merger, we analyse the possible impact of dark matter on such signals. We show that dark matter cores in merging neutron stars may yield an observable supplementary peak in the gravitational wave power spectral density following the merger, which could be distinguished from the features produced by the neutron components.

© 2018 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The discovery of gravitational waves (GW) by the LIGO and VIRGO collaborations has opened a new observational window on the Universe. The first measured GW signals were emitted from the merging of black hole binaries [1–3] and spectacularly confirmed the predictions of general relativity and constrained its extensions in the strong-field regime [4,5]. Recently the LIGO and VIRGO collaborations have also detected a GW signal originating during the inspiral leading to the merger of two neutron stars (NS), GW170817 [6]. On this occasion, the observation of the GW signal was accompanied by counterparts in many bands of the electromagnetic spectrum [7], as was expected for a NS-NS merger, notably including the observation of a gamma-ray burst, GRB170817A [8]. This new discovery is an important confirmation that NS-NS mergers are the originators of short gamma-ray bursts [9–12], and play a major role in the r-process nucleosynthesis as demonstrated by the ejection of synthesised material via subsequent kilonova explosions [13–18].

Conventional models of NS-NS merger yields three separate peaks in the GW emission following the merger [19–28]: for a review with a comprehensive list of references, see [29]. The

strongest observable peak in these simulations results from the rotation of the bar-deformed hyper-massive neutron star in the first instants after the merger. This strong peak is flanked by two less pronounced peaks that are produced by the dynamics of the remnants of the original stellar cores. Despite the complexity of the simulations, the above-mentioned qualitative features of the GW signal are captured by a simple mechanical model [22,23] that reproduces qualitatively the detailed calculation of the GW waveform emitted in these events. In the case of GW170817, no post-merger GW signal was observed, but future observations of such signals would have the capability to constrain models of dark matter as well nuclear matter, as we now discuss.

We show in this Letter how observations of GW signals from NS-NS mergers also offer a new window on the dark matter (DM) component of the Universe [30], through a possible supplementary peak in the power spectral density (PSD) of the GW emission following a NS-NS merger. We introduce additional features into the simple mechanical model of NS-NS mergers to account for the possible presence of DM cores within the neutron stars. We show that these DM cores may generate an additional well-defined peak in the PSD of the GW signal emitted after the merger. If the DM cores contribute about 5–10% of the NS mass, we find that the DM peak is observable and can be clearly distinguished from the features of the signal induced by the remnant neutron matter cores for a wide range of effective parameters.

It should be noted that ordinary WIMP DM cannot condense inside stars or compact stellar remnants in quantities as large as

E-mail addresses: john.ellis@cern.ch (J. Ellis), andi.hektor@cern.ch (A. Hektor), gert.hutsi@to.ee (G. Hütsi), kristjan.kannike@cern.ch (K. Kannike), luca.marzola@cern.ch (L. Marzola), martti.raidal@cern.ch (M. Raidal), ville.vaskonen@kbf.ee (V. Vaskonen).

<https://doi.org/10.1016/j.physletb.2018.04.048>

0370-2693/© 2018 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.




Fig. 1. A schematic diagram of the mechanical model adopted to investigate the possible effects of DM components on the GW signal emitted by NS-NS mergers. The DM cores (in black) move in an environment constituted of the remnant neutron stellar core (in white) and a disk of neutron matter enclosing the latter.

discussed in this study. For example, during a period of ~ 10 Gyr, even after allowing for DM densities $\sim 10^4$ times larger than the local value (as may be reachable in the central parts of the Galaxy), the typical amount of accreted WIMP DM does not significantly exceed $\sim 10^{-10} M_\odot$ [31–34]. However, having a nonstandard dark sector with dissipative (but subdominant) DM component (e.g., as in [35–37]) might lead to the formation of DM-admixed stars with noticeable DM content. For example, DM-admixed NSs have been discussed previously in [38–44]. Moreover, the formation of asymmetric DM stars, which might potentially serve as cores for the NSs studied in this paper, have been investigated in [45,46]. Although the details how these objects might have formed require much deeper investigation, which is beyond the scope of this study, we propose the following illustrative scenario. If a small fraction of DM ($\lesssim 10\%$) is strongly self-interacting then, as demonstrated in [37], it can lead to early formation of black holes (BHs) that can serve as seeds for subsequent supermassive BH formation, alleviating tension with the small amount of time available for their formation in the standard LCDM picture. Depending on the shape of the initial perturbation spectrum, in addition to BH formation, one would also expect formation of smaller structures analogous to visible-sector NSs. Later on, these compact dark objects can serve as accretion centres for baryonic matter, which when accreted in sufficient quantities can lead to the formation of DM admixed NSs. For this scenario to work no interaction beyond gravity is needed between the dark and the visible sectors.

The exploratory results in this Letter clearly indicate that observations of NS-NS mergers have the potential to probe the properties of such self-interacting DM, complementing dedicated searches. Our work underlines the need for further simulations to assess with more precision the possible impact of a DM component on the evolution of NS-NS mergers and disentangle its effects from phenomena associated with different nuclear equations of state.

2. A mechanical model

In order to assess the impact of a possible DM component on the GW signal emitted by a NS-NS merger, we extend the mechanical model originally presented in [22,23]. We include two additional DM cores of (for simplicity) equal mass $m_d/2$, which move in the gravitational potential of the hyper-massive NS generated in the merger. The latter is modelled as a disk of mass M containing two coupled oscillators of (again for simplicity) equal mass $m_n/2$, which represent the neutron components of the remnant stellar cores, as illustrated in Fig. 1. The disk and the neutron stellar cores rotate at a common angular velocity $\Omega = \dot{\theta}_n$, whereas the dark cores rotate at an angular velocity $\dot{\theta}_d$. We account for differences in the strength of DM and neutron interactions by considering different spring constants and damping factors.

The Lagrangian of the mechanical system takes the form

$$L = \frac{m_n}{2} (\dot{r}_n^2 + (r_n \dot{\theta}_n)^2) + \frac{m_d}{2} (\dot{r}_d^2 + (r_d \dot{\theta}_d)^2) + \frac{MR^2 \dot{\theta}_n^2}{4} + 2k_n (r_n - a_n)^2 + 2k_d (r_d - a_d)^2, \quad (1)$$

where an overdot indicates differentiation with respect to time and the subscripts n and d distinguish quantities related to the nuclear and dark cores, respectively. The parameters k_n and k_d are effective ‘spring constants’ that characterize the oscillations of the nuclear and DM cores in the gravitational potential and the medium. The effective damping induced by interactions is encoded in the parameters b_n and b_d , and that due to GW emission, c_n and c_d , is included by adding the term

$$D = -b_n \dot{r}_n^2 - b_d (\dot{r}_d^2 + (r_d (\dot{\theta}_n - \dot{\theta}_d))^2) - c_n (\dot{r}_n^2 + (r_n \dot{\theta}_n)^2) - c_d (\dot{r}_d^2 + (r_d \dot{\theta}_d)^2). \quad (2)$$

The equations of motion for the relevant degrees of freedom are obtained from the dissipative Euler–Lagrange equation

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_j} - \frac{\partial L}{\partial q_j} = \frac{\partial D}{\partial \dot{q}_j}. \quad (3)$$

In order to investigate the resulting GW signal, we compute the strain induced by the plus and cross polarisations at a distance D from the merger

$$h_+ = \frac{\ddot{I}_{xx} + \ddot{I}_{yy}}{D}, \quad h_\times = \frac{\ddot{I}_{xy}}{D}, \quad (4)$$

where I_{kl} , $k, l \in \{x, y, z\}$, are the components of the quadrupole-moment tensor:

$$I_{kl} = \sum_{j=n,d} \left(\bar{I}_{j,kl} - \frac{1}{3} \delta_{kl} \delta^{ab} \bar{I}_{j,ab} \right), \quad (5)$$

$$\bar{I}_{j,kl} = 3m_j x_{j,k} x_{j,l}.$$

The PSD is then obtained as

$$\tilde{h}(f) = \sqrt{\frac{|\tilde{h}_+(f)|^2 + |\tilde{h}_\times(f)|^2}{2}}, \quad (6)$$

where the tilded quantities are the Fourier transforms of the corresponding GW strains.

3. The effect of dark matter

As was shown in [22,23], the simple mechanical model without the DM components reproduces semi-quantitatively the peaks in the PSD found in a detailed simulation including general-relativistic and nuclear effects. Fig. 2 shows that, for suitable values of the model parameters, our implementation also reproduces well the full numerical-relativity strain h_+ for the full simulation reported in [22,23]. This comparison gives us confidence that such a simple model can capture correctly the dominant dynamical effects.

The PSD of the GW emission resulting from the above model (1) including the DM components is shown in Fig. 3. The top panel shows the PSD immediately following the merger of two equal-mass neutron stars, whereas the middle shows the PSD during a later time period. In calculating this representative point we set the parameters of the nuclear matter to $M = 10$, $m_n = 10$, $k_n = 0.25$, $b_n = 0.1$, $c_n = 0.02$ and $a_n = 1$, and the DM ones to

Download English Version:

<https://daneshyari.com/en/article/8186448>

Download Persian Version:

<https://daneshyari.com/article/8186448>

[Daneshyari.com](https://daneshyari.com)