Long-term treadmill exercise improves memory impairment through restoration of decreased synaptic adhesion molecule 1/2/3 induced by transient cerebral ischemia in the aged gerbil hippocampus

Ji Hyeon Ahn, Joon Ha Park, Jinseu Park, Myoung Cheol Shin, Jun Hwi Cho, In Hye Kim, Jeong-Hwi Cho, Tae-Kyeong Lee, Jae-Chul Lee, Bich Na Shin, Young-Myeong Kim, Choong Hyun Lee, In Koo Hwang, Il Jun Kang, Bai Hui Chen, Bing Chun Yan, Young Joo Lee, Moo-Ho Won, Soo Young Choi

PII: S0531-5565(17)30584-3
DOI: https://doi.org/10.1016/j.exger.2018.01.015
Reference: EXG 10259

To appear in: *Experimental Gerontology*

Received date: 2 August 2017
Revised date: 5 January 2018
Accepted date: 12 January 2018

Please cite this article as: Ji Hyeon Ahn, Joon Ha Park, Jinseu Park, Myoung Cheol Shin, Jun Hwi Cho, In Hye Kim, Jeong-Hwi Cho, Tae-Kyeong Lee, Jae-Chul Lee, Bich Na Shin, Young-Myeong Kim, Choong Hyun Lee, In Koo Hwang, Il Jun Kang, Bai Hui Chen, Bing Chun Yan, Young Joo Lee, Moo-Ho Won, Soo Young Choi , Long-term treadmill exercise improves memory impairment through restoration of decreased synaptic adhesion molecule 1/2/3 induced by transient cerebral ischemia in the aged gerbil hippocampus. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Exg(2017), https://doi.org/10.1016/j.exger.2018.01.015

Long-term treadmill exercise improves memory impairment through restoration of decreased synaptic adhesion molecule 1/2/3 induced by transient cerebral ischemia in the aged gerbil hippocampus

Ji Hyeon Ahn[1], Joon Ha Park[1], Jinseu Park[1], Myoung Cheol Shin[2], Jun Hwi Cho[2], In Hye Kim[3], Jeong-Hwi Cho[3], Tae-Kyeong Lee[3], Jae-Chul Lee[3], Bich Na Shin[4], Young-Myeong Kim[5], Choong Hyun Lee[6], In Koo Hwang[7], Il Jun Kang[8], Bai Hui Chen[9], Bing Chun Yan[10], Young Joo Lee[11], Moo-Ho Won[3*], Soo Young Choi[1*]

[1]*Department of Biomedical Science and Research Institute of Bioscience and Biotechnology, Hallym University, Chuncheon 24252, South Korea*

[2]*Department of Emergency Medicine, School of Medicine, Kangwon National University, Chuncheon 24341, South Korea*

[3]*Department of Neurobiology, School of Medicine, Kangwon National University, Chuncheon 24341, South Korea*

[4]*Department of Physiology, College of Medicine, Hallym University, Chuncheon 24252, South Korea*

[5]*Department of Molecular and Cellular Biochemistry, School of Medicine, Kangwon National University, Chuncheon 24341, South Korea*

[6]*Department of Pharmacy, College of Pharmacy, Dankook University, Cheonan 31116, South Korea*

[7]*Department of Anatomy and Cell Biology, College of Veterinary Medicine, and Research Institute for Veterinary Science, Seoul National University, Seoul 08826, South Korea*

[8]*Department of Food Science and Nutrition, Hallym University, Chuncheon 24252, South Korea*

[9]*Department of Histology and Embryology, Institute of Neuroscience, Wenzhou Medical University, Wenzhou, Zhejiang 325035, P.R. China*

[10] *Jiangsu Key Laboratory of Integrated Traditional Chinese and Western Medicine for Prevention and Treatment of Senile Diseases, Yangzhou 225001, People's Republic of China*

[11]*Department of Emergency Medicine, Seoul Hospital, College of Medicine, Sooncheonhyang University, Seoul 04401, South Korea.*

- Co-firsts: Ji Hyeon Ahn[1] and Joon Ha Park[1] have contributed equally to this article.