



Original article

Breaking seed dormancy in smooth loofah (*Luffa cylindrica* (L.) M. Roem.) using scarification and dry heat treatment



Sajee Chaodumrikul,^a Pichitra Kaewsonr,^{a,*} Pariyanuj Chulaka,^a Wanchai Chanprasert^b

^a Department of Horticulture, Faculty of Agriculture, Kasetsart University, Bangkok 10900, Thailand

^b Department of Agronomy, Faculty of Agriculture, Kasetsart University, Bangkok 10900, Thailand

ARTICLE INFO

Article history:

Received 10 April 2014

Accepted 30 September 2015

Available online 9 June 2016

Keywords:

Germination

Mean germination time

Seed coat structure

Seed dormancy

Sponge gourd

ABSTRACT

Hard seed dormancy is a common problem with smooth loofah seed. Scarification, clipping and dry heat were used to break dormancy in smooth loofah seed at the Department of Horticulture, Kasetsart University, Bangkok, Thailand. A completely randomized design with 20 treatments was used involving: untreated seed (control), clipping, scarifying by scarifier at 40, 70 and 100 revolutions per minute (rpm) for 1 min, and dry heat at 60 °C, 70 °C and 80 °C for 1–5 h. After breaking dormancy, seed germination was tested in four replicates, with 50 seeds per replicate. The thickness of the seed coat was measured under a digital microscope. The results showed that clipped seeds gave the highest germination (100%) and decreased mean germination time (3.58 d). Scarified seed using a scarifier at 70 and 100 rpm for 1 min resulted in germination rates of 67.0–75.5%, which was higher than for seeds scarified at 40 rpm. The dry heat-treated seeds at 60 °C for 3–5 h and at 70 °C for 2–5 h had germination of 71.0–80.5%. The outer layer of seed coat scarified at 100 rpm for 1 min was thinner than those of un-scarified seed samples. Dry heat had no effect on the seed coat thickness, but affected cells of the inner seed coat as the sclerenchymous cells showed disordered characteristics and were non-uniform and seemed to have been torn off. Dry heat treatment and scarification significantly improved germination compared to the control treatment. However, 80% germination may not be considered as an effective method at a commercial scale where 100% germination is needed. Further investigation of more accessions that may have different seed coat thicknesses may be needed.

Copyright © 2016, Kasetsart University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Luffa cylindrica (L.) M. Roem syn *L. aegyptiaca* Mill, commonly called smooth loofah or sponge gourd, is a member of the Cucurbitaceae family. The plants are economically important in many parts of the world—for instance, in China, Korea, Japan, India, Central America, as well as in Thailand—for their young fruits, which are used as ingredients for food, while ripening fruits are also used to produce consumer goods such as cleaning materials and engine oil filters (Oboh and Aluyor, 2009). The seeds are composed of 46% oil and 40% protein (Siemonsma and Piluek, 1993).

Some cucurbit species have severe problems with seed dormancy and viable seeds cannot germinate even in favorable environments because of seed coat impermeability is considered a major mechanism causing hard seed dormancy (Bradbeer, 1988).

Smooth loofah seed is considered hard-seeded with its thick seed structure and a hard seed coat; moreover, phenolic compounds including pectin or suberin on the surface of the seed coat restrict water uptake into the seed (Dojjode, 2001). Physical dormancy is caused by one or more water-impermeable layers of palisades (Baskin and Baskin, 2004). In addition, Singh and Dathan (1998) found that the seed coat of smooth loofah is characterized by upright epidermal cells with rod-like thickenings and narrow, palisade-like osteosclereids which cause physical dormancy of smooth loofah seed. Seed dormancy is the most important factor limiting germination and there are various ways to break hard seed coat such as clipping, scarifying, and dry heat (Bradbeer, 1988).

One technique that has been widely and successfully used for breaking hard seed dormancy is scarification, which involves removing the seed coat or rubbing it with sandpaper or subjecting it to a temperature treatment (Bradbeer, 1988). Pinmanee et al. (2001) reported that cutting the bottom of the bitter melon seed, but not removing the seed coat completely, increased germination from 40.5% to 70%, while germination was increased to 90% by removing the entire seed coat. The germination of watermelon seed

* Corresponding author.

E-mail address: agrptck@ku.ac.th (P. Kaewsonr).

has also been increased by removing the seed coat (Nerson, 2002). Loyma et al. (2009) reported that removing the seed coat of wax gourd seeds increased radicle emergence with scarification of 'Feang' and 'Fakkheaw' seeds by removing the outer testa and the outer and inner testa, significantly improving radicle emergence to 96.0% and 99.5%, respectively, of 'Feang' seeds and to 93.0% and 95.5%, respectively, of 'Fakkheaw' seeds, compared to the un-scarified seeds (83.0% for 'Feang' and 80% for 'Fakkheaw'). However, scarification may lead to embryo damage, abnormal seedlings, and dead seeds (Bradbeer, 1988).

Another technique for breaking hard seed dormancy is dry heat, which causes seed coat and perisperm dehydration and allows water and gases to enter the seed more quickly (Khan, 1980). The heat temperature for breaking dormancy in tropical and subtropical seeds is about 40–50 °C (ISTA, 2010). Sinviriyanon et al. (2011) reported that preheating at 70 °C could increase the seed germination of smooth loofah and decrease the mean germination time; however, the effect of preheating on seed germination depends on the accession. Moreover, a preheating treatment at 70 °C and 80 °C for 2, 4 and 6 h tended to increase the seed germination of smooth loofah but the germination was lower than 50% (Chamnongrit et al., 2011).

Seed dormancy is often a problem when planting smooth loofah as seeds have a low germination percentage and uniformity (Doijode, 2001). Therefore, the objective of this study was to determine the suitability of various techniques for breaking the dormancy of smooth loofah seed.

Materials and methods

Seed materials

Smooth loofah seeds of LF-01 were obtained from Chia Tai Co., Ltd (Bangkok, Thailand). Seeds were harvested and processed in September 2010. This study was conducted from October 2010 to January 2012 in a laboratory at the Department of Horticulture, Faculty of Agriculture, Kasetsart University, Bangkok, Thailand. The initial seed qualities tested were: 8.4% seed moisture content, 97.1 g per 1000 seed weight; 100% seed viability by tetrazolium test, 56.0% germination and 35.5% hard seed.

Methods of breaking seed dormancy

Seeds were treated in the laboratory to study methods of breaking dormancy. The experiment was arranged in a completely randomized design with 20 treatments. It consisted of two methods for breaking dormancy. The first method involved mechanical scarification and was carried out by clipping and scarification. Seeds were clipped opposite the embryo using a scalpel. Seeds were scarified using a scarifier at 40 revolutions per minute (rpm), 70 rpm and 100 rpm for 1 min. The second method was a dry heat treatment. Seeds were dried in a hot air oven at 60 °C, 70 °C and 80 °C for 1–5 h and then the seeds were kept in a desiccator for 30 min. Seeds after breaking dormancy were tested for quality in four replicates with 50 seeds per replicate.

Germination

Fifty seeds per replicate for four replicates were germinated in moist sand in a plastic box and kept in the germinator at 30 °C. First and final counts were done at 4 and 14 d after testing, respectively (ISTA, 2010). Normal seedlings were calculated as the germination percentage. The percentage of hard seeds was reported as that part of the germination percentage that remained viable with hard seed that could not absorb water.

Table 1

Germination and hard seed of smooth loofah by different methods of breaking dormancy.

Method of breaking dormancy	Germination (%)	Hard seed (%)
1. Untreated seeds (control)	56.0 ^{hi}	35.5 ^a
2. Clipping	100.0 ^a	0.0 ^j
3. Scarifier at 40 rpm for 1 min	63.0 ^{gh}	20.0 ^{bcd}
4. Scarifier at 70 rpm for 1 min	67.0 ^{efg}	23.5 ^b
5. Scarifier at 100 rpm for 1 min	75.5 ^{bcd}	16.5 ^{bcdef}
6. Dry heat at 60 °C for 1 h	66.0 ^{fg}	22.0 ^{bc}
7. Dry heat at 60 °C for 2 h	68.5 ^{defg}	19.5 ^{bcde}
8. Dry heat at 60 °C for 3 h	71.5 ^{cdef}	18.5 ^{bcdef}
9. Dry heat at 60 °C for 4 h	73.0 ^{bcdef}	17.5 ^{bcdef}
10. Dry heat at 60 °C for 5 h	80.5 ^b	12.5 ^{defgh}
11. Dry heat at 70 °C for 1 h	74.0 ^{bcdef}	17.5 ^{bcdef}
12. Dry heat at 70 °C for 2 h	78.0 ^{bc}	11.5 ^{efghi}
13. Dry heat at 70 °C for 3 h	80.0 ^b	11.0 ^{fghi}
14. Dry heat at 70 °C for 4 h	77.5 ^{bc}	7.5 ^{hi}
15. Dry heat at 70 °C for 5 h	75.0 ^{bcde}	13.0 ^{defgh}
16. Dry heat at 80 °C for 1 h	70.0 ^{cdefg}	15.5 ^{cdefg}
17. Dry heat at 80 °C for 2 h	26.5 ⁱ	4.0 ^{ij}
18. Dry heat at 80 °C for 3 h	22.5 ⁱ	4.0 ^{ij}
19. Dry heat at 80 °C for 4 h	21.0 ⁱ	8.5 ^{ghi}
20. Dry heat at 80 °C for 5 h	9.5 ^j	13.5 ^{defgh}
F-test	*	*
Coefficient of variation (%)	8.03	36.45

rpm = revolutions per minute.

* = significantly different at $p < 0.05$.

† = mean values in the same column followed by the same letter are not significantly different at $p < 0.05$ by Duncan's Multiple Range Test.

Mean germination time

Seeds were germinated as in germination test. The number of normal seedlings was counted from the day of the first count up to the day of the final count (14 d after testing). The mean germination time (MGT) was calculated using Equation (1) (Ellis and Roberts, 1981):

$$MGT = \frac{\sum nd}{\sum n} \quad (1)$$

Table 2

Mean germination time of smooth loofah seeds by different methods of breaking dormancy.

Method of breaking dormancy	Mean germination time (d)
1. Untreated seed (control)	5.17 ^{efg†}
2. Clipping	3.58 ^h
3. Scarifier at 40 rpm for 1 min	5.92 ^{cde}
4. Scarifier at 70 rpm for 1 min	5.45 ^{defg}
5. Scarifier at 100 rpm for 1 min	5.43 ^{defg}
6. Dry heat at 60 °C for 1 h	4.67 ^g
7. Dry heat at 60 °C for 2 h	5.38 ^{defg}
8. Dry heat at 60 °C for 3 h	5.37 ^{defg}
9. Dry heat at 60 °C for 4 h	4.95 ^{fg}
10. Dry heat at 60 °C for 5 h	4.87 ^g
11. Dry heat at 70 °C for 1 h	4.77 ^g
12. Dry heat at 70 °C for 2 h	5.75 ^{cdef}
13. Dry heat at 70 °C for 3 h	6.02 ^{cd}
14. Dry heat at 70 °C for 4 h	6.58 ^{bc}
15. Dry heat at 70 °C for 5 h	5.11 ^{efg}
16. Dry heat at 80 °C for 1 h	6.45 ^{bc}
17. Dry heat at 80 °C for 2 h	6.41 ^{bc}
18. Dry heat at 80 °C for 3 h	6.87 ^b
19. Dry heat at 80 °C for 4 h	8.16 ^a
20. Dry heat at 80 °C for 5 h	8.20 ^a
F-test	*
Coefficient of variation (%)	9.03

rpm = revolutions per minute.

* = significantly different at $p < 0.05$.

† = mean values in the same column followed by the same letter are not significantly different at $p < 0.05$ by Duncan's Multiple Range Test.

Download English Version:

<https://daneshyari.com/en/article/83080>

Download Persian Version:

<https://daneshyari.com/article/83080>

[Daneshyari.com](https://daneshyari.com)