



Original article

Growth performance of clonal rubber rootstocks and combining ability test with the scion of clone RRIM 600

Narid Khotcharat,^a Sayan Sdoodee,^{a,*} Upatham Meesawat^b^a Department of Plant Science, Faculty of Natural Resources, Prince of Songkla University, Songkhla 90112, Thailand^b Department of Biology, Faculty of Science, Prince of Songkla University, Songkhla 90112, Thailand

ARTICLE INFO

Article history:

Received 24 December 2014

Accepted 29 July 2015

Available online 9 June 2016

Keywords:

Compatibility

Root length density

Rootstocks

Rubber

Scion

ABSTRACT

Past screening of rubber rootstocks in southern Thailand has revealed five clones that showed good plant growth with tentative tolerance to white root disease. The current study tested the combining ability of these clones with the scion of clone RRIM 600, commonly grown in Thailand. The five clonal rootstocks and the clone RRIM 600 were grown in rhizoboxes (10 × 45 × 100 cm) in a completely randomized design, consisting of six treatments each with four replicates, making a total of 24 rhizoboxes. After growing for 6 mth, each seedling (1 cm stem diameter) was bud grafted with the scion RRIM 600. Root growth of the rootstocks was assessed by scanning a glass panel in the side of each rhizobox after 3 mth. It was found that the seedlings of clone#5 exhibited significantly higher root growth with tentative high shoot growth compared with the remaining treatments. The results showed that clone#5 exhibited the highest average root length density at 1.26 cm/cm² followed by clone#2 (0.90 cm/cm²) with a significant difference from the other clones. The investigation of graft union development showed that callus tissues developed on the graft union within 5 d after bud grafting (DAB). The graft union growth was almost fully developed over the wound areas and callus bridge formation was evident within 10 DAB. The new vascular tissues were connected with old cambium and vascular tissues of the scion and rootstocks within 20 DAB. The photosynthetic rate and stomatal conductance were assessed and showed 100% successful budding of rootstocks of clone RRIM 600 and clone#4, whereas the other clones achieved 75% success.

Copyright © 2016, Production and hosting by Elsevier B.V. on behalf of Kasetsart University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Southern Thailand is the country's main area of rubber planting, clonal material production and rubber production with Thailand's area of rubber plantation and the export of Thailand in 2013 being 3,548,274 ha and 3,664,941 t, respectively (Rubber Research Institute of Thailand, 2014). Normally, rubber seeds are collected from the smallholder's rubber plantation and therefore, most seeds are from rubber clone RRIM 600, which is mainly grown in southern Thailand and is sensitive to white root disease (Wattanasilakorn et al., 2012). Kaewchai and Soyong (2010) also noted that white root disease is a severe epidemic in southern Thailand. Therefore, some early introduced rubber clones were collected for rootstock screening of rootstock white root disease

resistance and it was found that there were some clones that exhibited tentative tolerance to white root disease (Wattanasilakorn et al., 2012). However, those rootstocks needed to be tested for their combining ability. Donald (1973) reported that 91% of the grafts were compatible when seedling families were grafted upon rootstocks which were compatible with both parents. During the formation of the graft union, researchers have observed callus proliferation (from both the rootstock and the scion), callus bridge formation, differentiation of new vascular tissue from callus cells and the production of secondary xylem and phloem (Hartmann et al., 2002). A low level or incorrect callus formation between the rootstock and scion could lead to defoliation, a reduction in scion growth and low survival of grafted plants (Oda et al., 2005; Johkan et al., 2009). The objectives of this study were: 1) to study the growth and development of the rubber clones used as rootstocks; 2) to study the compatibility of the rubber rootstocks with clone RRIM 600; and 3) to study the physiological responses of the RRIM 600 scion on the screening rootstocks.

* Corresponding author.

E-mail address: sayan.s@psu.ac.th (S. Sdoodee).

Materials and methods

The study was carried out from April 2012 to August 2013 in a glasshouse of the Faculty of Natural Resources, Prince of Songkla University, Songkhla province, Thailand.

Plant materials

The root growth of five early introduced clones collected from different areas in Songkhla province was tested. Collected seeds from each tree were georeferenced using a global positioning system (Table 1). Clone RRIM 600 was used for comparison. The experiment was established as a completely randomized design with six treatments and four replicates (one plant per replication). The seedlings (aged 2 mth) were grown in rhizoboxes (10 × 45 × 10 cm). One seedling was grown in each rhizobox (Fig. 1).

Root growth

A panel in each rhizobox was made of clear acrylic and covered with a black plastic sheet to avoid light exposure. Root images were photographed through the panel at every 20 cm depth. The images were analyzed using the Rootfly root analysis software which is a free, open-source software application to aid researchers in mini-rhizotron image analysis under the GNU General Public License (Stanley and Christina, 2011). The length, diameter and color of roots, as well as numbers alive and dead were recorded. All the experimental data were stored in a single file using the Rootfly software format.

Development of graft union

The graft union formation between 8-month-old rootstocks (clone#5) and scion (RRIM 600) was examined histologically using the standard method with paraffin described below. The samples (measured at 0.5 cm above and below the graft union) were collected at 5, 10 and 20 days after bud grafting (DAB). They were fixed in FAA II (formaldehyde:glacial acetic acid:70% ethyl alcohol; 5:5:90 v/v) for 48 h (Ruzin, 1999). The fixed samples were dehydrated in a tertiary-butyl-alcohol series and embedded in paraffin wax (Histoplast PE; Richard-Allan Scientific, Kalamazoo, MI, USA). Sections (12 μm thick) were cut using a rotary microtome and stained with safranin and fast green (Ruzin, 1999). All sections were observed under a light microscope.

Successful budding percentage

Four weeks after budding, a green bud indicated that budding had been successful. Then, the top of the rootstock was cut to



Fig. 1. Rhizoboxes used in the experiment.

induce shoot emergence. The percentage of successful buds remaining green was determined using Equation (1):

$$\text{Successful budding(\%)} = \frac{\text{Rootstock with successful buds}}{\text{Total number of rootstock}} \times 100 \quad (1)$$

Growth of the rubber tree

The plant height was measured at 10 cm from the soil level to the top of the plant shoot. The plant trunk diameter was measured 10 cm from the soil level. The number of leaves was determined by counting the number of compound leaves per plant.

Physiological responses

The photosynthetic rate (*A*) and stomatal conductance (*g_s*) were measured for 1000–1200 h using a portable photosynthesis system (LICOR-6400; LI-COR; Lincoln, NE, USA). Measurements were made on three fully expanded leaves for each treatment.

Results and discussion

Root growth of rubber trees

Most of the active root growth in the rhizoboxes was located within 20–40 cm depth from the soil surface with rapid proliferation. Clone#5 and RRIM 600 exhibited high extension root growth to deeper layers of 60–100 cm as shown in Fig. 2. Clone#5 had the highest

Table 1

Location of rubber clones collected in this study.

Name	Coordinates	Place of collection
Clone#1	7° 0' 23.1" N 100° 29' 52.8" E	Faculty of Environmental Management, Prince of Songkla University, Hat Yai, Songkhla
Clone#2	7° 0' 33.1" N 100° 29' 57.2" E	Khunying Long Athakravisunthon Learning Resources Center, Prince of Songkla University, Hat Yai, Songkhla
Clone#3	7° 0' 36.0" N 100° 29' 53.0" E	Office of the President, Prince of Songkla University, Hat Yai, Songkhla
Clone#4	7° 0' 37.8" N 100° 29' 56.7" E	Roundabout entrance halls, Prince of Songkla University, Hat Yai, Songkhla
Clone#5	7° 0' 29.6" N 100° 30' 2.2" E	Sritrang Reservoir, Prince of Songkla University, Hat Yai, Songkhla
RRIM 600	6° 8' 64.27" N 100° 42' 69.38" E	Klonghoykhong, Rubber Plantation, Songkhla

دانلود مقاله



<http://daneshyari.com/article/83082>



- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات