

Accepted Manuscript

Supervised pelvic floor muscle training versus attention-control massage treatment in patients with faecal incontinence: Statistical analysis plan for a randomised controlled trial

Anja Ussing, Inge Dahn, Ulla Due, Michael Sørensen, Janne Petersen, Thomas Bandholm

PII: S2451-8654(17)30048-0

DOI: [10.1016/j.conctc.2017.07.006](https://doi.org/10.1016/j.conctc.2017.07.006)

Reference: CONCTC 154

To appear in: *Contemporary Clinical Trials Communications*


Received Date: 14 March 2017

Revised Date: 12 July 2017

Accepted Date: 20 July 2017

Please cite this article as: A. Ussing, I. Dahn, U. Due, M. Sørensen, J. Petersen, T. Bandholm, Supervised pelvic floor muscle training versus attention-control massage treatment in patients with faecal incontinence: Statistical analysis plan for a randomised controlled trial, *Contemporary Clinical Trials Communications* (2017), doi: 10.1016/j.conctc.2017.07.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title page

Supervised pelvic floor muscle training versus attention-control massage treatment in patients with faecal incontinence: statistical analysis plan for a randomised controlled trial

Author information:

Anja Ussing ^{a,b,*}, Inge Dahn ^c, Ulla Due ^d, Michael Sørensen ^c, Janne Petersen ^b, Thomas Bandholm ^{a,b,e}

^a Department of Physiotherapy- and Occupational Therapy, Copenhagen University Hospital
Copenhagen University Hospital, Hvidovre, Denmark, Kettegaards Allé 30, DK-2650 Hvidovre

^b Optimed, Clinical Research Center, Copenhagen University Hospital
Copenhagen University Hospital, Hvidovre, Denmark, Kettegaards Allé 30, DK-2650 Hvidovre

^c Department of Surgical and Medical Gastroenterology, Copenhagen University Hospital,
Hvidovre, Denmark, Kettegaards Allé 30, DK-2650 Hvidovre

^d Department of Obstetrics and Gynaecology, Herlev Hospital, Herlev Ringvej 75, DK-2730 Herlev,
Department of Occupational and Physical Therapy, Herlev Hospital, Herlev Ringvej 75, DK-2730
Herlev, Denmark

^e Physical Medicine & Rehabilitation Research – Copenhagen (PMR-C), Copenhagen University
Hospital, Hvidovre, Denmark, Kettegaards Allé 30, DK-2650 Hvidovre,

* Corresponding author: Department of Physiotherapy- and Occupational Therapy, Optimed,
Clinical Research Center, Copenhagen University Hospital
Copenhagen University Hospital, Hvidovre, Denmark, Kettegaards Allé 30, DK-2650, anja.ussing@regionh.dk

Abstract

Introduction: Faecal incontinence affects approximately 8-9% of the adult population. The condition is surrounded by taboo; it can have a devastating impact on quality of life and lead to major limitations in daily life. Pelvic floor muscle training in combination with information and fibre supplements is recommended as first-line treatment for faecal incontinence. Despite this, the effect of pelvic floor muscle training for faecal incontinence is unclear. No previous trials have investigated the efficacy of supervised pelvic floor muscle training in combination with conservative treatment and compared this to an attention-control massage treatment including conservative treatment. The aim of this trial is to investigate if 16 weeks of supervised pelvic floor muscle training in combination with conservative treatment is superior to attention-control massage treatment and conservative treatment in patients with faecal incontinence.

Download English Version:

<https://daneshyari.com/en/article/8519376>

Download Persian Version:

<https://daneshyari.com/article/8519376>

[Daneshyari.com](https://daneshyari.com)