



Brazilian Journal
of Pharmacognosy

REVISTA BRASILEIRA DE FARMACOGNOSIA

www.elsevier.com/locate/bjp



Original Article

Development and clinical efficacy evaluation of anti-greasy green tea tonner on facial skin

Pathompong Meetham^a, Mayuree Kanlayavattanukul^{a,b}, Nattaya Lourith^{a,b,*}

^a School of Cosmetic Science, Mae Fah Luang University, Chiang Rai, Thailand

^b Phytocosmetics and Cosmeceuticals Research Group, Mae Fah Luang University, Chiang Rai, Thailand

ARTICLE INFO

Article history:

Received 19 November 2017

Accepted 16 January 2018

Available online xxx

Keywords:

Formulation

Stability

Green tea

Sebum

Tonner

Phytocosmetics

ABSTRACT

Green tea (*Camellia sinensis* (L.) Kuntze, Theaceae) polyphenols have activities against skin disorders; however, anti-sebum efficacy of green tea cosmetic has sparsely been reported. The facial tonner containing green tea was therefore developed and clinically evaluated. The base formulas which stabled following accelerated tests were sensorially assessed in ten volunteers. The base with hydroxyethyl cellulose, glycerin and panthenol (totally 3.6%) with the significant ($p < 0.05$) preference ($82.3 \pm 0.55\%$) over the others was further developed to green tea preparations. All of the products were stable and caused none of skin irritation as closed patch tested in twenty volunteers. Thereafter, they were clinically evaluated in the same group of the volunteers and monitored with Sebumeter[®] by means of a split-face, randomized single-blind, placebo-controlled study. Anti-greasy efficacy of 2, 4.5 and 7% green tea tonners were 3.47 ± 0.10 , 8.18 ± 0.44 and $17.87 \pm 0.46\%$ following 14 days of facial treatment. The efficiency was pronounced at the end of the study; day 28 (8.48 ± 0.13 , 20.26 ± 1.03 and $31.57 \pm 1.22\%$). Anti-sebum efficacy of the 4.5 and 7% green tea tonners were significantly better than the base formula (day 14; $p < 0.05$, day 28; $p < 0.01$). The efficacy of 28 days treatment was significantly better than 14 days ($p < 0.05$). The safe and efficient green tea tonner for oily face treatment was therefore approved in this context.

© 2018 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Green tea or *Camellia sinensis* (L.) Kuntze, Theaceae, is widely used for anti-aging (Elmets et al., 2001; Kartiyar and Elmets, 2001) and hydrating (Aburjai and Natsheh, 2003; Hsu, 2005) cosmetics either as the single active or in a combination with others (Chuarienthong et al., 2010). In addition, green tea possesses inhibitory effects against 5α -reductase (Lourith and Kanlayavattanukul, 2013) and androgen esthetic disorders. It is therefore applicable for body and oral malodor treatments (Lourith and Kanlayavattanukul, 2010; Kanlayavattanukul and Lourith, 2011) accordingly.

Oily skin usually refers to a facial condition with a shiny appearance and greasiness and heaviness due to excessive sebum production and secretion. This esthetic problem leads skin disorders such as acne, and significantly adverse quality of life (Wu et al., 2013) consequently. Thus, an application of cosmetics to reduce greasiness of skin is of challenge. Green tea facial tonner was therefore developed. The stable preparations were preference

assessed and skin irritation by means of a single closed patch test was conducted as consequently. A split-face, randomized single-blind, placebo-control clinical evaluation of the green tea tonners was thereafter examined with Sebumeter[®].

Materials and methods

Formulation and stability evaluation

Base tonners containing gelling agent selected from xanthan gum, carbopol and hydroxyethyl cellulose at different proportions, and tween 80, glycerine, panthenol, PEG/PPG-18/18 dimethicone, triethanolamine, preservative (Namsiang, Thailand) and water were formulated. Those with hydroxyethyl cellulose afforded the clear color less tonner with a preferred texture were therefore developed as shown in Table 1. pH of the tonners were determined (QIS, B200, the Netherlands). The obtaining three base formulas that were stable following accelerated stability test by means of a centrifugation assay ($805 \times g$ for 30 min) using Micromax RF (Thermo, USA), and six cycles of acceleration test (heating and cooling at 4°C and 45°C for 48 h, each cycle) (Chairsripipat et al., 2015), were included for preference test. The preferred tonner base

* Corresponding author.
E-mail: nattayal@mfu.ac.th (N. Lourith).

<https://doi.org/10.1016/j.bjp.2018.01.001>

0102-695X/© 2018 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Development of the stable green tea tonners.

Ingredient	Base (% w/w)			Green tea tonner (% w/w)		
	A	B	C	B1	B2	B3
Hydroxyethyl cellulose	0.1	0.1	0.1	0.1	0.1	0.1
Glycerin	1	3	4	3	3	3
Panthenol	↑	↑	↑	↑	↑	↑
PEG/PPG-18/18 dimethicone	3.5	3	4.5	3	3	3
Tween 80	↓	↓	↓	↓	↓	↓
Preservative	0.5	0.5	0.5	0.5	0.5	0.5
Triethanolamine	0.2	0.2	0.4	0.2	0.2	0.2
Water	94.7	93.2	90.5	91.2	88.7	86.2
Green tea extract	–	–	–	2	4.5	7
pH						
Int.	5.51 ± 0.03	5.55 ± 0.05	5.56 ± 0.02	5.56 ± 0.01	5.61 ± 0.02	5.51 ± 0.01
HC	5.60 ± 0.02	5.62 ± 0.03	5.62 ± 0.03	5.61 ± 0.01	5.67 ± 0.03	5.62 ± 0.09
Preference (%)						
Spreadability	88 ± 1.03	94 ± 0.96	80 ± 1.33			
Skin absorption	78 ± 1.47	88 ± 1.03	70 ± 1.69			
Greasiness	84 ± 0.84	92 ± 1.03	72 ± 1.03	NA	NA	NA
Color	82 ± 1.47	94 ± 0.96	68 ± 1.93			
Odor	60 ± 0.94	66 ± 0.96	58 ± 1.47			
Overall preference (%)	77.6 ± 0.63	82.3 ± 0.84	73.6 ± 10.3			

HC, heating and cooling; NA, not available.

was incorporated with green tea extract with 100 mg% of polyphenols (Product code: HE-EL11-CAS) supplied from Specialty Natural Products (Thailand). The obtaining green tea tonners were stability evaluated as above.

Preference test

Preference test for the base tonner was carried out in six female and four male Thai volunteers aged between 21 and 35 years old by means of the interview questionnaires with the hedonic system scoring from 1 to 5 (dislike – most prefer). The preferred base which passed the stability test was chosen for further development (Chaisripipat et al., 2015).

Clinical evaluation

Inclusion criteria

Thai healthy volunteers aged between 20 and 35 years old were enrolled in the study. The inclusion criteria for the volunteer was based on the facial sebum content between 180–250 and 160–250 $\mu\text{g}/\text{cm}^2$ for male and female subjects, respectively, as assessed by Sebumeter[®] (SM810, CK, Germany). All recruited subjects were informed about the study both in writing and verbally and signed a written consent form, which was approved by the ethical committee of Mae Fah Luang University prior to enrollment (REH-59083). All of the study in human volunteers was delineated by the Declaration of Helsinki.

Irritation test

Base tonner was comparatively examined with the green tea tonners. Water was used as a negative control, whereas 3% sodium lauryl sulfate (Namsiang) was a positive control. Observation was undertaken immediately, 24, 48 and 72 h following Finn chamber[®] (8 mm, SmartPractice, USA) removal. Mean Irritation Index (MII) was calculated (Schnuch et al., 2008).

Efficacy evaluation

Twenty healthy Thai volunteers (sixteen female and four male) aged between 20 and 35 years old were included. All subjects were allergy-free for one week, had not used steroids and topical products for oily or greasy face treatment for four weeks prior to study

enrollment. Subjects who were pregnant or lactating or dieting were excluded from the study as well as smoking and liquor drinking. The subjects were asked not to apply any products onto the face the night before starting the study. They were all facial greasiness determined with Sebumeter[®] at the baseline. The subjects were acclimatized in the waiting room ($20 \pm 1^\circ\text{C}$ and 40–60% relative humidity) for 30 min before the measurements were carried out under the same condition. A randomized single-blind placebo-controlled study was designed and the volunteers were requested to apply the tonner once in the morning by a split-face procedure for intra-individual comparative trails. The volunteers were directed to use ten drops of the tonner absorbed on the cotton pad (Ambulance, Thailand) on one face side and the different tonner on another side at the same quantity. The subject who used the assigned product less than 90% of the guideline was listed in the discontinuation criteria. The efficacy evaluation was monitored on days 14 and 28 by Sebumeter[®] under the same condition with the baseline. Anti-sebum efficacy was calculated as followings:

$$\text{Anti-sebum efficacy (\%)} = 100 - \left(\frac{St}{S0} \times 100 \right)$$

where

St = sebum content ($\mu\text{g}/\text{cm}^2$) at time interval
S0 = sebum content ($\mu\text{g}/\text{cm}^2$) at base line

Statistical analysis

The parameters were compared and analyzed using Post Hoc test for stability and preference tests. Independent sample t test was for efficacy evaluations. The significance was set at a reliability of 95%, and expressed as mean \pm SEM.

Results and discussion

Formulation, stability evaluation and preference test

The base formulation was firstly developed by a variation of gelling agents, at which hydroxyethyl cellulose was chosen for further development (Table 1) due to its appearance and texture. All of the tonners were clear, colorless liquid that stabled following

Download English Version:

<https://daneshyari.com/en/article/8542907>

Download Persian Version:

<https://daneshyari.com/article/8542907>

[Daneshyari.com](https://daneshyari.com)