


## Accepted Manuscript

Title: Establishment of a drug-induced rhabdomyolysis mouse model by co-administration of ciprofloxacin and atorvastatin

Authors: Akiko Matsubara, Shingo Oda, Sho Akai, Koichi Tsuneyama, Tsuyoshi Yokoi



PII: S0378-4274(18)30153-X  
DOI: <https://doi.org/10.1016/j.toxlet.2018.04.016>  
Reference: TOXLET 10164

To appear in: *Toxicology Letters*

Received date: 9-1-2018  
Revised date: 12-4-2018  
Accepted date: 14-4-2018

Please cite this article as: Matsubara, Akiko, Oda, Shingo, Akai, Sho, Tsuneyama, Koichi, Yokoi, Tsuyoshi, Establishment of a drug-induced rhabdomyolysis mouse model by co-administration of ciprofloxacin and atorvastatin. *Toxicology Letters* <https://doi.org/10.1016/j.toxlet.2018.04.016>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Establishment of a drug-induced rhabdomyolysis mouse model by  
co-administration of ciprofloxacin and atorvastatin**

Akiko Matsubara<sup>a</sup>, Shingo Oda<sup>a</sup>, Sho Akai<sup>a</sup>, Koichi Tsuneyama<sup>b</sup>, Tsuyoshi Yokoi<sup>a\*</sup>

<sup>a</sup>Department of Drug Safety Sciences, Division of Clinical Pharmacology, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya, 466-8550, Japan.

<sup>b</sup>Department of Pathology and Laboratory Medicine, Institute of Biomedical Sciences, Tokushima University Graduate School, 3-18-5 Kuramoto-cho, Tokushima, 770-8503, Japan.

\* **Corresponding author:** Tsuyoshi Yokoi, Ph.D.

Department of Drug Safety Sciences, Nagoya University Graduate School of Medicine, 65 Tsururui-cho, Showa-ku, Nagoya, 466-8550, Japan.

E-mail: [tyokoi@med.nagoya-u.ac.jp](mailto:tyokoi@med.nagoya-u.ac.jp)

Tel: +81-52-744-2110, Fax: +81-52-744-2114

## **HIGHLIGHTS**

- A rhabdomyolysis mouse model was established by co-administration of ciprofloxacin and atorvastatin.
- Drugs were orally administered for 4 days to the glutathione-depleted normal mouse.
- An increase of plasma creatinine phosphokinase (CPK) value and degeneration in

Download English Version:

<https://daneshyari.com/en/article/8553290>

Download Persian Version:

<https://daneshyari.com/article/8553290>

[Daneshyari.com](https://daneshyari.com)