

## Accepted Manuscript

Title: Epididymal phospholipidosis is a possible mechanism for spermatotoxicity induced by the organophosphorus insecticide fenitrothion in rats

Authors: Mio Miyake, Yuki Ito, Himiko Suzuki, Motohiro Tomizawa, Hirotaka Sato, Ming Liu, Ai Okamura, Tamie Nakajima, Katsumi Ohtani, Hisashi Takino, Hiroshi Inagaki, Michihiro Kamijima


PII: S0378-4274(17)31533-3  
DOI: <https://doi.org/10.1016/j.toxlet.2017.12.023>  
Reference: TOXLET 10055

To appear in: *Toxicology Letters*

Received date: 11-10-2017  
Revised date: 11-12-2017  
Accepted date: 27-12-2017

Please cite this article as: Miyake, Mio, Ito, Yuki, Suzuki, Himiko, Tomizawa, Motohiro, Sato, Hirotaka, Liu, Ming, Okamura, Ai, Nakajima, Tamie, Ohtani, Katsumi, Takino, Hisashi, Inagaki, Hiroshi, Kamijima, Michihiro, Epididymal phospholipidosis is a possible mechanism for spermatotoxicity induced by the organophosphorus insecticide fenitrothion in rats. *Toxicology Letters* <https://doi.org/10.1016/j.toxlet.2017.12.023>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Epididymal phospholipidosis is a possible mechanism for spermatotoxicity induced by the organophosphorus insecticide fenitrothion in rats

Mio Miyake<sup>a</sup>, Yuki Ito<sup>a,\*</sup>, Himiko Suzuki<sup>a</sup>, Motohiro Tomizawa<sup>a,b</sup>, Hirotaka Sato<sup>a</sup>, Ming Liu<sup>a</sup>, Ai Okamura<sup>c</sup>, Tamie Nakajima<sup>c,d</sup>, Katsumi Ohtani<sup>e</sup>, Hisashi Takino<sup>f</sup>, Hiroshi Inagaki<sup>f</sup>, and Michihiro Kamijima<sup>a,\*</sup>

<sup>a</sup> *Department of Occupational and Environmental Health, Nagoya City University Graduate School of Medical Sciences, Nagoya 467-8601, Japan*

<sup>b</sup> *Department of Chemistry and Food Safety, Faculty of Life Sciences, Tokyo University of Agriculture, Tokyo 156-8502, Japan*

<sup>c</sup> *Nagoya University Graduate School of Medicine, Nagoya 466-8550, Japan*

<sup>d</sup> *College of Life and Health Sciences, Chubu University, Kasugai 487-8501, Japan*

<sup>e</sup> *National Institute of Occupational Safety and Health, Kawasaki 214-8585, Japan*

<sup>f</sup> *Department of Pathology and Molecular Diagnostics, Nagoya City University Graduate School of Medical Sciences, Nagoya 467-8601, Japan*

\* Corresponding authors.

*E-mail addresses:* [yukey@med.nagoya-cu.ac.jp](mailto:yukey@med.nagoya-cu.ac.jp) (Y. Ito), [kamijima@med.nagoya-cu.ac.jp](mailto:kamijima@med.nagoya-cu.ac.jp) (M. Kamijima).

Running title: Epididymal phospholipidosis induced by fenitrothion in rats.

*Abbreviations:* 2-AG, 2-arachidonoylglycerol;  $\beta$ , standardized partial regression coefficient; FAAH, fatty acid amide hydrolase; FNT, fenitrothion; MNP, 3-methyl-4-nitrophenol; OP, organophosphate;  $R^2$ , coefficient of determination adjusted for the degrees of freedom; sPLA2,

Download English Version:

<https://daneshyari.com/en/article/8553424>

Download Persian Version:

<https://daneshyari.com/article/8553424>

[Daneshyari.com](https://daneshyari.com)