

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/TJEM)

Turkish Journal of Emergency Medicine

journal homepage: <http://www.elsevier.com/locate/TJEM>

Original Article

Analysis of mean platelet volume and red blood cell distribution width in recurrent epistaxis

Ali Ekber Karabulut ^a, Yunsur Çevik ^b, Emine Emektar ^b, Şeref Kerem Çorbacıoğlu ^b, Seda Dağar ^{b,*}, Oğuz Yardim ^c

^a Emergency Department, Van Training and Research Hospital, Van, Turkey

^b Department of Emergency Medicine, Keçioren Training and Research Hospital, Ankara, Turkey

^c Emergency Department, Bayburt State Hospital, Bayburt, Turkey

ARTICLE INFO

Article history:

Received 11 December 2017

Received in revised form

9 February 2018

Accepted 11 February 2018

Available online xxx

Keywords:

Emergency department

Mean platelet volume

Recurrent epistaxis

Red blood cell distribution width

ABSTRACT

Objectives: Recurrent epistaxis is one of the most common causes of emergency department visits. Although several localized and systemic conditions has been described, the exact cause is unknown in the majority of cases. In our study, we aimed to determine the effect of mean platelet volume (MPV) and red blood cell distribution width (RDW) levels on recurrent epistaxis.

Method: One hundred and thirty six patients with recurrent epistaxis and 170 healthy cases as control group were included in the study. Demographic data, vital signs and the results of complete blood counts were recorded. The patients who had clinical conditions that might affect the levels of MPV or RDW, were excluded. MPV and RDW levels were compared between the two groups.

Results: The median level of MPV was 7.6 fL (IQR25–75%: 7.1–8.4) in the study group and 8.2 fL (IQR25–75%: 7.8–8.9) in the control group ($p < 0.001$). The median levels of RDW were found in the patient and control groups in order %15.4 (IQR25–75%: 14.5–15.4) and %14.3 (IQR25–75%: 13.4–15.4) ($p < 0.001$). Systolic blood pressure, leukocyte count, age, sex, RDW and MPV levels that were variables with p levels < 0.2 , were included in the multivariate analyses. It was determined that high RDW levels increased epistaxis ($p < 0.001$; OR: 1.89 [95% CI: 1.53–2.33]) and high MPV levels decreased epistaxis ($p < 0.001$; OR: 0.54 [95% CI: 0.39–0.72]).

Conclusion: Low MPV and high RDW levels caused an increased bleeding tendency in patients with recurrent epistaxis. Although exact mechanism is not known, referring those patients for the assessment of etiologic causes would be proper.

Copyright © 2018 The Emergency Medicine Association of Turkey. Production and hosting by Elsevier B.V. on behalf of the Owner. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Epistaxis is one of the most common causes of bleeding which is managed and treated in emergency departments frequently. Although it has been described several localized conditions (such as digital trauma, allergic rhinitis, nasal septal deformity, septal perforations, nasal sprays with antihistaminic or steroid, nasal foreign bodies, aneurysm, nasal papilloma, adenocarcinoma) and systemic conditions (such as hypertension, clotting disorders, bleeding

diathesis, vascular and cardiovascular diseases), the cause of epistaxis are unknown in the majority of cases.¹

Several studies have been published recently that research whether there are any relationships between epistaxis and various blood parameters, the most obvious of which are mean platelet volume (MPV) and red blood cell distribution width (RDW).^{2–4} It was revealed that high MPV values lower the frequency of epistaxis.² Platelets are made up of megakaryocytes in the form of a disc and responsible for coagulation. MPV value is one of the important parameters that determine the function of platelet. MPV is an important variable, and in metabolic and enzymatic aspects, big platelets were shown to be more active than small platelets.⁵ Big platelets involve more prothrombic factors, such as thromboxane A₂, beta-thromboglobulin, and adhesion molecules. Thromboxane A₂ functions in the aggregation of vasoconstrictors

* Corresponding author. Keçiören Training and Research Hospital, Emergency Department, Ankara, Turkey.

E-mail address: sedadagar@gmail.com (S. Dağar).

Peer review under responsibility of The Emergency Medicine Association of Turkey.

<https://doi.org/10.1016/j.tjem.2018.02.001>

2452–2473/Copyright © 2018 The Emergency Medicine Association of Turkey. Production and hosting by Elsevier B.V. on behalf of the Owner. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

and platelets.^{6–8} Increased MPV value is associated with shortened bleeding period.⁹ It has also been reported that increase in platelet volume is associated with acute myocardial infarction (AMI), acute cerebral ischemia and transient ischemic attacks.^{6–8} RDW value is a criterion for the variability in the erythrocyte population.¹⁰ Increased RDW can be seen in a large number of diseases which cause reticulocytes to be released into circulation before maturation. As well as being the distinctive diagnosis of anemia, increased RDW values were found to be associated with heart failure, cardiovascular events, celiac disease and inflammatory intestinal disease activity.¹¹ Also, increase in RDW level was determined to be associated with increased inflammatory markers.^{12,13}

In this study, our purpose is to find out whether there is any relationships between recurrent epistaxis and MPV and RDW values.

2. Material and methods

2.1. Study design

This retrospective, case-control study was conducted by examining the files of the patients who were presented to the Emergency Department of Keçiören Training and Research Hospital between November 15, 2014 and November 15, 2015 due to epistaxis. Approval from the local ethical committee was obtained before the study began. The presentation of the patients to hospital with epistaxis at least twice during the previous year was considered as recurrent epistaxis. Demographic data of these patients, their vital signs and the results of their complete blood count were recorded. The patients aged under 18 and over 65, those with traumatic epistaxis, pregnant women, the patients with diseases that may lead to changes in MPV and RDW parameters (diabetes mellitus, hypertension, anemia, bleeding disorders, cardiovascular diseases, stroke, peripheral vascular diseases, acute infections, thyroid disorders, chronic inflammatory skin diseases, malignancy and chronic drug use) were excluded from the study. The control group was composed of healthy individuals at similar ages and with similar genders, presented to the primary care physician for routine control.

2.2. Statistical analyses

Statistical analyses were performed using SPSS version 15.0 (SPSS, Inc., Chicago, IL, USA). The Shapiro-Wilk test was used to assess the normal distribution of the variables. The data consistent with the normal distribution were expressed as mean \pm standard deviation (SD), and those that were not consistent were expressed as median and 25% and 75% quarters (IQR). Pearson chi-square test was used for the comparison of categorical data between the groups, Student-T test was used for the comparison of parametric continuous data and Mann-Whitney-U test was used for non-parametric data. The variables with a p value under 0.2 as well as age and gender were included in the model for a multivariable regression analysis. Of these variables, the ones that could correlate with each other were evaluated with Spearman correlation test. The suitability of the model was assessed with Hosmer-Lemeshow test. $p < 0.05$ was considered statistically significant.

3. Results

In total, 136 patients and 170 control cases were included in the study. 76 patients (55.3%) were male in the patient group, while there were 94 males (55.3%) in the control group. The median age was 38.5 (IQR25–75%: 23.0–51.0) in the patient group and 35 (IQR25–75%: 27.0–46.2) in the control group. No statistically

significant difference was detected between the groups ($p = 0.6$). Demographic data and vital signs of the patients are shown at Table 1.

When whole blood count parameters were examined, MPV median value was found to be 7.6 fL (IQR25–75%: 7.1–8.4) in the patient group and 8.2 fL (IQR25–75%: 7.8–8.9) in the control group. There was a statistically significant difference between the groups ($p < 0.001$). Similarly, RDW median value was 15.4% in the patient group (IQR25–75%: 14.5–15.4) and 14.3% in the control group (IQR25–75%: 13.4–15.4). There was also a statistically significant difference between the groups ($p < 0.001$). The other whole blood count parameters which a statistically significant difference was detected between the groups were hemoglobin ($p = 0.010$), hematocrit ($p = 0.002$) and MCV ($p < 0.001$). No statistically significant difference was found between the other variables (Table 2). A multivariable regression analysis was conducted in order to determine the effects of these parameters on epistaxis. Systolic blood pressure, leukocyte, RDW, MPV values with $p \leq 0.2$ as well as age and gender were included in this multivariable analysis. It was found that a high RDW value increased epistaxis ($p < 0.001$; OR:1.89 [95% CI:1.53–2.33]) and a high MPV value decreased it ($p < 0.001$; OR:0.54 [95% CI:0.39–0.72]) (Tables 3 and 4).

4. Discussion

The most important result of this study is that MPV value in the patients who were presented to the emergency department with epistaxis, was lower than that of the healthy controls, whereas RDW value in patients with epistaxis was higher. We found that a high MPV value decreased epistaxis by 0.54-fold (OR:0.54 [95% CI:0.39–0.72]), while a high RDW value increased epistaxis by 1.89-fold (OR:1.89 [95% CI:1.53–2.33]). We believe that a low MPV value and a high RDW value could be predictive for recurrent epistaxis.

RDW is an indicator showing the variability of the sizes of red blood cells in circulation. The increase in RDW value, which traditionally suggests the presence of anemia, in fact shows that the heterogeneity of erythrocyte size in the peripheral blood. Consequently, studies have been conducted to show that RDW value is associated with several vascular incidents. In a study by Çetin et al., the patients with stable coronary arterial disease and the cases with normal coronary arteries in their angiography were compared, and they reported that a high RDW value was associated with the spread and severity of coronary arterial disease.¹⁴ In another study, a correlation between a high RDW value and atherosclerosis of carotid artery was established in patients with hypertension.¹⁵ Some of the researchers found that a high RDW value could be a reflection of chronic inflammation that might lead to vascular incidents.^{12,13} In other studies, the existence of a correlation between celiac disease and the activity of inflammatory intestinal disease gives support to the idea that a high RDW value might hint the presence of inflammation.^{11,16} It was noted that the effect of inflammation on erythropoiesis and the half-life of erythrocytes in circulation could lead to a high RDW.¹⁷ The only study which identified lower RDW values in patients with recurrent epistaxis than in normal population, belongs to Kemal et al.² However, in contrast to this result, we found that high RDW values increased epistaxis episodes by 1.89-fold ($p < 0.001$; OR:1.89 [95% CI:1.53–2.33]) in our study. Just as in the recurrent hidden gastrointestinal bleedings in colorectal carcinomas, an increased blood loss that might result from recurring epistaxis attacks could increase erythropoiesis, which might be the reason why we identified high RDW values in these patients.

MPV is one of the most important markers to show the function and activity of platelets. Increased MPV value is associated with shortened bleeding period.⁹ Big platelets are more active than small

Download English Version:

<https://daneshyari.com/en/article/8554234>

Download Persian Version:

<https://daneshyari.com/article/8554234>

[Daneshyari.com](https://daneshyari.com)