



Correlation between parent-adolescent communication and adolescents' premarital sex risk

Widyatuti*, Citra Hafilah Shabrina and Astuti Yuni Nursasi

Faculty of Nursing, Universitas Indonesia, Depok, Jawa Barat, Indonesia

KEYWORDS

Parents;
Adolescents;
Communication;
Parent-Adolescent
Communication
Scale;
Premarital sex

Abstract

Objective: Previous studies have indicated the parent-adolescent relationship has a correlation to adolescents' premarital sex behavior risk. Therefore, the aim of this study was to discover the influence of parent-adolescent communication on adolescents' risk of sexual issues.

Method: This was a quantitative study with a cross-sectional design. The population of this study consisted of students from a high school in Jakarta. A purposive sampling technique was used, which resulted in the selection of 253 students as samples. A PACS (Parent-Adolescent Communication Scale) questionnaire was applied.

Results: The results showed that 59.3% of the adolescents studied were at risk for engaging in premarital sex, while the risk for adolescents with positive communication with their parents was 56.5%. Bivariate analysis also showed a significant correlation between gender and parent-adolescent communication and the risk of adolescent premarital sex behavior ($\alpha < 0.05$). This study's result was expected, and nurses can follow up by providing health education on how to maintain good communication between parents and adolescents.

Conclusions: Communication must align with adolescents' developmental tasks. Nurses can also create a promotion program on the topic of communication for parents and adolescents.

© 2018 Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Adolescents account for a large portion of the population in Indonesia. BKKBN explained that, in 2010, the population of Indonesia was 237.6 million, and 26.67% of it was adolescents¹. Adolescence is the most vulnerable period in an individual's life span because of the many physical, cognitive, and psychosocial² changes that occur at this stage. Indeed, adolescence marks the transition from childhood to adulthood. During this stage, individuals expand their social relationships, so their usual daily interactions can affect their lives. These changes that occur during adoles-

cents' growth and development make them vulnerable to many problems.

The problems that can occur during adolescence are many and include emotional disorders and suicide, violence, substance abuse, pregnancy, sexually transmitted infections, dermatological issues, nutritional issues such as obesity or deficiency of micronutrient, and premarital sex. Arist said that research by the Child Protection National Commission in 2012 showed that 93.7% of 4,726 respondents had engaged in premarital sex, and 21.26% had undergone abortions³. The study showed an increase from 2008, when 62.7% had reported having engaged in premarital sex, and 21.2% had undergone abortion. In Jakarta,

*Corresponding author.

Email: tuti_cw@yahoo.com (Widyatuti).

51% of adolescents reported having engaged in premarital sex in 2014⁴. Based on that number, the number of adolescents engaging in premarital sex is still high and increasing. This increase may have an effect on the rates of adolescent pregnancy, early marriage, and maternal and infant mortality.

Premarital sex is sex before or without marriage. Some adolescent sexual behaviors, such as dating, may be a precursor to sex⁵. Sexual behavior has some steps, and we need to understand these steps to determine the risk of adolescents engaging in premarital sex. Duvall and Miller identified the stages of sexual behavior as physical touch, kissing, flirting, and sex⁶. There are some factors that may contribute to sexual activity in adolescents: if they are part of a couple or if they have a friend who supports the idea of premarital sex⁷. Some of the factors that affect adolescents' choices about premarital sex include education, reproductive knowledge, personal attitude, media exposure, societal pressures, parents, school, and peers⁸. Parents act as a main system of support, educators, role models, supervisors, friends, counselors, and communicators⁹. Thus, parents can have an impact on adolescents' premarital sex risk. Based on previous research, bad family relationships may increase the risk of premarital sex to around 72.7%¹⁰. Barnes and Olson explained that family relationships may also have an impact on the communication between parents and adolescents¹¹.

Previous research explains there is a correlation between the quality of parent-adolescent communication and premarital sex among adolescents^{12,13}. However, there is also research that explains there is no correlation between parent-adolescent communication about sexuality and premarital sex among adolescents¹⁴. Based on Olson theory about family correlation and Anwar's research result about premarital sex factor, there should be a correlation between parent-adolescent communication and premarital sex among adolescents. The differences between Anwar's and this research were populations studied. Therefore, there is a need for conducting research related to the connection between parent-adolescent communication and premarital sex among adolescents within population. This population have good school environment that is school with minimum risk or no student smoking, supportive peers and no premarital sex history. The hypothesis of this research is that there is a correlation between parent-adolescent communication and adolescents' premarital sex risk.

Method

This quantitative research applied a cross-sectional design with independent t-test to determine if a correlation exists between parent-adolescent communication, and the risk of premarital sex among adolescents. The population studied in this research consisted of students at one senior high school at Jakarta, with a total student body of 1,060. The researcher used the Slovin formula to set the sample number, with class X and XI as a population. Purposive sampling will be applied to 253 participants. The inclusion criteria are willing to be participated, and living with parents.

The variables in this study included parent-adolescent communication (numeric) and premarital sex risk (category). The instrument was a questionnaire, which consisted of three parts: a) demographics such as age and gender; b) risk of premarital sex, and c) the David Olson Parent-Adolescent Communication Scale which has 2 aspects (positive communication such as honesty to parents and negative communication such as avoiding some topics with parents), ten questions of which had been modified with the Likert Scale (a five choice scale ranging from strongly disagree to strongly agree).

The data collection procedure began with a research proposal letter for collecting data that was then submitted to the school. The researcher met the teacher and some students, who helped the researcher to identify suitable respondents. After that, the researcher met respondents and explained the aim of the study and also obtained informed consent before handing out the questionnaires. The researcher then collected the questionnaires and ensured that each item had been answered in full. Univariate and bivariate analysis was performed with the SPSS program. In terms of ethical considerations, the researcher obtained permission from the school and the respondents to conduct the study. All results study were kept privately, only researcher who can access to the data.

Results

The results of this study indicated that the adolescents surveyed have a high risk of engaging in premarital sex. Of the total respondents, 150 (59.3%) are at risk for engaging in premarital sex, and 103 (40.7%) are not at risk. Parent-adolescent communication was found mostly positive. According to data analysis, 110 respondents (43.5%) received a score lower than 34, indicating a negative communication pattern between parents and adolescents. In contrast there were 143 respondents (56.5%), who reported having good communication with their parents such as tell their feeling to their parents voluntarily.

Further analysis showed that there is no correlation between age and risk of premarital sex. This can be seen from the p value of 0.37 or more than α with Levene's test with the same variant (Table 1).

The results from further analysis between gender and the risk of premarital sex indicate a correlation exists between the two. This can be seen from the p value 0.005 or less than α with Levene's test with a different variant (Table 2).

A correlation was also identified between parent-adolescent communication and the risk of premarital sex. This is can be seen from the p value of 0.021 or less than α with Levene's test with a different variant (Table 3).

Risks of premarital sex	Mean	Standard deviation	P value
Risk	16.16	0.795	0.37
No risk	16.07	0.808	

Download English Version:

<https://daneshyari.com/en/article/8568863>

Download Persian Version:

<https://daneshyari.com/article/8568863>

[Daneshyari.com](https://daneshyari.com)