Saudi Journal of Ophthalmology (2017) xxx, xxx-xxx

Original Article

Knowledge and practice regarding contact lens among Saudi urban contact lens users

Omar S. Alobaidan ^{b,*}; Mohammed K. Alkhalifah ^b; Ammar A. AlSayegh ^b; Fahad A. Alhumaid ^b; Ahmad S. Ashammery ^b; Khalid Alghamdi ^b; Ahmed Mousa ^c; Rajiv Khandekar ^d; Waleed AlRashid ^a

Abstract

Purpose: To assess the level and determinants of knowledge and practice regarding contact lens (CL) and its accessories among adult Saudi CL users.

Methods: This survey was conducted in 2015 for Saudi contact lens users visiting a Mall in Riyadh, Saudi Arabia. The questions were related to knowledge about contact lenses, correct CL and accessories usage. We inquired indication, initial prescription, CL use, complications experienced and their management. The knowledge and practice responses were correlated to their determinants.

Results: In this survey, 510 contact lens users participated. The level of knowledge was of excellent grade among 279 [54.7% (95% Confidence Interval 50.4–59)] CL users. The practice was of excellent grade in 210 [41.4% (95% CI 37.1–45.7)] CL users. Variation in knowledge was not associated to gender (P = 0.1), education (P = 0.4), type of work (P = 0.3), funding for CL (P = 0.1). Occasional users and those not having prescribed CL had significantly poor knowledge (P < 0.001). The excellent level of practice was associated to younger CL users (P = 0.004). While it was not associated to gender (P = 0.8), type of CL (P = 0.9), funding for CL (P = 0.2), education (P = 0.7) and occupation (P = 0.1).

Conclusions: The knowledge and practice among CL users was less than desired. Those using contact lens for cosmetic purpose and procuring them without prescription need special focus for health promotion so that their CL related practice improve and eye complications are reduced.

Keywords: Contact lens, Knowledge and practice, Abuse of contact lens, Survey

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of Saudi Ophthalmological Society, King Saud University.

This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

https://doi.org/10.1016/j.sjopt.2017.09.008

Introduction

Contact lenses (CL) are widely used for cosmetic purpose, in addition to correction of refractive error. ^{1,2} They are dispensed by qualified contact lens practitioners and are sold

as 'over the counter product' in many countries. Serious eye complications like keratitis and endophthalmitis, are often not explained to CL users at the time of dispensing. Minor complications like conjunctival hyperaemia and papillae formation causes discomfort and may need attention by

Received 30 January 2017; received in revised form 12 June 2017; accepted 25 September 2017; available online xxxx.

- ^a Department of Ophthalmology, Al Imam Mohammad Ibn Saud Islamic University, College of Medicine, Riyadh, Saudi Arabia
- ^b Al Imam Mohammad Ibn Saud Islamic University, College of Medicine, Riyadh, Saudi Arabia
- ^c Ocular Epidemiology at King Saud University, Saudi Arabia
- ^d King Khaled Eye Specialist Hospital, Riyadh, Saudi Arabia
- * Corresponding author at: P.O. Box 245, Riyadh 11411, Saudi Arabia. e-mail address: omaralobaidan@gmail.com (O.S. Alobaidan).





Peer review under responsibility of Saudi Ophthalmological Society, King Saud University



Access this article online: www.saudiophthaljournal.com www.sciencedirect.com 2 O.S. Alobaidan et al.

caregiver during periodic follow-ups. Many contact lens wearers face sight-threatening eye complications because of not taking proper care or poor compliances with the advices given by the eye care professionals.^{3,4}.

The knowledge about CL, its accessories and eye care among contact lens users therefore is very crucial. Even among health staff using CL, the level of knowledge and compliance to good practice was less than desired.⁵ A study of Saudi female university students, 70% of interviewed students were using CL and two third of them were using them for cosmetic purpose. As many as 39% of them were using it without consulting eye care professionals.⁶

To the best of our knowledge, the practice among adult Saudi CL users including males has not been evaluated. Therefore, we surveyed adult Saudi population. We used a questionnaire as survey tool to assess level of knowledge about care of CL and their accessories, type of CL usage practice among them and untoward events taking place due to abuse of contact lens in Riyadh, Saudi Arabia.

Material and methods

This survey was approved by the ethical and research committee of our institute. Informed verbal consent was obtained from each participant. Adult Saudi population of Riyadh city and using contact lenses coming to the eye health promotion campaign that was held in Granada Mall, Riyadh, Saudi Arabia during the period from 29 th to 31 st of January 2015, were our study population.

We assumed based on annual supply of CL by international brand CL companies that there could be as many as 50,000 adult contact lens wearers could be in the study area. The strict protocol for hygiene is followed among 75% of them.⁶ For a survey with 95% confidence interval, 5% acceptable error margin and design effect factor of 2, we need at least 441 participants. To compensate for not completing the entire survey, we increased the sample by 15%. Thus final sample size for the present study was 507. We used STAT calculator of Open-EPI software for calculating the sample of a cross sectional study.⁷

A pretested data collection form included demographics, questions related to knowledge about CL and its accessories and practice of CL usage. The demography included agegroup, gender, education level and type of work for CL wearers. The inquiry also included source of purchase of CL, indication for CL usage, source of information regarding CL, frequency and type of CL usage.

There were six questions related to knowledge about CL and allied products. There were 10 questions related to practice regarding hygiene, CL and accessories care, need for seeking urgent and routine eye care for discomfort or acute eye complications they experienced. The correctness of response was determined by comparing to the reply of two experienced contact lens practitioners (Gold standard). If the reply of participant matched to the expert's response, one point was given. If it did not match '0' score was given. For knowledge related questions, there was maximum 14 score. If score of each participant was more than 10, we considered it to be excellent grade of knowledge. For practice related responses, the maximum score was 10 and the minimum score was 0. If the overall score of participant was seven

or more, we considered the level of practice as of 'excellent' grade.

The survey data was compiled in spreadsheet of Microsoft XL^{\circledast} . The outcome variable of excellent grade of knowledge and excellent grade of practice was compared with different epidemiological and CL related variables. For qualitative variables, we calculated frequencies and percentage proportions. The outcome variables were also presented as percentage with its 95% confidence interval. For Quantitative variable, we first plotted distribution curve and if it was normal, we calculated mean and standard deviation. To compare outcome variable in two independent variables we used 2×2 tables and calculated Odds ratio and two sided P value. For more than 2 variables, we calculated chi-square value, degree of freedom and two sided P values. The P value of <0.05 was considered as significant.

Results

We surveyed 510 adult CL users. Their demographic profile suggested that more than half of the CL users were of less than 25 years of age. Male constituted less than 10% of total CL users. Nearly two-third of participants had college education. Most of the CL users were spending their own money for CL related products.

The information about the contact lenses being used by participants is given in Table 1. Six out of ten use them for cosmetic purpose. As many as one fifth had stopped using CL. Daily wear soft lenses were the main type of CL used by participants. Only one fourth of CL users received information regarding CL from eye care professionals.

The level of knowledge was of excellent grade among 279 [54.7% (95% Confidence Interval 50.4–59)]. The knowledge about CL was self-perceived in 312 (61.2%) of participants. 40% of the participants were overestimating their level of knowledge regarding CL. The knowledge regarding possible complications due to abuse of contact lenses was very poor in 26.2% of CL users. Nearly 45% of CL users were not knowing contraindications of CL usage.

Table 1. Information about contact lens used by Saudi participants.

		Number	Percentage
Indication for contact lens use	Medical Cosmetic Medical + cosmetic	137 297 76	26.9 58.3 14.8
Type of CL usage	All time Part time Not using now	91 324 85	17.8 63.5 16.7
Type of lens using	Soft Semisoft Hard Don't know	440 29 21 20	86.3 5.7 4.1 3.9
Source of cl information	Doctor Optometrist Vendor Friends Family member Internet Brochure Self Other	63 68 130 26 31 108 64 11	12.4 13.3 25.5 5.1 6.1 21.2 12.5 2.2 1.8

Download English Version:

https://daneshyari.com/en/article/8591893

Download Persian Version:

https://daneshyari.com/article/8591893

<u>Daneshyari.com</u>