

Accepted Manuscript

Title: Resting Functional Connectivity of the Periaqueductal Gray is Associated with Normal Inhibition and Pathological Facilitation in Conditioned Pain Modulation

Author: Daniel E. Harper, Eric Ichesco, Andrew Schrepf, Johnson P. Hampson, Daniel J. Clauw, Tobias Schmidt-Wilcke, Richard E. Harris, Steven E. Harte


PII: S1526-5900(18)30024-5
DOI: <https://doi.org/10.1016/j.jpain.2018.01.001>
Reference: YJPAI 3517

To appear in: *The Journal of Pain*

Received date: 19-6-2017
Revised date: 18-12-2017
Accepted date: 3-1-2018

Please cite this article as: Daniel E. Harper, Eric Ichesco, Andrew Schrepf, Johnson P. Hampson, Daniel J. Clauw, Tobias Schmidt-Wilcke, Richard E. Harris, Steven E. Harte, Resting Functional Connectivity of the Periaqueductal Gray is Associated with Normal Inhibition and Pathological Facilitation in Conditioned Pain Modulation, *The Journal of Pain* (2018), <https://doi.org/10.1016/j.jpain.2018.01.001>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Resting functional connectivity of the periaqueductal gray is associated with normal inhibition and pathological facilitation in conditioned pain modulation

Authors: Daniel E. Harper^{1*}, Eric Ichesco¹, Andrew Schrepf¹, Johnson P. Hampson¹, Daniel J. Clauw¹, Tobias Schmidt-Wilcke^{2,3}, Richard E. Harris^{1,4}, Steven E. Harte^{1,4}

¹Chronic Pain and Fatigue Research Center, Department of Anesthesiology, University of Michigan, Ann Arbor, MI, USA

²Department of Neurology, St. Mauritius Therapieklinik, Meerbusch, Germany

³Department of Clinical Neuroscience and Medical Psychology, Medical Faculty, Heinrich-Heine-University, Düsseldorf, Germany

⁴Equal author contribution

*Corresponding author

Daniel E Harper, Ph.D.

Chronic Pain and Fatigue Research Center

Department of Anesthesiology

24 Frank Lloyd Wright Drive

PO Box 385, Lobby M

Ann Arbor, MI 48106

Phone: (734)-998-0176

Fax: (734)-998-6900

Email: harperd@med.umich.edu

www.med.umich.edu/painresearch

4 Tables; 6 Figures; 2 Supplemental Figures

Disclosures: This was an investigator initiated study funded by Forest Laboratories (MD-SAV-09). Dr. Clauw has consulted for Forest Laboratories, Pfizer, Inc., Cerephex Corporation, Eli Lilly and Company, Merck & Co., Nuvo Research Inc., Tonix Pharmaceuticals, Johnson & Johnson, Pierre Fabre, Cypress Biosciences, Wyeth Pharmaceuticals, UCB, AstraZeneca, Jazz Pharmaceuticals, Abbott Laboratories, and Iroko Pharmaceuticals. Dr. Harris has consulted for Pfizer, Inc. Dr. Harte has received research funding from Aptinyx, Cerephex, Eli Lilly, Forest Laboratories, and Merck; and served as a consultant for Pfizer, Analgesic Solutions, Aptinyx, and deCode Genetics. He is also a member of Arbor Medical Innovations, LLC (Ann Arbor, MI), current licensee of the MAST pain testing device from the University of Michigan. Dr. Harper is supported by NIDCR grant K99 DE026810. Tobias Schmidt-Wilcke was supported by the Deutsche Forschungsgemeinschaft (SFB874 A8).

Download English Version:

<https://daneshyari.com/en/article/8605012>

Download Persian Version:

<https://daneshyari.com/article/8605012>

[Daneshyari.com](https://daneshyari.com)