

Accepted Manuscript

Serum Adiponectin/Ferritin Ratio in Relation to the Risk of Type 2 diabetes and Insulin Sensitivity

Alex Aregbesola, Vanessa D.F. de Mello, Jaana Lindström, Sari Voutilainen, Jyrki K. Virtanen, Sirkka Keinänen-Kiukaanniemi, Tomi-Pekka Tuomainen, Jaakko Tuomilehto, Uusitupa Matti

PII: S0168-8227(17)31608-X
DOI: <https://doi.org/10.1016/j.diabres.2018.05.012>
Reference: DIAB 7373

To appear in: *Diabetes Research and Clinical Practice*

Received Date: 23 October 2017
Revised Date: 21 April 2018
Accepted Date: 8 May 2018

Please cite this article as: A. Aregbesola, V.D.F. de Mello, J. Lindström, S. Voutilainen, J.K. Virtanen, S. Keinänen-Kiukaanniemi, T-P. Tuomainen, J. Tuomilehto, U. Matti, Serum Adiponectin/Ferritin Ratio in Relation to the Risk of Type 2 diabetes and Insulin Sensitivity, *Diabetes Research and Clinical Practice* (2018), doi: <https://doi.org/10.1016/j.diabres.2018.05.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Original contribution

Title

Serum Adiponectin/Ferritin Ratio in Relation to the Risk of Type 2 diabetes and Insulin Sensitivity

Alex Aregbesola ^a, Vanessa D.F. de Mello ^a, Jaana Lindström ^b, Sari Voutilainen ^a, Jyrki K. Virtanen ^a, Sirkka Keinänen-Kiukaanniemi ^{c,d}, Tomi-Pekka Tuomainen ^a, Jaakko Tuomilehto ^{e,f,g}, Uusitupa Matti ^a

^a Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Yliopistonranta 1C, P.O. Box 1627, FI70211 Kuopio, Finland

^b Diabetes Prevention Unit, National Institute for Health and Welfare, Helsinki, Finland

^c Center for Life Course Health Research, University of Oulu, Finland

^d Medical Research Center and Unit of General Practice, Oulu University Hospital and Oulu Health Center, Oulu, Finland

^e Centre for Vascular Prevention, Danube-University Krems, Krems, Austria

^f Dasman Diabetes Institute, Dasman, Kuwait

^g Diabetes Research Group, King Abdulaziz University, Jeddah, Saudi Arabia

Details of the corresponding author

Alex Aregbesola

Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Yliopistonranta 1C, P.O. Box 1627, FI70211 Kuopio, Finland

Email: alex.aregbesola@uef.fi; Tel: +358 (0) 449788099; Fax: +35817162936

Running title: Serum adiponectin/ferritin ratio and type 2 diabetes

Word count: abstract 250/250; text 3352

Download English Version:

<https://daneshyari.com/en/article/8629920>

Download Persian Version:

<https://daneshyari.com/article/8629920>

[Daneshyari.com](https://daneshyari.com)