

Accepted Manuscript

Brief Report

Serum Meteorin-like protein levels decreased in patients newly diagnosed with type 2 diabetes

Ju Hee Lee, Yea Eun Kang, Ji Min Kim, Sorim Choung, Kyong Hye Joung, Hyun Jin Kim, Bon Jeong Ku


PII: S0168-8227(17)31088-4
DOI: <https://doi.org/10.1016/j.diabres.2017.10.005>
Reference: DIAB 7093

To appear in: *Diabetes Research and Clinical Practice*

Received Date: 5 July 2017
Revised Date: 10 August 2017
Accepted Date: 2 October 2017

Please cite this article as: J. Hee Lee, Y. Eun Kang, J. Min Kim, S. Choung, K. Hye Joung, H. Jin Kim, B. Jeong Ku, Serum Meteorin-like protein levels decreased in patients newly diagnosed with type 2 diabetes, *Diabetes Research and Clinical Practice* (2017), doi: <https://doi.org/10.1016/j.diabres.2017.10.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Serum Meteorin-like protein levels decreased in patients newly diagnosed with type 2 diabetes

Ju Hee Lee¹, Yea Eun Kang¹, Ji Min Kim¹, Sorim Choung², Kyong Hye Joung¹, Hyun Jin Kim¹, Bon Jeong Ku¹

Department of Internal Medicine, Chungnam National University School of Medicine, 266 Munwha-ro, Jung-gu, Daejeon, 35015, Republic of Korea¹

Department of Medical Science, Chungnam National University School of Medicine, 266 Munwha-ro, Jung-gu, Daejeon, 35015, Republic of Korea²

Correspondence to: Bon Jeong Ku

Department of Internal Medicine, Chungnam National University School of Medicine, 266 Munwha-ro, Jung-gu, Daejeon, 35015, Republic of Korea.

Tel: +82-42-280-7149

Fax: +82-42-280-8173

E-mail: bonjeong@cnu.ac.kr

Highlights

1. Serum METRNL levels were significantly lower in newly diagnosed type 2 diabetes mellitus patients than in subjects with normal glucose tolerance or prediabetes.
2. Serum METRNL was negatively correlated with fasting plasma glucose, post-load 2-h glucose, HbA1c, and HOMA-IR.
3. This is the first study on serum METRNL levels in human diabetes.

Download English Version:

<https://daneshyari.com/en/article/8630405>

Download Persian Version:

<https://daneshyari.com/article/8630405>

[Daneshyari.com](https://daneshyari.com)