Accepted Manuscript

Incidence of epileptiform discharges in children during induction of anaesthesia using Propofol versus Sevoflurane

Susanne Koch, Leopold Rupp, Christine Prager, Rudolf Mörgeli, Sylvia Kramer, Klaus Dieter Wernecke, Astrid Fahlenkamp, Claudia Spies


PII: S1388-2457(18)31097-6

DOI: https://doi.org/10.1016/j.clinph.2018.05.013

Reference: CLINPH 2008549

To appear in: Clinical Neurophysiology

Received Date: 18 January 2018 Revised Date: 26 May 2018 Accepted Date: 29 May 2018



Please cite this article as: Koch, S., Rupp, L., Prager, C., Mörgeli, R., Kramer, S., Dieter Wernecke, K., Fahlenkamp, A., Spies, C., Incidence of epileptiform discharges in children during induction of anaesthesia using Propofol versus Sevoflurane, *Clinical Neurophysiology* (2018), doi: https://doi.org/10.1016/j.clinph.2018.05.013

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Epileptiform discharges during anaesthesia induction

Incidence of epileptiform discharges in children during induction of anaesthesia using Propofol versus Sevoflurane

Susanne Koch^{1*}, Leopold Rupp^{1*}, Christine Prager², Rudolf Mörgeli¹, Sylvia Kramer¹, Klaus Dieter Wernecke³, Astrid Fahlenkamp¹, Claudia Spies¹

¹Department of Anaesthesiology and Operative Intensive Care Medicine (CCM, CVK), Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Campus Virchow-Klinikum, Augustenburger Platz 1, 13353 Berlin, Germany ²Department of Paediatria and Neurology (CVK), Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Campus Virchow-Klinikum, Augustenburger Platz 1, 13353 Berlin, Germany

Corresponding author:

Dr. Susanne Koch

Department of Anaesthesiology and Intensive Care Medicine CCM / CVK

Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-

Universität zu Berlin, and Berlin Institute of Health

Campus Virchow-Klinikum

Augustenburger Platz 1; 13353 Berlin, Germany

Tel.: +49 30 450 651 078 FAX: +49 30 450551909

E-mail: susanne.koch@charite.de

Abbreviations: DSP = delta with spikes, PSR = rhythmic polyspikes, PED = periodic, epileptiform discharges, SSP = suppression with spikes.

³Institute of Medical Biometry, Charité-Universitätsmedizin Berlin, Sostana GmbH, Berlin, Germany

^{*} Authors contribute equally.

Download English Version:

https://daneshyari.com/en/article/8682217

Download Persian Version:

https://daneshyari.com/article/8682217

<u>Daneshyari.com</u>