

Accepted Manuscript

Title: Preclinical and Clinical Studies for Transplant Tolerance via the Mixed Chimerism Approach

Hajime Sasaki, Tetsu Oura, Thomas R Spitzer, Yi-Bin Chen, Joren C. Madsen, James Allan, David H. Sachs, A.B. Cosimi, Tatsuo Kawai

PII: S0198-8859(17)30549-9

DOI: <https://doi.org/10.1016/j.humimm.2017.11.008>

Reference: HIM 9993

To appear in: *Human Immunology*

Received Date: 13 July 2017

Revised Date: 2 November 2017

Accepted Date: 20 November 2017

Please cite this article as: Sasaki, H., Oura, T., Spitzer, T.R., Chen, Y-B., Madsen, J.C., Allan, J., Sachs, D.H., Cosimi, A.B., Kawai, T., Title: Preclinical and Clinical Studies for Transplant Tolerance via the Mixed Chimerism Approach, *Human Immunology* (2017), doi: <https://doi.org/10.1016/j.humimm.2017.11.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title: Preclinical and Clinical Studies for Transplant Tolerance via the Mixed Chimerism Approach

Hajime Sasaki, Tetsu Oura, Thomas R Spitzer, Yi-Bin Chen, Joren C. Madsen, James Allan, David H. Sachs, A. B. Cosimi and Tatsuo Kawai

Department of surgery, Center for transplant science, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

*Correspondence to:

Tatsuo Kawai, MD, PhD, Massachusetts General Hospital, MGH White 521, 55 Fruit Street, Boston, MA 02114, USA

Tel: +1 617 726 0289

Fax: +1 617 724 8634

E-mail: tkawai@mgh.harvard.edu

Abstract

Based upon observations in murine models, we have developed protocols to induce renal allograft tolerance by combined kidney and bone marrow transplantation (CKBMT) in non-human primates (NHP) and in humans. Induction of persistent mixed chimerism has proved to be extremely difficult in major histocompatibility complex (MHC)-mismatched primates, with detectable

Download English Version:

<https://daneshyari.com/en/article/8737606>

Download Persian Version:

<https://daneshyari.com/article/8737606>

[Daneshyari.com](https://daneshyari.com)