

Accepted Manuscript

Title: Outcome of Infective Endocarditis in the Current Era:
Early Predictors of Poor Prognosis,

Authors: Maria Carmo Pereira Nunes, Milton Henriques
Guimarães-Júnior, Pedro Henrique Oliveira Murta Pinto,
Rodrigo Matos Pinto Coelho, Thais Lins Souza Barros, Nicole
de Paula Aarão Faleiro Maia, Dayane Amaral Madureira,
Rodrigo Citton Padilha Reis, Paulo Henrique Nogueira Costa,
Renato Bráulio, Cláudio Léo Gelape, Teresa Cristina Abreu
Ferrari



PII: S1201-9712(18)30017-1
DOI: <https://doi.org/10.1016/j.ijid.2018.01.016>
Reference: IJID 3146

To appear in: *International Journal of Infectious Diseases*

Received date: 31-10-2017
Revised date: 17-12-2017
Accepted date: 17-1-2018

Please cite this article as: Nunes Maria Carmo Pereira, Guimarães-Júnior Milton Henriques, Murta Pinto Pedro Henrique Oliveira, Coelho Rodrigo Matos Pinto, Souza Barros Thais Lins, Faleiro Maia Nicole de Paula Aarão, Madureira Dayane Amaral, Padilha Reis Rodrigo Citton, Costa Paulo Henrique Nogueira, Bráulio Renato, Gelape Cláudio Léo, Ferrari Teresa Cristina Abreu. Outcome of Infective Endocarditis in the Current Era: Early Predictors of Poor Prognosis, *International Journal of Infectious Diseases* <https://doi.org/10.1016/j.ijid.2018.01.016>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Outcome of Infective Endocarditis in the Current Era: Early Predictors of Poor Prognosis

Maria Carmo Pereira Nunes^{a,b}; Milton Henriques Guimarães-Júnior^b; Pedro Henrique Oliveira Murta Pinto^b; Rodrigo Matos Pinto Coelho^a; Thais Lins Souza Barros^b; Nicole de Paula Aarão Faleiro Maia^b; Dayane Amaral Madureira^b; Rodrigo Cíton Padilha Reis^c; Paulo Henrique Nogueira Costa^b; Renato Bráulio^b, Cláudio Léo Gelape^b; Teresa Cristina Abreu Ferrari^{a,b}

^aDepartment of Internal Medicine, School of Medicine, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^b Department of Cardiology and Cardiovascular Surgery of the Hospital das Clínicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^cDepartment of Statistics, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

Address for correspondence: Maria do Carmo P Nunes
Departamento de Clínica Médica - UFMG
Av. Professor Alfredo Balena, 190, Santa Efigênia
30130 100 – Belo Horizonte, MG
Phone: +55 31 34099746, Fax: +55 31 34099437
Email: mcarmo@waymail.com.br

Abstract

Background: Early identification of patients at risk of IE complications using parameters obtained routinely is essential for clinical decision making guidance. This study aimed to identify a parameter at hospital admission that predicts outcome, adding value to other well-known factors of poor prognosis in IE.

Download English Version:

<https://daneshyari.com/en/article/8739057>

Download Persian Version:

<https://daneshyari.com/article/8739057>

[Daneshyari.com](https://daneshyari.com)