Accepted Manuscript

Kaposi Sarcoma Incidence in Females is nearly Four-Fold Higher in the Lower Rio Grande Valley compared to the Texas Average

Wendy Innis-Whitehouse, Xiaohui Wang, Nicolas Restrepo, Carlos Salas, Katia Moreno, Alvaro Restrepo, Megan Keniry

PII: S2468-2942(17)30189-2 DOI: 10.1016/j.ctarc.2018.06.003

Reference: CTARC 90

To appear in: Cancer Treatment and Research Communications

9 June 2018 Revised date: Accepted date:



Cancer Treatment and Research Communications (2018), doi: 10.1016/j.ctarc.2018.06.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Highlights

- The incidence of Kaposi sarcoma is nearly four-fold higher in females in Hidalgo County Texas compared to the average for females in the State of Texas
- An increased incidence of Kaposi sarcoma is found on the United States-Mexican border in the State of Texas
- The increased incidence of Kaposi sarcoma in South Texas females has significant health implications: environmental factors, genetic contributions and possible links to diabetes need further investigation



Download English Version:

https://daneshyari.com/en/article/8785738

Download Persian Version:

https://daneshyari.com/article/8785738

<u>Daneshyari.com</u>