


## Accepted Manuscript

The association of duration and severity of disease with executive function: differences between drug-naïve patients with bipolar and unipolar depression

Tao Yang , Guo-Qing Zhao , Rui-Zhi Mao , Jia Huang ,  
Xian-Rong Xu , You-Song Su , Na Zhu , Ru-Bai Zhou , Xiao Lin ,  
Wei-Ping Xia , Fan Wang , Rui Liu , Xing Wang , Zhi-Jia Huang ,  
Yong Wang , Ying-Yan Hu , Lan Cao , Cheng-Mei Yuan ,  
Zuo-Wei Wang , Raymond W. Lam , Jun Chen , Yi-Ru Fang



PII: S0165-0327(18)30107-1  
DOI: [10.1016/j.jad.2018.05.051](https://doi.org/10.1016/j.jad.2018.05.051)  
Reference: JAD 9827

To appear in: *Journal of Affective Disorders*

Received date: 17 January 2018  
Revised date: 20 March 2018  
Accepted date: 28 May 2018

Please cite this article as: Tao Yang , Guo-Qing Zhao , Rui-Zhi Mao , Jia Huang , Xian-Rong Xu , You-Song Su , Na Zhu , Ru-Bai Zhou , Xiao Lin , Wei-Ping Xia , Fan Wang , Rui Liu , Xing Wang , Zhi-Jia Huang , Yong Wang , Ying-Yan Hu , Lan Cao , Cheng-Mei Yuan , Zuo-Wei Wang , Raymond W. Lam , Jun Chen , Yi-Ru Fang , The association of duration and severity of disease with executive function: differences between drug-naïve patients with bipolar and unipolar depression , *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.05.051](https://doi.org/10.1016/j.jad.2018.05.051)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- The executive function in drug-naïve UPD patients was worse than that in BDD patients.
- The executive function of drug-naïve UDP patients was lower than that of HC significantly.
- The executive function of drug-naïve BDD patients was likely to influence by the duration of current depressive episode. However, the number of mood episode was the main influence factor for executive function in UPD patients.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8815272>

Download Persian Version:

<https://daneshyari.com/article/8815272>

[Daneshyari.com](https://daneshyari.com)