

www.elsevier.com/locate/jad

Co-morbid Depressive Disorder is Associated with Better Neurocognitive Performance in First Episode Schizophrenia Spectrum

Sarah E. Herniman, Sue M. Cotton, Eoin Killackey, Robert Hester, Kelly A. Allott

PII: S0165-0327(17)31019-4
DOI: https://doi.org/10.1016/j.jad.2017.12.088
Reference: JAD9496

To appear in: *Journal of Affective Disorders*

Received date: 22 May 2017
Revised date: 11 November 2017
Accepted date: 31 December 2017

Cite this article as: Sarah E. Herniman, Sue M. Cotton, Eoin Killackey, Robert Hester and Kelly A. Allott, Co-morbid Depressive Disorder is Associated with Better Neurocognitive Performance in First Episode Schizophrenia Spectrum, *Journal of Affective Disorders,* https://doi.org/10.1016/j.jad.2017.12.088

Co-morbid Depressive Disorder is Associated with Better Neurocognitive Performance in First Episode Schizophrenia Spectrum

Sarah E. Herniman[123]
sarah.herniman@unimelb.edu.au

Sue M. Cotton[12]
sue.cotton@orygen.org.au

Eoin Killackey[12]
eoin.killackey@orygen.org.au

Robert Hester[2]
hesterr@unimelb.edu.au

Kelly A. Allott[12]
kelly.allott@orygen.org.au

[1] Orygen, the National Centre of Excellence in Youth Mental Health, Melbourne, Australia Postal address: Locked Bag 10, Parkville VIC 3052 Australia

[2] Centre for Youth Mental Health, University of Melbourne, Melbourne, Australia Postal address: Locked Bag 10, Parkville VIC 3052 Australia

[3] Melbourne School of Psychological Sciences, University of Melbourne, Melbourne, Australia Postal address: Level 2, Redmond Barry Building 115, University of Melbourne, Parkville VIC 3010

Corresponding author:
Sarah E. Herniman, Orygen, The National Centre of Excellence in Youth Mental Health, University of Melbourne, 35 Poplar Road, Parkville VIC 3052, Australia
Email: sarah.herniman@unimelb.edu.au
Mob: +61412778046

Abstract

Background: Both major depressive disorder (MDD) and first episode schizophrenia spectrum (FES) are associated with significant neurocognitive deficits. However, it remains unclear whether the neurocognitive deficits in individuals with FES are more severe if there