

Accepted Manuscript

Title: Bipolar Fracture Dislocation of Clavicle: A Report of Osteosynthesis and Early Soft Tissue Reconstruction

Authors: Renaldi Prasetia, Hermawan Nagar Rasyid

PII: S2210-2612(17)30533-3
DOI: <https://doi.org/10.1016/j.ijscr.2017.10.025>
Reference: IJSCR 2828



To appear in:

Received date: 19-7-2017
Revised date: 13-10-2017
Accepted date: 18-10-2017

Please cite this article as: Prasetia Renaldi, Rasyid Hermawan Nagar. Bipolar Fracture Dislocation of Clavicle: A Report of Osteosynthesis and Early Soft Tissue Reconstruction. *International Journal of Surgery Case Reports* <https://doi.org/10.1016/j.ijscr.2017.10.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Bipolar Fracture Dislocation of Clavicle: A Report of Osteosynthesis and Early Soft Tissue Reconstruction

Renaldi Prasetia, MD., MMedSc

Hermawan Nagar Rasyid, MD., MT., PhD

Department of Orthopedics and Traumatology Universitas Padjadjaran, Dr. Hasan Sadikin Teaching Hospital, Bandung, Indonesia

Corresponding Author:

Renaldi Prasetia, MD., MMedSc

Department of Orthopedics and Traumatology

Faculty of Medicine, Universitas Padjadjaran

Dr. Hasan Sadikin Teaching Hospital

Jl. Pasteur no. 38, Bandung, West Java, Indonesia, 40161

Email: renaldi.prasetia@gmail.com

Highlights

- Bipolar fracture-dislocation of the clavicle in concomitant with coracoid fracture are uncommon traumatic injuries.
- The scarce number of published cases have made the treatment of these injuries still controversial.
- We proposed osteosynthesis and early soft tissue joint reconstruction as an option treatment to be considered.
- Anatomic reconstruction is the key to achieve success and good functional outcome in bipolar fracture- dislocation as seen in various ligament reconstruction and joint restoration surgeries.
- Biological Healing in soft tissue reconstruction can prevent acromioclavicular and sternoclavicular joint re-dislocation as bipolar clavicle fracture-dislocation complication

Download English Version:

<https://daneshyari.com/en/article/8833040>

Download Persian Version:

<https://daneshyari.com/article/8833040>

[Daneshyari.com](https://daneshyari.com)