



# Natureza & Conservação

Brazilian Journal of Nature Conservation

Supported by Boticário Group Foundation for Nature Protection

<http://www.naturezaeconservacao.com.br>



## Research Letters

# Evaluation of Global Positioning System telemetry collar performance in the tropical Andes of southern Ecuador

Meghan J. Camp<sup>a,b,\*</sup>, Janet L. Rachlow<sup>a</sup>, Rodrigo Cisneros<sup>c</sup>, David Roon<sup>d</sup>, Reid J. Camp<sup>e</sup>

<sup>a</sup> Department of Fish and Wildlife Sciences, University of Idaho, Moscow, United States

<sup>b</sup> School of the Environment, Washington State University, Pullman, United States

<sup>c</sup> Department of Biology, Universidad Técnica Particular de Loja, Loja, Ecuador

<sup>d</sup> Department of Ecology and Conservation Biology, University of Idaho, Moscow, United States

<sup>e</sup> Eco Logical Research, Inc., Logan, United States

### ARTICLE INFO

#### Article history:

Received 6 January 2015

Accepted 28 July 2016

Available online xxx

#### Keywords:

Advanced Telemetry Systems

Data screening

Fix rate

Location error

Páramo

### ABSTRACT

The objective of this study was to evaluate if canopy cover, topographic obstruction of the sky, and differences among vegetation type affected performance of global positioning system (GPS) telemetry in southern Ecuador. A GPS collar was placed at 30 test sites in Podocarpus National Park, Ecuador, and we estimated canopy cover and topographic obstruction at each site. The mean fix success rate was 87.7% (SD = 22.3%), and the mean location error for test sites was 9.7 m (SD = 4.17 m). Canopy cover significantly reduced the performance of GPS telemetry in our study area both in terms of location error and fix acquisition rate. However, topographic obstruction did not significantly influence location errors or fix acquisition rate. Screening data to remove less accurate, two-dimensional fixes resulted in an 11% data loss and reduced the mean location error by 3.8 m. Understanding how habitat variables influence fix acquisition and location errors of GPS collars will assist researchers in evaluating potential biases and developing methods to correct for the effect of such biases on analyses of habitat use and animal movements in the tropical Andes in southern Ecuador.

© 2016 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Global positioning system (GPS) telemetry enables remote collection of large quantities of animal locations at all times over broad geographic areas. Despite the advantages of GPS technology, habitat characteristics can influence performance of

GPS telemetry in ways that can potentially bias analyses that rely on GPS locations (Dussault et al., 1999; Frair et al., 2010; Lewis et al., 2007). Therefore, the capability of GPS telemetry devices should be evaluated in the field before they are used in studies of animal ecology, and appropriate corrective measures (e.g., correction or weighing factors) taken prior to analysis (Dussault et al., 1999; Webb et al., 2013).

\* Corresponding author.

E-mail address: [meghan.camp@wsu.edu](mailto:meghan.camp@wsu.edu) (M.J. Camp).

<http://dx.doi.org/10.1016/j.ncon.2016.07.002>

1679-0073/© 2016 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

There are two types of errors associated with GPS telemetry. One type is unsuccessful fix acquisition, which results in missing location data (D'Eon et al., 2002; Frair et al., 2004). A second type of error is location inaccuracy that occurs when the GPS unit records a location that is inaccurate (Frair et al., 2004). Habitat features that influence the ability of a GPS unit to receive satellite signals can result in both types of location errors; these include percent of vegetation canopy cover, vegetation cover type, and rugged terrain (D'Eon et al., 2002; Frair et al., 2004; Rempel et al., 1995). Other factors that can influence the errors of GPS collars include weather conditions, time of day, season, animal behavior, as well as the time interval between attempted fixes, device orientation, time in the field, and device manufacturer and model (Dussault et al., 2001; Frair et al., 2004; Matthews et al., 2013; Sager-Fradkin et al., 2007). Furthermore, GPS units are capable of recording either three-dimensional (3-D) fixes or two-dimensional fixes (2-D). A 2-D fix is recorded when signals are received from only three satellites, and a 3-D fix is achieved when signals are received from  $\geq 4$  satellites; 3-D fixes are typically more accurate than 2-D fixes (Di Orio et al., 2003).

Our objective was to evaluate how canopy cover and topographic obstruction affected the total fix rate, 3-D fix rate, and location error in the tropical Andes of southern Ecuador. We expected that both location accuracy and fix success rate would be negatively influenced by increasing vegetation canopy cover and topographic obstruction of the sky. Such evaluations have been conducted frequently in North America and have been reported for tropical lowlands; however, to our knowledge, performance of GPS telemetry collars has not been systematically evaluated and reported for lower latitudes and rugged terrain in South America.

## Methods

### Study site

The study area is located in Podocarpus National Park (PNP) in southern Ecuador ( $4^{\circ}17'0''$  S,  $79^{\circ}0'0''$  W). Elevation in the Park varies between 800 and 3695 m, and the area receives up to 6000 mm of rain per year. The grass páramo occurs between 3335 and 3385 m, and the vegetation is dominated by forbs, *Chusquea* bamboo and other grasses, and woody plants less than 10 cm tall (Keating, 1999). The shrub páramo occurs below 3335 m and nearly 100% of the area is covered by shrubs and bromeliads. The vegetation below 3150 m is the cloud forest and is dominated by trees including three species of the genus *Podocarpus*.

### GPS testing

We stratified the study area by vegetation type and then selected representative locations at random within each stratum, with the constraint that locations were within 1 km of a road or trail. We tested 30 different locations. We randomized the order of locations during the deployment period to avoid a potential bias associated with variation in GPS performance due to weather, which changed frequently, even within days. We classified the vegetation types as grass, which included the

grass páramo; shrub, which included the shrub páramo type; and tree, which included pine forest, cloud forest and rain forest. The minimum distance between test sites was 650 m, and the maximum distance was 31 km. The entire study region covered approximately 47,900 ha.

We used an Iridium/GPS Location Collar (Model G2110E; Advanced Telemetry Systems, Isanti, MN, USA) weighing 825 g programmed to record a location every 30 min to evaluate the influence of habitat on GPS performance. We placed the collar on a stand approximately 1 m above the ground in a vertical position with the antenna facing upward at each of the 30 test sites for a 24-h period during 29 May–12 July 2012. The GPS collar attempted to record at least 48 fixes per test site, and our sampling unit was the 30 independent test sites.

At each test site, we recorded the percent canopy cover above the collar using a spherical densiometer, and we calculated the available sky (i.e., the inverse of terrain obstruction) from each test site using the viewshed function in 3-D spatial analyst in ArcGIS 10 (ESRI, Redlands, CA). To do this, we created a layer of points in the sky 100 m above the highest point in the study area, and from each test location, we calculated the percentage of points that were visible from the test site to provide a value of available sky (D'Eon et al., 2002).

### Data analyses

We recorded the actual location of collar test sites with a handheld GPS unit (Garmin, GPSMAP 60CSx). For each location we allowed the handheld GPS unit to average multiple fixes to improve location accuracy. We recorded a location after the estimated accuracy was  $<6$  m. We calculated location errors of the GPS collar by determining the distance between the actual location and the estimated location recorded by the collar using Spatial Analyst in ArcGIS 10 (ESRI, Redlands, CA). The fix rate was quantified as the proportion of attempted fixes that were successful. We compared location error and fix rate among habitat types using a one-way analysis of variance (ANOVA). We tested if percent canopy cover and available sky were correlated using a Pearson product-moment correlation. We evaluated multiple linear regression models of location error and fix rate that included the continuous variables of percent canopy cover and available sky (PROC REG; SAS Institute, Inc., Cary, NC, USA). We performed an arcsin transformation on the response variables for overall fix rate and 3-D fix rate because they were proportional data (Zar, 1999). The test location was the experimental unit ( $n = 30$ ); however, individual fixes were considered independent samples to compare 2-D and 3-D location errors using Student's two-sample *t*-test.

## Results

Mean percent canopy cover was 0% in the grass vegetation type 76% in the shrub type (range = 0–93%, SE = 14%), and 88% in the tree type (range = 34–96%, SE = 10%). Missed fixes tended to occur in areas with high canopy cover. The mean fix rate for the GPS collar at the 30 test sites, across all three vegetation types, was 87.7% (SD = 22.3%, range = 6–100%). The mean fix rate was 100% in the grass vegetation type, 70% in shrub

Download English Version:

<https://daneshyari.com/en/article/8849457>

Download Persian Version:

<https://daneshyari.com/article/8849457>

[Daneshyari.com](https://daneshyari.com)