



ELSEVIER

Contents lists available at ScienceDirect

Environment International

journal homepage: www.elsevier.com/locate/envint

Prenatal exposure to polychlorinated biphenyls and fetal growth in British girls

Jill F. Patel^{a,b}, Terry J. Hartman^{a,b,c,*}, Andreas Sjodin^a, Kate Northstone^{d,e}, Ethel V. Taylor^a

^a Centers for Disease Control and Prevention, Atlanta, GA 30341, United States

^b Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA 30322, United States

^c Winship Cancer Institute, Emory University, Atlanta, GA 30322, United States

^d The National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West (NIHR CLAHRC West) at University Hospitals Bristol NHS Foundation Trust, UK

^e School of Social and Community Medicine, University of Bristol, Bristol, UK

ARTICLE INFO

Handling Editor: Olga-Ioanna Kalantzi

Keywords:

Polychlorinated biphenyls
Endocrine disrupting chemicals
Pregnancy
Birthweight

ABSTRACT

Polychlorinated biphenyls (PCBs) are synthetic chemicals that bioaccumulate in the food chain. PCBs were used primarily for industrial applications due to their insulating and fire retardant properties, but were banned in the 1970s in the United States and in the 1980s in the United Kingdom, as adverse health effects following exposure were identified. Previous studies of populations with high PCB exposure have reported inverse associations with birth weight and gestational length. Birth weight is a powerful predictor of infant survival, and low birth weight can predispose infants to chronic conditions in adult life such as diabetes and cardiovascular diseases.

Using data from the Avon Longitudinal Study of Parents and Children, we investigated the association between prenatal exposure to PCBs and fetal growth in a sample of 448 mother-daughter dyads. Concentrations of three common PCB analytes, PCB-118, PCB-153 and PCB-187, were measured in maternal serum collected during pregnancy, and fetal growth was measured by birth weight and birth length. Multivariable linear regression was used to examine the associations between PCB analytes and measures of fetal growth, after adjusting for parity, maternal age, pre-pregnancy BMI, educational status, tobacco use and gestational age of infant at sample collection. Birth length, ponderal index and gestational age were not associated with any of the PCB analytes. Mothers' educational status modified associations for PCB analytes with birthweight. We observed significant inverse associations with birth weight only among daughters of mothers with less education. Daughter's birth weight was -138.4 g lower (95% CI: -218.0 , -58.9) for each 10 ng/g lipid increase in maternal serum PCB-118. Similarly, every 10 ng/g lipid increase in maternal serum PCB-153 was associated with a -41.9 g (95% CI: -71.6 , -12.2) lower birth weight. Every 10 ng/g lipids increase in maternal serum PCB-187, was associated with a -170.4 g (95% CI: -306.1 , -34.7) lower birth weight, among girls with mothers in the lowest education group.

Our findings suggest that prenatal exposure to PCBs is inversely associated with daughters' birth weight and that mothers' education, which is a possible marker for socioeconomic status, significantly modified the association between maternal PCB concentrations and birth weight in female newborns.

1. Introduction

Polychlorinated biphenyls (PCBs) are a family of synthetic organic chemicals, comprising 209 chemically related compounds that were used between 1930 and 1977 for various industrial applications because of their insulating and fire-retardant properties (Ross, 2004). PCBs were banned in the 1970s in the United States and in the 1980s in the United Kingdom, as adverse health effects following exposure were identified. PCBs are biphenyls with between one and ten chlorine atoms

attached, and the degree of chlorination determines the stability and lipophilicity of the specific PCB analyte (Kezios et al., 2012). PCBs with lower degrees of chlorination tend to be more rapidly excreted from the body, while more chlorinated PCB compounds (e.g., PCB 153) are retained for a longer period of time, many with biological half-lives in the order of years (Taylor et al., 1989; El Majidi et al., 2012). There are no known natural sources of PCBs (Public Health, 2000) and once PCBs are released into the environment, they do not readily break down and can easily cycle between air, water and soil. Furthermore, they can be

* Corresponding author at: Rollins School of Public Health, Winship Cancer Institute, Emory University, 1518 Clifton RD NE, CNR 3035, Atlanta, GA 30322, United States.
E-mail address: tjhartm@emory.edu (T.J. Hartman).

carried long distances and have been found in areas of snow and sea water far from the original release site (Taylor et al., 1989).

The most common sources of environmental exposure to these substances are dairy products, meat and especially fish (Koopman-Esseboom et al., 1994). PCBs are stored mainly in human adipose tissue, and their poor metabolism results in elimination half-lives of approximately 10–15 years (Patandin et al., 1998; Ritter et al., 2011). Animal and human studies have shown that PCBs cross the placenta (Kezios et al., 2012) and the quantities of PCBs found in cord serum may be considerable relative to the size of the developing fetus (Jacobson et al., 1984). Endocrine pathways that are important for fetal development, such as thyroid hormone signaling, can be disrupted by PCBs, potentially leading to decreased in utero growth (Public Health, 2000).

Birth weight is one of the most, if not the most, powerful predictors of infant survival (WHO, 1992), with low birth weight contributing to about 9.1 million infant deaths each year. Globally, 17% of total births are to low birth weight newborns (WHO, 1992). Low birth weight in early childhood can be associated with adiposity in adolescents and earlier pubertal maturation, and is a strong predictor of the development of obesity, hypertension and cardiovascular disease in adults (Wohlfahrt-Veje et al., 2014). Additionally, low birth weight is a predictor for other adverse outcomes such as poor school performance, high blood pressure and cardiovascular diseases (Law et al., 2002; Barker, 2002; Barker, 1998; Thompson et al., 2001). Therefore, it is important to determine whether prenatal PCB exposure is associated with birth outcomes such as low birth weight. The objective of this study was to investigate the association between prenatal exposure to PCBs and fetal growth in a well-characterized British sample of mother-daughter dyads.

2. Methods

2.1. Population

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a prospective birth-cohort study designed to evaluate the influences of genetic and environmental factors on health. During the years 1990–1992, 14,541 pregnant women residing in Avon, Great Britain with an estimated delivery date between April 1991 and December 1992 were enrolled in the study. These (initial) pregnancies resulted in a total of 14,062 live births and 13,988 children alive at one year of age. An additional 713 eligible children were enrolled at approximately 7 years of age and their data are available for analyses when including variables collected from the age of seven and later. Details of recruitment methods are described in detail elsewhere (Fraser et al., 2013; Boyd et al., 2013).

This study examined associations between maternal exposure to PCBs and fetal growth in girls in an ancillary study designed to look at the association between maternal serum concentrations of environmental exposures and daughter's puberty characteristics (Christensen et al., 2011). To be considered, girls had to have at least two pubertal assessments to allow for classification of age at menarche. The ancillary study included all girls with early menarche (< 11.5 years; $n = 218$) and a random sample of girls without early menarche ≥ 11.5 years ($n = 230$). Informed consent was provided at the time of enrollment by the mothers. Human subjects' protection and ethical approval were provided by the ALSPAC Law and Ethics Committee, the Local Research Ethics Committees, and the Centers for Disease Control and Prevention (CDC) Institutional Review Board. Please note that the study website contains details of all the data that are available through a fully searchable data dictionary, <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary>.

2.2. Data collection

Outcomes of interest included birth weight (in grams), birth length (in centimeters), and gestational age (in weeks) which were abstracted

from medical records. Low birth weight was considered < 2500 g. Ponderal index was calculated using the following formula: (weight in g/height in cm³) \times 100 (Maisonet et al., 2012). Self-reported data on maternal prenatal characteristics and behaviors were obtained from questionnaires completed during pregnancy. Data collection and methods have been described in detail elsewhere (Golding et al., 2001).

2.3. Laboratory analyses

Blood samples were collected during pregnancy, processed and serum was stored frozen at -20°C . In 2008, samples were shipped to the National Center for Environmental Health's Division of Laboratory Sciences, Centers for Disease Control and Prevention where PCB-118, PCB-153 and PCB-187 were measured by gas chromatography isotope dilution high resolution mass spectrometry (GC-IDHRMS) (Hovander et al., 2000). This analysis presents lipid-adjusted exposures and measurements recorded as 0 ng/g are noted as < LOD.

2.4. Statistical analyses

Pearson correlation coefficients were used to assess the relationship between PCB analytes. Stratum-weighted linear models, which accounted for the nested case-control study design, were used to estimate the association between individual PCB concentrations (PCB-118, PCB-153, and PCB-187) and fetal growth markers (birth weight, birth length, gestational age, and ponderal index). To adjust for the original selection criteria for the nested case-control study, cases (all girls who attained menarche < 11.5 years) and controls (girls who attained menarche ≥ 11.5 years) were assigned weights of 1 and 15.1, respectively. Each PCB analyte was examined individually in a stratum-weighted model for each birth outcome, and backwards elimination was used to identify potential covariates which appreciably contributed to model fit or interpretation. Covariates considered included previous births (0/ ≥ 1), maternal age (continuous), maternal race (white/non-white), pre-pregnancy BMI (continuous), educational status (categorical), tobacco use (binary), and gestational age when maternal serum sample was obtained (continuous). In this analysis, not attaining any General Certificates of Secondary Education (GCSEs, at 16 years of age) was coded as "< O" (low) educational level, obtaining GCSEs as "O" (medium) and completing GCSEs and/or vocational training with additional education (e.g., University or Advanced) was considered "> O level" (high). After backwards elimination, the remaining covariates for birth weight outcome included previous births, maternal BMI, race, education, tobacco use during pregnancy and gestational ages at sample collection. For birth length, the confounders included in the model were parity and maternal BMI. The ponderal index model included the confounders, parity, maternal BMI, and tobacco use during pregnancy. For gestational age at birth, the only confounder remaining in the model was gestational age when serum sample was collected. Effect modification by maternal smoking and maternal education was also examined by testing appropriate interaction terms for statistical significance. SAS version 9.3 (SAS Institute Inc., Cary, NC) was used to conduct all analyses. All statistical tests were 2-tailed; a p -value of < 0.05 was considered statistically significant.

3. Results

More than half of the mothers were 29 years of age or younger at delivery and reported having normal pre-pregnancy BMI. About half of the mothers had an educational level of less than or equal to O level and most mothers were white (94%) (Table 1). The median gestational age at serum sample collection was 15 weeks with an interquartile range of 10–28 weeks. Among 448 daughters included in the study population, mean birth weight (\pm SD) was 3396.7 (\pm 498.7) g; birth length was 50.4 (\pm 2.2) cm; ponderal index was 2.7 (\pm 0.2) kg/m³; and gestational age was 39.8 (\pm 1.6) weeks (data not presented). Fewer than 4% of daughters were born preterm or weighed < 2500 g.

Download English Version:

<https://daneshyari.com/en/article/8855204>

Download Persian Version:

<https://daneshyari.com/article/8855204>

[Daneshyari.com](https://daneshyari.com)