



Contents lists available at ScienceDirect

## Environmental Pollution

journal homepage: [www.elsevier.com/locate/envpol](http://www.elsevier.com/locate/envpol)

## Commentary

Emission of volatile organic compounds from plants shows a biphasic pattern within an hormetic context<sup>☆</sup>Evgenios Agathokleous<sup>a, b, \*</sup>, Mitsutoshi Kitao<sup>a</sup>, Edward J. Calabrese<sup>c</sup><sup>a</sup> Hokkaido Research Center, Forestry and Forest Products Research Institute (FFPRI), Forest Research and Management Organization, 7 Hitsujigaoka, Sapporo, Hokkaido, 062-8516, Japan<sup>b</sup> Research Faculty of Agriculture, Hokkaido University, Kita 9 Nishi 9, Sapporo, Hokkaido, 060-8589, Japan<sup>c</sup> Department of Environmental Health Sciences, Morrill I, N344, University of Massachusetts, Amherst, MA 01003, USA

## ARTICLE INFO

## Article history:

Received 24 February 2018

Received in revised form

4 April 2018

Accepted 6 April 2018

## Keywords:

Dose-response

Hormesis

Risk assessment

U-shape curve

Volatile organic compounds

## ABSTRACT

Biogenic volatile organic compounds (BVOCs) are released to the atmosphere from vegetation. BVOCs aid in maintaining ecosystem sustainability via a series of functions, however, VOCs can alter tropospheric photochemistry and negatively affect biological organisms at high concentrations. Due to their critical role in ecosystem and environmental sustainability, BVOCs receive particular attention by global change biologists. To understand how plant VOC emissions affect stress responses within a dose-response context, dose responses should be evaluated. This commentary collectively documents hormetic-like responses of plant-emitted VOCs to external stimuli. Hormesis is a generalizable biphasic dose response phenomenon where the response to low doses acts in an opposite way at high doses. These collective findings suggest that ecological implications of low-level stress that may alter BVOC emissions should be considered in future studies. This commentary promotes new insights into the interface between biological systems and environmental change that influence several parts of the globe, and provide a base for advancing hazard assessment testing strategies and protocols to provide decision makers with adequate data for generating environmental standards.

© 2018 Elsevier Ltd. All rights reserved.

## 1. Introduction

Volatile organic compounds (VOCs) are abundant in the Earth's atmosphere as a result of natural (e.g. vegetation) and anthropogenic emissions. Recently, VOCs received particular attention due to their contribution in the formation of ground-level ozone (O<sub>3</sub>). Such O<sub>3</sub> levels can be significantly elevated in the Northern Hemisphere, posing risks to animals and vegetation (Agathokleous et al., 2016; Kampa and Castanas, 2014; Li et al., 2017).

Despite their potential to affect the formation of O<sub>3</sub>, VOCs help to maintain ecosystem sustainability (Peñuelas and Staudt, 2010; Tumlinson, 2014). For example, VOCs are part of an external communications network which transfers plant-to-plant signals

(Peñuelas and Staudt, 2010; Tumlinson, 2014). A further crucial role is mediating interactions between plants and (a)biotic factors (Peñuelas and Staudt, 2010; Tumlinson, 2014). For instance, when a plant is attacked by herbivores or other organisms such as fungi, it releases a complex VOC blend which will be transferred to neighboring plants preconditioning against potential threats within a defined time window (Himanen et al., 2010; Piesik et al., 2007). Plant-emitted VOCs can be of profound significance to insects which rely on them for tracing their route to host plants (Piesik et al., 2014); plants may also rely on VOCs to attract insects as is the case of the Venus flytrap (*Dionaea muscipula*) which relies on food smell mimicry to attract prey. In addition to VOC utilization by animals to trace host/prey, animals utilize VOCs for sexual communication, i.e. pheromones (Tumlinson, 2014). VOCs, therefore, are relevant to a variety of scientific disciplines, including agriculture, behavioral science, biology, chemistry, earth and planetary sciences, ecology, engineering, environmental science, and physiology.

Within the context of global change, it is necessary to understand how plant VOC emissions affect stress responses in a dose-response framework. Three common dose-response models are

<sup>☆</sup> This paper has been recommended for acceptance by B. Nowack.

\* Corresponding author. Hokkaido Research Center, Forestry and Forest Products Research Institute (FFPRI), Forest Research and Management Organization, 7 Hitsujigaoka, Sapporo, Hokkaido, 062-8516, Japan.

E-mail addresses: [globalscience@frontier.hokudai.ac.jp](mailto:globalscience@frontier.hokudai.ac.jp), [evgenios@ffpri.affrc.go.jp](mailto:evgenios@ffpri.affrc.go.jp), [evgenios@for.agr.hokudai.ac.jp](mailto:evgenios@for.agr.hokudai.ac.jp) (E. Agathokleous), [kitao@ffpri.affrc.go.jp](mailto:kitao@ffpri.affrc.go.jp) (M. Kitao), [edwardc@schoolph.umass.edu](mailto:edwardc@schoolph.umass.edu) (E.J. Calabrese).

the threshold, linear-non-threshold (LNT) and hormetic (Fig. 1). The threshold and LNT models have dominated the literature throughout the 20th century (Calabrese, 2017a; b). Hormesis appeared in the mainstream only after the 2000s, creating a dose-response revolution via extensive documentation and challenging the belief that the fundamental nature of dose-response is biphasic rather than a type of combined set of threshold and linear frameworks (Agathokleous, 2018; Calabrese, 2011; Calabrese and Blain, 2011; Calabrese and Mattson, 2017).

Assessment of more than ten thousand dose responses revealed that hormesis is a highly generalizable biological phenomenon, with hormetic dose responses being independent of biological model, endpoint, and stressor, and having maximum response in the low dose zone less than 2-fold the control and width of the low-dose, potentially beneficial, zone lower than 10-fold in dose range (Agathokleous, 2018; Agathokleous et al., 2018; Calabrese, 2011, 2013; 2014, 2017c; Calabrese and Blain, 2009, 2011; Calabrese and Mattson, 2011, 2017; Cedergreen et al., 2007; Hashmi et al., 2014; Kim et al., 2018; Poschenrieder et al., 2013; Vargas-Hernandez et al., 2017).

## 2. Analysis

An earlier hypothesis that VOCs emitted by plants may display a hormetic response to environmental factors (Calfapietra et al., 2009) along with recently-reported biphasic emission kinetics of the lipoxygenase (LOX) pathway in cucumber (*Cucumis sativus* L.) in response to time-dependent exposure to methyl jasmonate (MeJA) stimulated the need to assess whether VOCs emission by plants is biphasic by time and by dose in response to stress (Jiang et al., 2017). Emission kinetics of benzenoid methyl salicylate, 4,8-dimethyl-nona-1,3,7-triene (DMNT), sesquiterpene (*E,E*- $\alpha$ -farnesene, LOX, or monoterpene (*E*)- $\beta$ -ocimene tended to be biphasic in response to the time *Alnus glutinosa* plants were exposed to: feeding from moth *Cabera pusaria* larvae; feeding from sawfly *Monsoma pulveratum* larvae (a); drought (b); or combination of (a) and (b) (Copolovici et al., 2011, 2014). Data suggesting a biphasic emission of a variety of green leaf VOCs were also found in other plants in response to the time exposed to herbivory or mechanical wounding (Harren and Cristescu, 2013; Piesik et al., 2013), drought (Jubany-Marí et al., 2010), or light (Rasulov et al., 2009, 2011).

In addition to the time-dependent biphasic VOCs responses, concentration-dependent biphasic VOCs responses to external stimuli were also reported (Carriero et al., 2016; Rasulov et al., 2009, 2011). For instance, the response of isoprene emission rate to CO<sub>2</sub> levels was biphasic in aspen (*Populus tremula*  $\times$  *P. tremuloides*) leaves (Fig. 2A); isoprene emission rate was closely related to the chloroplastic DMADP pool (Rasulov et al.,

2009). In the same aspen hybrid, signs for a biphasic dose response of isoprene emission rate to CO<sub>2</sub> (Fig. 2B), O<sub>2</sub> (Fig. 2C), and leaf temperature (Fig. 2D) were observed (Rasulov et al., 2011). Biphasic emission of isoprenoids is supported by further studies, particularly in response to O<sub>3</sub> (Calfapietra et al., 2009; Tani et al., 2017; Yuan et al., 2016). Emission rates of several compounds ( $\alpha$ -pinene,  $\beta$ -pinene, limonene, ocimene, hexanal and DMNT) from birch (*Betula pendula* Roth) plants exposed to O<sub>3</sub> and 10, 30 or 70 kg N ha<sup>-1</sup> yr<sup>-1</sup> consistently indicate the occurrence of hormesis (Carriero et al., 2016), however, unlike isoprene, with reduced emission in the low-dose region (Fig. 2E,F,G,H). Interestingly, the latter results (Carriero et al., 2016) raised the question that some biphasic dose responses may have two maxima, in agreement with results from other studies (Jiang et al., 2017). Reduced emission potential of sesquiterpenes in the low-dose region has been also reported for Norway spruce (*Picea abies* (L.) H. Karst.) which displayed typical hormetic dose response of sesquiterpenes to O<sub>3</sub> (Bourtsoukidis et al., 2012). Similarly, evidence for O<sub>3</sub>-induced hormesis in the emission rate of several VOCs was revealed in tobacco (*Nicotiana tabacum* L. 'Wisconsin') plants which were exposed to 0, 400, 600, 800, and 1000 ppb for 30 min and their VOCs emission rates were measured at 0.5, 3, 10, 24, and 48 h (Kanagendran et al., 2018).


Biphasic isoprene emissions, with maximum in the low-dose zone (Fig. 2A,B,C,D), are particularly important to air quality and environmental health as isoprene is non-specifically stored in the plants and has the largest emission rate among VOCs thus being the major biogenic VOC affecting tropospheric photochemistry. Hence, higher emissions under low-level stress would result in more isoprene in the atmosphere. However, biphasic emissions of VOCs specifically stored in plant tissues are also important to ecological health because reduced emission in the low-dose zone may have implications for plant tolerance against environmental stress and to the interactions among plants, herbivores, and other organisms, therefore with potential consequences to ecosystem sustainability (Ormeño et al., 2011).

The biphasic dose responses of plant-emitted VOCs as noted above reveal hormesis, a highly generalizable dose response phenomenon (Agathokleous, 2018; Calabrese and Blain, 2009, 2011; Calabrese and Mattson, 2011, 2017). The evolutionary basis of hormesis (Agathokleous, 2018; Calabrese and Mattson, 2017; Kim et al., 2018) suggests that plants/VOCs biology and accompanying VOC emissions have evolved within the context of enhancing communication, conditioning, protection, and defense functions.

## 3. Conclusions & perspectives

This commentary promotes new insights into the interface between biological systems and environmental change that influences several parts of the globe, by collectively documenting hormetic-like responses of plant-emitted VOCs to stimuli. Hormetic acting VOCs challenge the environmental pollution research community to consider the ecological implications of low-level stress that may alter VOC emissions. The blend of VOC emissions also warrants further examination within the context of their evolutionary adaptations and how these may be mediated within a dose response framework.

The study of hormetic dose responses will be challenging, requiring stronger study designs with more doses, especially in the low dose zone, and probably with repeat measure-time components. However, the continued ignoring of the possibility of hormesis in ecological system responses by inadequate hazard assessment testing strategies and protocols will provide decision makers with inadequate data upon which to base environment standards. With the use of more rigorous and demanding protocols that permit an evaluation of the entire dose response continuum,



**Fig. 1.** Hypothetical hormetic, threshold, and linear non-threshold (LNT) dose-response models. The five-point star indicates the no-observed-adverse-effect-level (NOAEL) for the threshold and hormesis models. The LNT model assumes that response increases linearly with increasing dose.

Download English Version:

<https://daneshyari.com/en/article/8856513>

Download Persian Version:

<https://daneshyari.com/article/8856513>

[Daneshyari.com](https://daneshyari.com)