


## Accepted Manuscript

Messinian salinity crisis record under strong freshwater input in marginal, intermediate, and deep environments: The case of the North Aegean

Vasileios Karakitsios, Jean-Jacques Cornée, Theodora Tsourou, Pierre Moissette, George Kontakiotis, Konstantina Agiadi, Emmanouil Manoutsoglou, Maria Triantaphyllou, Efterpi Koskeridou, Harikleia Drinia, Dimitrios Roussos



PII: S0031-0182(17)30116-5  
DOI: doi: [10.1016/j.palaeo.2017.06.023](https://doi.org/10.1016/j.palaeo.2017.06.023)  
Reference: PALAEO 8339

To appear in: *Palaeogeography, Palaeoclimatology, Palaeoecology*

Received date: 2 February 2017  
Revised date: 7 June 2017  
Accepted date: 16 June 2017

Please cite this article as: Vasileios Karakitsios, Jean-Jacques Cornée, Theodora Tsourou, Pierre Moissette, George Kontakiotis, Konstantina Agiadi, Emmanouil Manoutsoglou, Maria Triantaphyllou, Efterpi Koskeridou, Harikleia Drinia, Dimitrios Roussos, Messinian salinity crisis record under strong freshwater input in marginal, intermediate, and deep environments: The case of the North Aegean, *Palaeogeography, Palaeoclimatology, Palaeoecology* (2017), doi: [10.1016/j.palaeo.2017.06.023](https://doi.org/10.1016/j.palaeo.2017.06.023)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Messinian salinity crisis record under strong freshwater input in marginal, intermediate, and deep environments: the case of the North Aegean**

KARAKITSIOS Vasileios<sup>1</sup>, CORNÉE Jean-Jacques<sup>2</sup>, TSOUROU Theodora<sup>1</sup>, MOISSETTE Pierre<sup>3,1</sup>, KONTAKIOTIS George<sup>1</sup>, AGIADI Konstantina<sup>1</sup>, MANOUTSOGLOU Emmanouil<sup>4</sup>, TRIANTAPHYLLOU Maria<sup>1</sup>, KOSKERIDOU Efterpi<sup>1</sup>, DRINIA Harikleia<sup>1</sup>, ROUSSOS Dimitrios<sup>1</sup>

<sup>1</sup>National and Kapodistrian University of Athens, Faculty of Geology and Geoenvironment, Department of Historical Geology & Paleontology, 15784, Athens, Greece, vkarak@geol.uoa.gr

<sup>2</sup>Géosciences Montpellier, CNRS, Université de Montpellier, Montpellier, France

<sup>3</sup>Univ Lyon, Université Lyon 1, Ens de Lyon, CNRS, UMR 5276 LGL-TPE, 69622 Villeurbanne, France

<sup>4</sup>Technical University of Crete, School of Mineral Resources Engineering, GR - 731 00 Chania, Greece

**Abstract**

In the present study, we investigate the Mediterranean–Paratethys connection during the late Miocene in Strymon Basin (North Aegean, northeastern Mediterranean) and compare this onshore sequence with the adjacent offshore Prinos–Nestos sequence, before, during, and after the Messinian Salinity Crisis (MSC). Strymon Basin was a peripheral shallow-water basin during the first MSC stage. The Akropotamos sections expose a clastic sequence with gypsum intercalations, which is dated in the Messinian based on the ostracod and calcareous nannofossil assemblages. This sequence records the Primary Lower Gypsum deposition in a shallow marine environment and its passage via the Messinian erosional surface to a brackish environment with changing salinity conditions similar to the Paratethyan depositional environments. The sequence is capped by a travertine marker horizon observed across the entire Strymon Basin indicating freshwater environment. The Miocene–Pliocene transition is characterized by salinity changes caused by the interaction between Atlantic–Mediterranean and Paratethyan waters, predating the marine reflooding at the end of the MSC, which is attested by the overlying Pliocene open marine deposits. The offshore Prinos–Nestos basin incorporates the Nestos intermediate basin and the Prinos intermediate–deep basin. Borehole and seismic profile data from the offshore Prinos–Nestos oil field reflect a thick clastic

Download English Version:

<https://daneshyari.com/en/article/8868658>

Download Persian Version:

<https://daneshyari.com/article/8868658>

[Daneshyari.com](https://daneshyari.com)