ARTICLE IN PRESS

C. R. Acad. Sci. Paris, Ser. I ••• (••••) •••-••



Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I



CRASS1:6118

www.sciencedirect.com

Partial differential equations

Stability for entire radial solutions to the biharmonic equation with negative exponents

Stabilité des solutions radiales entières de l'équation biharmonique avec exposants négatifs

Xia Huang^a, Dong Ye^b, Feng Zhou^a

^a Center for Partial Differential Equations, School of Mathematical Sciences, East China Normal University, Shanghai, 200241, China ^b IECL, UMR 7502, Département de mathématiques, Université de Lorraine, 3, rue Augustin-Fresnel, 57073 Metz, France

ARTICLE INFO

Article history: Received 21 March 2018 Accepted after revision 3 May 2018 Available online xxxx

Presented by Haïm Brézis

ABSTRACT

In this note, we are interested in entire solutions to the semilinear biharmonic equation

 $\Delta^2 u = -u^{-p}, \ u > 0 \quad \text{in } \mathbb{R}^N,$

where p > 0 and $N \ge 3$. In particular, the stability outside a compact set of the entire radial solutions will be completely studied, which resolves the remaining case in [5]. © 2018 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

RÉSUMÉ

Dans cette note, on s'intéresse aux solutions radiales entières de l'équation semilinéaire biharmonique

$$\Delta^2 u = -u^{-p}, \ u > 0 \quad \text{dans } \mathbb{R}^N,$$

où p > 0 et $N \ge 3$. En particulier, on étudie la stabilité en dehors d'un compact des solutions radiales entières, et on résout un cas ouvert dans [5].

© 2018 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

In this note, we are interested in entire radial solutions to the biharmonic equation

$$\Delta^2 u = -u^{-p}, \ u > 0 \quad \text{in } \mathbb{R}^N$$

where p > 0 and $N \ge 3$.

https://doi.org/10.1016/j.crma.2018.05.001

Please cite this article in press as: X. Huang et al., Stability for entire radial solutions to the biharmonic equation with negative exponents, C. R. Acad. Sci. Paris, Ser. I (2018), https://doi.org/10.1016/j.crma.2018.05.001

(1.1)

E-mail addresses: xhuang1209@gmail.com (X. Huang), dong.ye@univ-lorraine.fr (D. Ye), fzhou@math.ecnu.edu.cn (F. Zhou).

¹⁶³¹⁻⁰⁷³X/ $\ensuremath{\mathbb{C}}$ 2018 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

2

ARTICLE IN PRESS

X. Huang et al. / C. R. Acad. Sci. Paris, Ser. I ••• (••••) •••-•••

Recently, the fourth-order equations have attracted the interest of many researchers. In particular, the existence, multiplicity, stability, and qualitative properties of solutions to equation (1.1) are studied in many works, especially for radial solutions. It has been proved in [6] that, if 0 , the equation (1.1) admits no entire smooth solution. It is showed in [4,7] that, for any <math>p > 1, there exist radial solutions to (1.1).

Definition 1. A solution *u* to (1.1) is said stable in $\Omega \subseteq \mathbb{R}^N$ if there holds

$$\int_{\Omega} |\Delta \phi|^2 \mathrm{d} x - p \int_{\Omega} u^{-p-1} \phi^2 \mathrm{d} x \ge 0 \quad \text{for any } \phi \in C_0^{\infty}(\Omega).$$

Moreover, a solution u to (1.1) is said stable outside a compact set K if u is stable in $\mathbb{R}^N \setminus K$. For simplicity, we say also that u is stable if $\Omega = \mathbb{R}^N$.

We consider the following initial value problem

$$\begin{cases} \Delta^2 u = -u^{-p} & \text{for } r \in [0, R_{\alpha,\beta}) \\ u'(0) = u'''(0) = 0, \\ u(0) = \alpha, \quad \Delta u(0) = \beta; \end{cases}$$
(1.2)

for any $\alpha, \beta \in \mathbb{R}$, we denote by $u_{\alpha,\beta}$ the (local) solution to (1.2) and by $[0, R_{\alpha,\beta})$ the maximal interval of existence. Notice that the equation (1.2) is invariant under the scaling transformation

$$u_{\lambda}(x) = \lambda^{-\frac{4}{p+1}} u(\lambda x), \ \lambda > 0.$$

Therefore, we need only to consider the case $\alpha = 1$. We will denote $u_{1,\beta}$ by u_{β} . Let p > 1, it is known from [3,5,7] that

- there is no global solution to (1.2) if $N \leq 2$;
- for $N \ge 3$, there exists $\beta_0 > 0$ depending on N such that the solution to (1.2) is globally defined if and only if $\beta \ge \beta_0$. Furthermore, $\lim_{r\to\infty} \Delta u_{\beta} \ge 0$ and $\lim_{r\to\infty} \Delta u_{\beta} = 0$ if and only if $\beta = \beta_0$;
- for $N \ge 3$, any entire solution u_{β} is stable outside a compact set if $\beta > \beta_0$;
- for N = 4, u_{β_0} is unstable outside every compact set;
- for $5 \le N \le 12$, there exists a critical value $p_N > 1$ (see below for the precise definition) such that, if $1 , <math>u_\beta$ is stable for every $\beta \ge \beta_0$, while for $p > p_N$, there exists $\beta_1 > \beta_0$ such that u_β is stable if and only if $\beta \ge \beta_1$, and u_{β_0} is unstable outside every compact set;
- for $N \ge 13$ and any p > 1, u_{β} is stable for every $\beta \ge \beta_0$.

Moreover, Warnault [8] proved that equation (1.1) admits no stable solution (radial no not) for $N \le 4$. So it remains to consider the eventual stability outside a compact set for N = 3 and $\beta = \beta_0$.

The stability property of entire radial solutions is closely related to their asymptotic behaviors. Let us recall the asymptotic behaviors showed in [2,3,5]. For N = 3 and $\beta = \beta_0$, the following hold:

$$\begin{cases} \lim_{r \to \infty} u_{\beta_0}(r)r^{-1} = \ell > 0, & \text{if } p > 3; \\ \lim_{r \to \infty} u_{\beta_0}(r)r^{-1}(\ln r)^{-\frac{1}{4}} = \sqrt[4]{2}, & \text{if } p = 3; \\ \lim_{r \to \infty} u_{\beta_0}(r)r^{-\frac{4}{p+1}} = \left[-Q_4\left(-\frac{4}{p+1}\right) \right]^{-\frac{1}{p+1}} =: L_0, & \text{if } 1 (1.3)$$

where Q₄ is defined by

$$Q_4(m) := m(m+2)(N-2-m)(N-4-m).$$
(1.4)

Remark that equation (1.1) has a singular solution $u_s(r) \equiv L_0 r^{\frac{4}{p+1}}$, if $Q_4\left(-\frac{4}{p+1}\right) < 0$.

From [2], we know that for N = 3, there exist $3 > p_c^+ > p_c > 1$ such that, if $p = p_c$ or $p = p_c^+$, then $-pQ_4(m) = \frac{9}{16}$ with $m = -\frac{4}{p+1}$, and if $p_c then <math>-pQ_4(m) > \frac{9}{16}$. For $N \ge 5$, p_N is the unique root of

$$-pQ_4\left(-\frac{4}{p+1}\right) = \frac{N^2(N-4)^2}{16}$$

in $(1, \infty)$.

Please cite this article in press as: X. Huang et al., Stability for entire radial solutions to the biharmonic equation with negative exponents, C. R. Acad. Sci. Paris, Ser. I (2018), https://doi.org/10.1016/j.crma.2018.05.001

Download English Version:

https://daneshyari.com/en/article/8905322

Download Persian Version:

https://daneshyari.com/article/8905322

Daneshyari.com