

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/TJEM)

Turkish Journal of Emergency Medicine

journal homepage: <http://www.elsevier.com/locate/TJEM>

Original Article

Factors predicting the need for tube thoracostomy in patients with iatrogenic pneumothorax associated with computed tomography-guided transthoracic needle biopsy

İbrahim Ulaş Özturan ^{a,*}, Nurettin Özgür Doğan ^a, Cansu Alyeşil ^a, Murat Pekdemir ^a, Serkan Yılmaz ^a, Hüseyin Fatih Sezer ^b

^a Kocaeli University, Faculty of Medicine, Department of Emergency Medicine, Kocaeli, Turkey

^b Kocaeli University, Faculty of Medicine, Department of Thoracic Surgery, Kocaeli, Turkey

ARTICLE INFO

Article history:

Received 25 March 2018

Received in revised form

10 May 2018

Accepted 17 May 2018

Available online xxx

Keywords:

Pneumothorax

Iatrogenic disease

Needle biopsy

Chest tubes

Tube thoracostomy

ABSTRACT

Objectives: Traumatic iatrogenic pneumothorax occurs most often after a transthoracic needle biopsy. Since this procedure has become a common outpatient intervention, emergency department admissions of post-biopsy pneumothorax patients have increased. The aim of this study was to determine the factors that predict the need for tube thoracostomy in patients with post-biopsy pneumothorax in the emergency department.

Methods: A retrospective cross-sectional study was conducted on 191 patients with post-biopsy pneumothorax who were admitted to the emergency department between 2010 and 2017. Patient characteristics, clinical findings at the emergency department presentation, and procedural and radiological features were reviewed. A multivariate logistic regression model was constructed using the variables from univariate comparisons to determine the need for tube thoracostomy in patients with iatrogenic pneumothorax, and the effect sizes were demonstrated with odds ratios.

Results: Tube thoracostomies were performed on 69 out of 191 patients (36.1%). A total of 122 patients (63.9%) were treated with supplemental oxygen therapy without any other intervention, and 126 patients (66.0%) were hospitalized. In the multivariate model, the variables predicting the need for a tube thoracostomy were decreased breath sounds, dyspnea, decreased systolic blood pressure, decreased oxygen saturation and increased pleura–lesion distance. A distance of 19.7 mm predicted the need with a sensitivity of 69.6% and a specificity of 62.3%.

Conclusion: Decreased breath sounds, dyspnea, decreased systolic blood pressure, decreased oxygen saturation, and increased pleura–lesion distance may predict the need for a tube thoracostomy in patients with post-biopsy pneumothorax.

Copyright © 2018 The Emergency Medicine Association of Turkey. Production and hosting by Elsevier B.V. on behalf of the Owner. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Pneumothorax is classified as either spontaneous or traumatic

depending on its etiology.^{1,2} The term “iatrogenic pneumothorax” (IP) is used when the pneumothorax results from an interventional procedure,³ including such common procedures such as transthoracic needle biopsy (TTNB; 24% of all IP’s), subclavian vein catheterization (22%), and thoracentesis (20%).⁴ Therefore, the incidence of IP is even higher than that of spontaneous pneumothorax.⁵ As the number of diagnostic or therapeutic procedures increases, the rate of IP increases. In order to better manage this condition, there is a need for intervention methods based on the clinical signs of IP.

TTNB is usually performed as an outpatient procedure, which is a safe procedure after which the patient can be discharged,

* Corresponding author.

E-mail addresses: ozturan.iu@gmail.com (İ.U. Özturan), nurettinozgurdogan@gmail.com (N.Ö. Doğan), drcansualyeshil@gmail.com (C. Alyeşil), mpekdemir@yahoo.com (M. Pekdemir), syilmazmd@gmail.com (S. Yılmaz), hfs.hfs@gmail.com (H.F. Sezer).

@dr_ozturanMD (İ.U. Özturan)

Peer review under responsibility of The Emergency Medicine Association of Turkey.

<https://doi.org/10.1016/j.tjem.2018.05.002>

2452–2473/Copyright © 2018 The Emergency Medicine Association of Turkey. Production and hosting by Elsevier B.V. on behalf of the Owner. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Özturan İU, et al., Factors predicting the need for tube thoracostomy in patients with iatrogenic pneumothorax associated with computed tomography-guided transthoracic needle biopsy, Turkish Journal of Emergency Medicine (2018), <https://doi.org/10.1016/j.tjem.2018.05.002>

following a short observation period.⁶ However, large surveys have shown that the most common complication of TTNB is pneumothorax,^{7,8} for which patients may require follow-up in an observation unit or ward. The patients may also be admitted to the emergency department (ED) if any symptoms present after discharge. Although most pneumothoraces are resolved without any intervention other than supplemental oxygen therapy, a small number of pneumothoraces can expand and may result in a tension pneumothorax,⁹ in which case immediate intervention is necessary before a cardiovascular collapse develops.¹⁰ Therefore, the early recognition of an expanding pneumothorax is of critical importance in order to be prepared for the necessary intervention.

The American College of Chest Physicians in 2001 and the British Thoracic Society in 2010 published guidelines for the management of spontaneous pneumothorax.^{5,11} Though the guidelines offer diverse recommendations, none of the guidelines address the management of IP. Consequently, the management of IP is still being debated. Therefore, to define the patient population who undergoes post-TTNB tube thoracostomy, we aimed to determine the predictive factors for tube thoracostomy in patients with postbiopsy pneumothorax.

2. Methods

2.1. Study design

This retrospective cross-sectional study was conducted in an academic ED serving approximately 45,000 patients per year between January 2010 and January 2017. The institution is also an interventional radiology center that performs about 4000 procedures per year. Institutional review board approval was obtained for the study and informed consent was waived.

2.2. Study setting and population

The study population included all patients older than 18 years who presented to the ED with post-TTNB pneumothorax. At our institution, patients with post-TTNB pneumothorax are referred to the ED for further evaluation and treatment. The exclusion criteria were patients who had an inpatient TTNB, patients who were not referred to the ED after TTNB, and patients who had missing data in their archive records.

According to our ED management, tube thoracostomy decision is made based on patient's clinical and radiological findings in conjunction with thoracic surgery department. All the patients who had a tube thoracostomy performed were admitted to the thoracic surgery ward. All the patients who had a small pneumothorax without symptoms were observed in the ED. After 6 h of observation, if no pneumothorax was seen on the control chest x-ray, the patients were discharged from the ED with instructions. Patients were admitted to the wards if the pneumothorax was enlarged or no improvement was seen during the ED observation.

2.3. Study protocol

Patient characteristics including age; gender; medical history; tobacco use; presence of COPD or malignancy; clinical findings which were documented by an emergency physician at the initial ED presentation including vital signs, symptoms, and physical examination features; ED observation duration or length of stay (LOS) at the hospital; procedural features including fine-needle or core biopsy; and radiological features including lesion size, cavitation of the lesion, pleura–lesion distance, fissure or atelectasis in the affected lung, and lesion location (central or peripheral) were reviewed from the hospital's archive records using international

classification of disease 9 (ICD-9) codes in conjunction with the hospital's imaging database. Computed tomography (CT) images were evaluated by a radiologist who was blind to the patient outcomes. All the variables that were available at the initial ED presentation were evaluated to predict the need for tube thoracostomy.

2.4. Outcome measures

The primary outcome measure was the definition of the patient population that may require tube thoracostomy after TTNB according to demographic parameters such as age, gender and past medical history; clinical parameters at the initial ED presentation such as symptoms, vital signs and physical examination findings and radiological parameters such as pleura–lesion distance or lesion characteristics. The secondary outcome measures were defined as the hospitalization rate and the hospital LOS.

2.5. Data analysis

All statistical analyses were performed with SPSS for Windows (version. 15.0, SPSS Inc., Chicago, IL, USA). The Kolmogorov–Smirnov test was used to test for a normal distribution of the variables. A logistic regression model was constructed to assess the factors predicting the need for tube thoracostomy in patients with IP. The fitness of the multivariable regression model was evaluated with the Hosmer–Lemeshow test. The univariate model included demographic characteristics, radiographic records, and the variables that were available upon ED presentation. Each variable was tested in the univariate model, and those that were significant at an alpha level of 0.2 were then tested in the multivariate model. Odds ratios (ORs) were presented with 95% confidence intervals (CIs). An ROC curve analysis was also performed to determine which pleura–lesion distance could predict the need for tube thoracostomy.

3. Results

Between 2010 and 2017, 306 patients had complications of pneumothorax, and 115 patients were excluded (Fig. 1). A total of 191 patients were included in the study. Tube thoracostomy was performed on 69 out of 191 patients (36.1%). A total of 122 patients (63.9%) were treated with supplemental oxygen therapy without any other intervention, and 126 patients (66.0%) were hospitalized. A total of 69 patients underwent tube thoracostomy; fine needle aspiration (FNA) was performed in 47 patients, core biopsies were performed in 10 patients, and both FNA and core biopsies were performed in 12 patients.

The overall incidence of pneumothorax was 12% (306/2445), and the tube thoracostomy rate was 3% (69/2330) of all the TTNB cases in our study after excluding the 115 patients.

Demographic features of the patients with post-TTNB pneumothorax and the univariate analyses of the variables are shown in Table 1. The demographic and radiological variables that were determined to be significant predictors of the need for tube thoracostomy by univariate analysis were included in the multivariate logistic regression.

The results of the multivariate logistic regression analysis to determine the factors predicting the need for tube thoracostomy are shown in Table 2. The Hosmer–Lemeshow test revealed that our regression results were fit ($p = 0.128$). According to the multivariate regression, decreased breath sounds (OR = 5.6, 95% CI = 2.4–12.8, $p < 0.001$), dyspnea (OR = 2.7, 95% CI = 1.2–6.2, $p = 0.015$), decreased SBP (OR = 0.9, 95% CI = 0.9–0.9, $p = 0.04$), decreased oxygen saturation (OR = 0.9, 95% CI = 0.7–0.9, $p = 0.011$), and

Download English Version:

<https://daneshyari.com/en/article/8950269>

Download Persian Version:

<https://daneshyari.com/article/8950269>

[Daneshyari.com](https://daneshyari.com)