

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

The Egyptian Heart Journal

journal homepage: www.elsevier.com/locate/ehj

Original Article

Waist height ratio and waist circumference in relation to hypertension, Framingham risk score in hospitalized elderly Egyptians [☆]

Heba M. Tawfik

Geriatric Medicine and Gerontology Department, Ain-Shams University, Cairo, Egypt

ARTICLE INFO

Article history:

Received 1 August 2017

Accepted 29 December 2017

Available online xxxxx

Keywords:

Hypertension
Waist to height ratio
Waist circumference
Framingham risk
Egyptians

ABSTRACT

Background: Hypertension is a public health problem and obesity is becoming an epidemic, increasing the risk of hypertension. Both are risk factors for cardiovascular diseases (CVD).

Methods: A case control study recruiting 102 patients aged ≥ 60 years, divided into 55 cases with hypertension and 47 controls without. Body mass index (BMI), waist circumference (WC) and waist to height ratio (WHtR) were measured as well as lipid profile then Framingham risk score (FRS) was calculated.

Results: Odds ratio (OR) for hypertension and medium to high risk cardiovascular events was the same in female patients using WC and WHtR. In male patients, only WHtR increased the risk for hypertension and for cardiovascular events, OR significantly increased with higher WHtR compared to WC.

Conclusion: WHtR and WC are strong risk factors for hypertension and cardiovascular events in Egyptian elderly female patients. WHtR is the best anthropometric predictor for hypertension and cardiovascular events in male patients.

© 2018 Egyptian Society of Cardiology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Hypertension prevalence is increasing, constituting a major public health problem. It is considered one of the most common chronic medical conditions in the United States, affecting about one third of the population.¹ It is a major risk factor for cardiovascular and cerebrovascular diseases, increasing morbidity and mortality. Therefore, prevention of hypertension is a public health challenge.² In Egypt, according to Egyptian National Hypertension Project (ENHP) 1991–1993, about 26.3% of adult Egyptians had high blood pressure, and more than 50% of people older than 60 years suffered from hypertension.³ Obesity and dyslipidemia are associated with hypertension, and visceral adiposity accounts for 65% to 75% of the risk for essential hypertension. All are major risk factors for coronary artery disease (CAD).^{4,5} Egyptians have one of the highest mortality rates worldwide attributed to CAD, further complicating this problem.⁶ Both generalized and abdominal obesity are associated with increased risk of morbidity and mortality. BMI was traditionally the chosen indicator measuring body composition. However, alternative measures that reflect visceral adi-

posity, such as waist circumference and waist–hip ratio (WHR), have been suggested to be superior in predicting CVD risk, because increased visceral adipose tissue is associated with metabolic abnormalities and inflammation.^{5,7} WHtR is considered in some studies now a proxy for central adipose tissue, and has recently been described as a marker of ‘early health risk’.⁸ The Framingham coronary heart disease (CHD) risk assessment tool has been validated and used widely to determine the 10 year risk of cardiovascular events.⁹ Population-based research focuses on the relationship between hypertension and obesity and abdominal obesity, but gender-specific approaches are less common¹⁰ and mostly no studies were done in Egypt approaching this issue.

2. Aim of the work

To study the relation between anthropometric measures (waist height ratio and waist circumference) and both the risk hypertension and Framingham risk score for cardiovascular diseases in male and female elderly Egyptians.

3. Study design

A case control study performed in Ain-Shams University hospital during the period between June and September 2016 after taking informed consent from patients to participate in the study.

[☆] The protocol for the research project has been approved by Ethics Committee of the Faculty of Medicine, Ain-Shams University.

Peer review under responsibility of Egyptian Society of Cardiology.

E-mail addresses: hmt_82@yahoo.com, hebatawfik82@med.asu.edu.eg

<https://doi.org/10.1016/j.ehj.2017.12.008>

1110-2608/© 2018 Egyptian Society of Cardiology. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

4. Participants

One hundred and two patients aged 60 years and more were recruited. Patients were divided into 55 cases and 47 controls according to the presence or absence of hypertension. Patients were considered to have hypertension if systolic blood pressure was ≥ 140 mmHg or diastolic blood pressure was ≥ 90 mmHg or both on two different occasions after complete physical or mental rest, or patients previously informed they have hypertension or on treatment for hypertension.

5. Exclusion criteria

Patients known to have diabetes mellitus, ischemic heart disease either on anti-ischemic treatment or experienced previous cardiac events were excluded from the study.

6. Methods and assessment

Resting blood pressure was measured in a sitting position after a 5-min of mental and physical rest using a mercury sphygmomanometer and according to standard procedures.¹¹

The first objective: to study the relation between anthropometric measures, lipid profile and hypertension. Anthropometric measurements were taken by a trained staff, in the morning, according to the world health organization (WHO) recommendations.¹² Weight was measured to the nearest 0.5 kg, height was measured with the patient barefoot in the standing position to the nearest 0.5 cm. BMI was calculated by dividing weight per kilograms by height per meter square (kg/m^2). WC was measured to the nearest centimeter, midway between lowest rib margin and iliac crest with the patient standing and breathing normally. It was used as a surrogate for central obesity. WC was categorized into high-risk (≥ 94 cm and ≥ 80 cm for males and females, respectively) and low-risk if below these cutoffs as established standards by WHO and Interna-

tional diabetes federation (IDF) for patients living the Middle East; including Egypt.¹³ WHtR was calculated as WC (in cm) divided by height (in cm). A ratio of (0.5) was considered a cutoff point for WHtR; according to a systematic review done in 2010, which stated that this cutoff can be used for men and women across different ethnic groups.¹⁴

Blood samples were obtained for lipid profile from patients by venipuncture after 12 h of fasting. Blood samples were assayed within 24 h.

The second objective: FRS for cardiovascular events⁹ was calculated; to study its relation to WC, BMI, WHtR. It takes into consideration the following factors: age; gender; systolic blood pressure value; whether or not the patient takes anti-hypertensive medications; the presence or absence of diabetes mellitus; smoking; high-density lipoproteins cholesterol (HDL-c) and total cholesterol values. Patients are considered to have low 10 year risk for cardiovascular events if FRS is less than 10%, intermediate-risk if between 10–20%, and high risk if 20% or more.¹⁵ Patients were classified in the study into two groups, a group with low risk (FRS <10%) and a group with intermediate to high risk (FRS $\geq 10\%$) and OR was calculated accordingly.

7. Statistical analysis

Age, BMI, WC and Lipid profile were studied in relation to cases and controls by Paired sample t test. Odds ratio regarding BMI, WHtR and WC were calculated using Chi square test. Values for P less than 0.05 were considered significant and values less than 0.001 were considered highly significant.

8. Results

A total of 102 patients divided into 55 cases (patients already having hypertension) and 47 controls (patients with normal blood pressure) as shown in Table 1. The study included 70 male (about

Table 1
Demography of the study population.

		Hypertension				X ²	P
		No controls number = 47		Yes cases number = 55			
		Count	N%	Count	N%		
Sex	Male N = 70	35	74.5%	35	63.6%	1.101	.294
	Female N = 32	12	25.5%	20	36.4%		
Smoking	No	25	53%	37	67%	4.453	.056
	Yes	22	47%	18	33%		

Table 2
Comparison between cases and controls regarding age, anthropometric measures and lipid profile.

	Controls/cases	N	Mean	Std. Deviation	t test	p
Age	Controls	47	67.94	6.664	0.682	.497
	Cases	55	67.11	5.490		
BMI	Controls	47	29.60	3.048	-2.376	.019
	Cases	55	31.37	4.257		
WC	Controls	47	93.40	8.9838	-2.140	.036
	Cases	55	98.05	13.3256		
Total Cholesterol	Controls	47	131.26	44.647	-2.327	.022
	Cases	55	154.72	56.565		
TG	Controls	47	114.79	38.483	-2.810	.006
	Cases	55	141.02	54.814		
HDL	Controls	47	45.00	5.469	2.251	.027
	Cases	55	42.50	5.649		
LDL	Controls	47	91.00	19.123	-2.332	.022
	Cases	55	100.59	21.829		

BMI = body mass index WC = waist circumference TG = triglycerides.
LDL = low density lipoproteins HDL = high density lipoproteins.

Download English Version:

<https://daneshyari.com/en/article/8957073>

Download Persian Version:

<https://daneshyari.com/article/8957073>

[Daneshyari.com](https://daneshyari.com)