## Accepted Manuscript

Title: Local canopy diversity does not influence phenotypic expression and plasticity of tree seedlings exposed to different resource availabilities

Authors: Cristina C. Bastias, Fernando Valladares, Natalia Ricote M., Raquel Benavides

PII: S0098-8472(18)30639-7

DOI: https://doi.org/10.1016/j.envexpbot.2018.08.023

Reference: EEB 3549

To appear in: Environmental and Experimental Botany

Received date: 26-4-2018 Revised date: 14-8-2018 Accepted date: 17-8-2018

Please cite this article as: Bastias CC, Valladares F, Ricote M. N, Benavides R, Local canopy diversity does not influence phenotypic expression and plasticity of tree seedlings exposed to different resource availabilities, *Environmental and Experimental Botany* (2018), https://doi.org/10.1016/j.envexpbot.2018.08.023

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## ACCEPTED MANUSCRIPT

Local canopy diversity does not influence phenotypic expression and plasticity of tree seedlings exposed to different resource availabilities.

**Author list:** Cristina C. Bastias <sup>a\*</sup>, Fernando Valladares <sup>a,b</sup>, Natalia Ricote M. <sup>c,d</sup> & Raquel Benavides <sup>a</sup>

- <sup>a</sup> Departamento de Biogeografía y Cambio Global. Museo Nacional de Ciencias Naturales, MNCN-CSIC, Madrid, Spain.
- <sup>b</sup> Área de Biodiversidad y Conservación, Universidad Rey Juan Carlos, Móstoles, Madrid, Spain.
- <sup>c</sup> Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile. Santiago, Chile.
- <sup>d</sup> Center of Applied Ecology & Sustainability (CAPES), Pontificia Universidad Católica de Chile, Santiago 6513677, Chile

### Highlights

- Similar plasticity for tree seedlings from communities differing in canopy richness
- Plants from the Mediterranean forest showed great plasticity to resource supply
- Differential plastic responses of functional groups in the Mediterranean forest
- Oaks have a higher capacity response to changing conditions than pines
- Nutrients affected most of the traits, while light only at leaf morphology level

<sup>\*</sup>Corresponding author, crbasc@gmail.com

#### Download English Version:

# https://daneshyari.com/en/article/8959438

Download Persian Version:

https://daneshyari.com/article/8959438

<u>Daneshyari.com</u>