




ELSEVIER

Contents lists available at ScienceDirect

Journal of Obsessive-Compulsive and Related Disorders

journal homepage: www.elsevier.com/locate/jocrd

Introduction: A global perspective on unwanted intrusive thoughts

David A. Clark^a, Adam S. Radomsky^{b,*}^a University of New Brunswick, Canada^b Concordia University, Canada

ARTICLE INFO

Article history:

Received 15 January 2014

Received in revised form

9 February 2014

Accepted 11 February 2014

Available online 18 February 2014

Keywords:

Unwanted intrusive thoughts

Obsessions

Cross-cultural differences

Cognitive behavioral theory of obsessions

OCD

ABSTRACT

Because of [Rachman and de Silva's \(1978\)](#) influential research on normal and abnormal obsessions, it is now a basic tenet of cognitive behavioral theories (CBT) that clinical obsessions have their origins in the normal intrusive thought phenomena that characterizes the stream of consciousness. However much of the empirical research on unwanted intrusive thoughts has utilized retrospective self-report questionnaires that have weak construct validity and the samples have been drawn primarily from Western European and North American populations. To enhance the measurement precision and investigate the universality of unwanted intrusions, a structured intrusive thoughts interview was developed and administered to 777 nonclinical individuals drawn from 13 countries. The three papers in this special issue present findings based on this large data set. Together it was found that unwanted intrusive thoughts are reported by the majority of individuals in all countries, that significant cross-cultural differences are apparent in primary intrusive thought content, that faulty appraisals and confrontational control strategies are related to the distress of intrusions, and that the frequency of dirt/contamination, doubt, and miscellaneous intrusions are specifically related to obsessive compulsive symptom distress. The authors discuss these findings in terms of their consistency with predictions derived from the CBT perspective on obsessions. The special issue concludes with a discussion paper by Professor Jack Rachman, originator of the obsessional intrusions concept.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

This year marks the 36th anniversary of [Rachman and de Silva's \(1978\)](#) seminal article that introduced the concept of unwanted intrusive thoughts as a potential etiological factor in the development of obsessive compulsive disorder (OCD). Their paper reported on three studies that sought to determine whether healthy, non-clinical individuals experience unwanted thoughts, images or impulses that are similar in content to clinical obsessions (i.e., thoughts of contamination, doubt, aggression, sexuality, etc.). In the first study based on an intrusive thoughts questionnaire 99/124 (80%) nonclinical individuals reported fairly frequent unwanted thoughts or images involving obsessional content, although they were considered fairly easy to dismiss. The second study consisted of a detailed interview of intrusive thoughts conducted on 40 nonclinical individuals and 8 patients with OCD. A list of 23 obsessions from the clinical sample and 58 obsessive-like intrusions from the nonclinical individuals were reproduced in the article and have since become the basis of numerous self-report measures of intrusive thoughts.

Comparison between the clinical and nonclinical samples revealed similarities in thought content, although the OCD patients reported greater frequency, discomfort and difficulty controlling their obsessions. In the third study OCD patients and nonclinical participants were asked to repeatedly form their obsession or intrusive thought. Habituation was evident in both samples, although the nonclinical group had more difficulty forming their intrusion upon request than the OCD patients. Although [Rachman and de Silva's \(1978\)](#) findings raised many questions about the nature of intrusive thoughts, it also provided the first empirical evidence that obsessions might have their origins in the normal thought processes that characterize the stream of consciousness.

The conceptual significance of “normal obsessions” was subsequently emphasized in [Rachman and Hodgson's \(1980\)](#) text, *Obsessions and Compulsions*, which in many respects provides the theoretical underpinnings for contemporary cognitive-behavioral (CB) models of OCD. In their chapter entitled *An Anatomy of Obsessions*, the occurrence of unwanted, unacceptable intrusive thoughts or images initiated a process of faulty meta-cognitive appraisal and control efforts that could spiral into the development of obsessions. Later publications elaborating on a CBT model of obsessions and compulsions also reiterated the occurrence of mental intrusions as a key determinant in the pathogenesis of obsessions (e.g., [Clark, 2004](#); [Freeston, Rhéaume, & Ladouceur, 1996](#); [Rachman, 1997, 1998](#);

* Correspondence to: Department of Psychology, Concordia University, 7141 Sherbrooke St. West, Montreal, QC, Canada.

E-mail address: adam.radomsky@concordia.ca (A.S. Radomsky).

Salkovskis, 1985, 1989). Today the universality of obsession-related intrusive thoughts, images and impulses is a central tenet of CBT explanations for the development of obsessions.

While the conceptual significance of the [Rachman and de Silva \(1978\)](#) publication cannot be overstated, their work has also had a significant impact on empirical research on unwanted intrusions. In the last three decades research on intrusions has taken two broad pathways; the most common approach is questionnaire based, whereas the other is more experimental, examining the impact of intentional mental control efforts on unwanted intrusive thoughts. These two broad research pathways are represented in the first and third studies reported in the [Rachman and de Silva \(1978\)](#) publication. Interestingly there has been little research that has employed the interview format; a neglect that we think is serious given the complexity of intrusive thoughts.

There are important conceptual and methodological weaknesses that have been raised with questionnaire and experimental measurement of unwanted intrusive thoughts. Questionnaire measures of intrusions present individuals with a predetermined list of thought statements which respondents are then instructed to rate for frequency of occurrence. However concerns have been raised about the veridicality of these item responses ([Brown & Clark, 2014](#)). When individuals indicate that they frequently experience an unwanted intrusive thought by endorsing an item statement, does this reflect their actual memory of specific thought occurrences, or might it reflect an emotional state or self-identity that is congruent with the item content ([Glass & Arnkoff, 1997](#))? In addition the content validity of many intrusive thought questionnaires is problematic because they can contain a broad range of negative thought content that often includes general anxiety and depressive thought content ([Clark & Purdon, 1995](#)). As well the lists of intrusions are always predetermined so how well these pre-established items represent the intrusive thought content of specific individuals is doubtful. [Rachman \(1981\)](#) has noted that unwanted mental intrusions are often quite idiosyncratic and triggered by external cues. Questions have also been raised about the specificity and discriminative validity of intrusive thoughts questionnaires ([Julien, O'Connor, & Aardema, 2007](#)). These are fundamental questions about the construct validity of retrospective self-report of unwanted intrusions that leave us uncertain about the interpretability of a high score on such measures. When a person indicates that s/he frequently has an intrusion (e.g., "I have unwanted intrusive thoughts, images or impulses of verbalizing something rude or embarrassing that would hurt a person's feelings"), is this based on memories of actual thought occurrences, recalling times when feeling anxious about one's verbal interaction, a single recent experience of "embarrassed verbalization", or a self-identity that acknowledges you are the type of person who is concerned about your impact on others? In sum questionnaires may not be the most accurate method for assessing people's actual experience of unwanted intrusive thoughts.

Unfortunately experimental approaches to the investigation of unwanted mental intrusions and their control also have significant drawbacks. The first problem is ecological validity. When an individual is brought into a laboratory setting and asked to produce an intrusive thought, it is no longer intrusive. The intentional production of the thought is no longer unwillful and so lacks generalizability to the spontaneous, unintended intrusions that occur in the natural world. As well, a diffusion of responsibility occurs such that the participant can now attribute responsibility to the experimenter for the generation of the intrusion rather than the self. This transfer of responsibility to the experimenter could significantly affect the participant's appraisal of the intrusive thought. Even if the intrusion is induced, difficulties arise because it may be difficult to ensure that the induction was

successful across all research participants. Finally, the control of unwanted negative thoughts under controlled laboratory conditions for very brief time periods will be quite different from the experience of repeated unwanted intrusions over an extended period of time in a context-rich, everchanging naturalistic setting. Thus the findings derived from laboratory-contrived intrusive phenomena raise questions about their relevance to the spontaneous, distressing intrusive thoughts, images and impulses that can "haunt" individuals in their everyday life.

2. Current research objectives

The present research was initiated to address two major shortcomings in the intrusive thoughts literature. From a methodological perspective, our first objective was to explore the use of a structured interview format to assess individuals' experience of unwanted intrusive thoughts, images and impulses specifically related to the major themes of OCD; dirt/contamination, doubt, harm/injury, sex, religion/morality, and minor or miscellaneous themes. [Though not normally represented in obsessional fears, we also assessed intrusive thoughts about being a victim of violence because these thoughts may be common in nonclinical populations]. The use of an interview would allow us to collect qualitative data to ensure that participants reported on intrusive thoughts relevant to OCD rather than a range of negative thoughts that might be more relevant to worry, dysphoria or anger. In addition the qualitative information would help interpret the quantitative data on the phenomenology of intrusions and their sequelae (see below). Trained interviewers were provided with strict definitions and examples of obsession-relevant intrusive thoughts and participants' responses were written down for subsequent verification that the intrusion fit the study criteria. In addition the interview ensured that participants responded to the six types of obsessive intrusive thoughts and the interviewer could probe to ensure that participants really did experience an intrusion. Another advantage of the interview is that participants were asked to describe specific instances of the unwanted intrusion. This increased the likelihood that endorsement reflected actual thought occurrences rather than personal conjectures on the types of thoughts a person experiences. Thus our intention was to develop an interview that would provide a more accurate measurement of intrusive thought content highly specific to OCD.

A second objective of the interview method was to provide a more comprehensive and detailed assessment of individual's appraisal and perceived control of their most distressing intrusive thought. First, general ratings of frequency, distress and control were collected on the six OCD-relevant intrusions. Then individuals selected their most distressing intrusion and detailed ratings of several appraisal and control constructs were obtained. This part of the interview utilized an endorsement approach with participants providing their own ratings on appraisal and control dimensions. Six point rating scales were used so individuals could rate their level of agreement to specific questions about their appraisal of the most distressing intrusion and frequency of using various mental control strategies. Our intent was to provide the participant with appraisal statements that accurately reflect the most common evaluative constructs in contemporary CBT theories of OCD, while keeping them broad enough to encompass a wide range of intrusive phenomena.

In addition to these methodological considerations, we also wanted to investigate the cross-cultural expression of unwanted intrusive thoughts. The universality of unwanted intrusive thoughts is a central assumption of CBT theories of obsessions ([Rachman, 2003](#)). However, there are no published studies that have compared the occurrence of unwanted intrusions across

Download English Version:

<https://daneshyari.com/en/article/912272>

Download Persian Version:

<https://daneshyari.com/article/912272>

[Daneshyari.com](https://daneshyari.com)