



Blood-Based Biomarkers

Stability of blood-based biomarkers of Alzheimer's disease over multiple freeze-thaw cycles

Ashvini Keshavan^{a,b,*}, Amanda Helsegrave^{b,c}, Henrik Zetterberg^{b,c,d,e}, Jonathan M. Schott^a

^aDementia Research Centre, Box 16 National Hospital for Neurology and Neurosurgery, London, UK

^bLeonard Wolfson Biomarkers Laboratory, Department of Molecular Neuroscience, UCL Institute of Neurology, London, UK

^cThe DRI Fluid Biomarker Laboratory, UK Dementia Research Institute at UCL, London, UK

^dDepartment of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Sahlgrenska University Hospital, Mölndal, Sweden

^eClinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

Abstract

Introduction: Freeze-thaw instability may contribute to preanalytical variation in blood-based biomarker studies. We investigated the effects of up to four freeze-thaw cycles on single molecule array immunoassays of serum neurofilament light chain and plasma total tau, amyloid beta 1–40 (A β ₄₀), and A β 1–42 (A β ₄₂).

Methods: Individuals who had peripheral venepuncture during investigation of suspected neurodegenerative disease were recruited. After standardized preprocessing, 200 μ L of plasma and serum aliquots were stored at -80°C within 60 minutes. Aliquots underwent one to four freeze-thaw cycles.

Results: There was no significant difference across four freeze-thaw cycles for serum neurofilament light chain ($n = 12$), plasma total tau ($n = 11$), or plasma A β ₄₂ ($n = 12$). For plasma A β ₄₀ ($n = 14$), there were significant median reductions by ratios of .96 and .92 at the third and fourth cycles, respectively.

Discussion: Up to four freeze-thaw cycles do not influence single molecule array blood biomarkers of neurofilament light chain, total tau, or A β ₄₂, with at most minor reductions in A β ₄₀.

© 2018 Published by Elsevier Inc. on behalf of the Alzheimer's Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords:

Freeze-thaw; Blood; Biomarker; Simoa

1. Introduction

Preanalytical factors may be an important source of variation among studies investigating putative blood-based biomarkers of Alzheimer's disease, in which replication across studies has posed a significant challenge. Guidelines for standardizing these preanalytic variables have stated that it is desirable to minimize the number of freeze-thaw cycles to which samples are exposed [1]. The evidence for this comes predominantly from studies showing that more than three freeze-thaw cycles reduce plasma amyloid beta 1–40 and 1–42 (A β ₄₀ and A β ₄₂) concentrations by as much as

20% [2] and also reduce cerebrospinal fluid (CSF) A β ₄₂ concentration [3] as quantified on immunoassay platforms. There is some conflicting evidence on the stability in CSF of total tau, which in one report remained stable over up to six freeze-thaw cycles [3], but in another study, both CSF total tau and phospho-tau-181 concentrations as measured by ELISA reduced over three freeze-thaw cycles [4]. Conversely, CSF neurofilament light chain (NFL) remains stable over up to four freeze-thaw cycles [5]. However, there has been no systematic examination of the stability of these biomarkers in blood. More recently, the Single molecule array (Simoa) HD-1 analyzer, a highly sensitive digital immunoassay platform, has been developed, which is capable of measuring these and other biomarkers with sensitivity in the femtomolar range [6]. In this study, we examined the effects of up to four freeze-thaw cycles on serum NFL and plasma total tau, A β ₄₀, and A β ₄₂.

The authors have declared that no conflict of interest exists.

*Corresponding author. Tel.: (+44) 2034 483011; Fax: ■■■■.

E-mail address: a.keshavan@ucl.ac.uk

<https://doi.org/10.1016/j.dadm.2018.06.001>

2352-8729/© 2018 Published by Elsevier Inc. on behalf of the Alzheimer's Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2. Methods

2.1. Participants

We recruited individuals seen at the specialist cognitive disorders clinics at the National Hospital for Neurology and Neurosurgery, who were having diagnostic lumbar puncture with paired venous blood testing for investigation of a suspected neurodegenerative condition. At the time of clinical sampling, all individuals donated an additional blood sample for research, and basic demographic data were recorded. All participants gave written informed consent, or for those who were deemed lacking in capacity to do so by the assessing clinician, proxy consent was obtained from a consultee. The study was conducted in accordance with local clinical research regulations and was approved by the local ethics committee (reference: 12/0344).

2.2. Sample handling

The plasma was collected in 8-mL EDTA tubes and serum in 6.5-mL SST gel separator tubes by peripheral venepuncture with a tourniquet, using either a 21G or 23G butterfly needle and BD Vacutainer® collecting system. Participants were not instructed to fast, but all blood samples for each participant were taken between 9 AM and 1 PM on a single venepuncture. Plasma and serum were centrifuged at $1800 \times g$ for 5 minutes at room temperature. Plasma was centrifuged within a median of 20 minutes after sampling, whereas serum was allowed for a median of 15 minutes to clot before centrifugation. The supernatant was pipetted into 200- μ L aliquots into identical polypropylene tubes, for freezing at -80°C , within 60 minutes of sampling, in keeping with current guidelines [1].

2.3. Freeze-thaw cycles

Each additional freeze-thaw cycle after the first consisted of 1 hour of direct thaw to room temperature, followed by 1 hour of direct replacement into -80°C . After the final thaw, samples were transferred to 1.5-mL polypropylene centrifuge tubes and centrifuged as per the kit-manufacturer's recommendation at $13,000 \times g$ for 10 minutes, and the supernatant of each sample was pipetted onto the plate for analysis. For each biomarker, all aliquots from the same participant were assayed using the same batch of reagents in the same run.

2.4. Blood fraction choice

To examine the effects of freeze-thaw cycle number, we chose to measure NFL in serum as we and others have shown that NFL may be measured reliably in serum to show differences between disease groups among neurodegenerative diseases [7,8] and that there is a linear relationship between plasma and serum NFL within individuals, with serum returning higher values when paired samples are analyzed on the same run (Supplementary Figure S1A). Similarly, we chose to measure total tau, $\text{A}\beta_{40}$, and $\text{A}\beta_{42}$ in plasma as it returns higher values than serum for these biomarkers (Supplementary Figure S1B–D).

2.5. Analysis

Commercially available Simoa total tau, NFL, $\text{A}\beta_{40}$, and $\text{A}\beta_{42}$ kits (Quanterix) were used according to the manufacturer's instructions. The principles of these assays are similar; a detailed account of the basis of Simoa technology has been provided by Wilson and colleagues [6].

All measurements were conducted on the same automated HD-1 analyzer (Quanterix). For serum NFL, plasma $\text{A}\beta_{40}$, and plasma $\text{A}\beta_{42}$, all samples were assayed in duplicate. For plasma tau, all samples were assayed in triplicate because of kit availability. The mean of the replicates for each sample was calculated and represented graphically. Results were included in the analysis if the coefficient of variation across replicates was $<15\%$. The number of individuals included was therefore as follows: serum NFL, 12; plasma total tau, 11; plasma $\text{A}\beta_{40}$, 14; and plasma $\text{A}\beta_{42}$, 12.

2.6. Statistics

A Friedman ANOVA test (nonparametric repeated measures) was used to assess differences within individuals across four freeze-thaw cycles (SPSS Statistics, version 24). Where statistically significant differences were found at the $P = .05$ level, the median ratio of the concentration at a specific freeze-thaw cycle to the concentration at the first cycle was calculated across individuals.

3. Results

Fig. 1 shows the measured concentration of each analyte against the number of freeze-thaw cycles.

The Friedman ANOVA test showed a two-tailed significance of $>.05$ for a change in the means for serum NFL, plasma total tau, and plasma $\text{A}\beta_{42}$ over four cycles.

For plasma $\text{A}\beta_{40}$, the median concentration ratio between the third and first cycles was .96 (interquartile range, .92–.99; Friedman ANOVA $P = .015$). The median concentration ratio between the fourth and first cycles was .92 (interquartile range, .90–.96; Friedman ANOVA $P = .001$), and this survived the exclusion of the one clear outlier individual.

4. Discussion

We show that for up to four freeze-thaw cycles, each consisting of at least 1 hour of thawing to room temperature, in aliquot volumes of 200 μL , there is no consistent trend for change in the concentrations of serum NFL, plasma total tau, or plasma $\text{A}\beta_{42}$ as measured using the Simoa digital immunoassay platform. We found some evidence for a small reduction in plasma $\text{A}\beta_{40}$ after the third and further after the fourth cycle. This latter finding is in keeping with the experience of Lachno and colleagues, who used a Luminex bead-based immunoassay platform and a 1-hour thaw per cycle to demonstrate a reduction in plasma $\text{A}\beta_{1-40}$ from the fourth cycle in five individuals [2]. Here, we show, in a larger sample size, a statistically significant, albeit very small, reduction by a median ratio of 0.96 at

Download English Version:

<https://daneshyari.com/en/article/9987982>

Download Persian Version:

<https://daneshyari.com/article/9987982>

[Daneshyari.com](https://daneshyari.com)