

HEMATOLOGY/ONCOLOGY CLINICS OF NORTH AMERICA

INDEX

Α

Acquired immunodeficiency syndrome (AIDS), primary central nervous system lymphoma related to, 665-687 diagnostic considerations, 671-674 baseline evaluations, 671 - 672biopsy for, 698-699 early biopsy versus empiric toxoplasmosis therapy, 673 - 674evaluation of cerebrospinal fluid, 672-673 epidemiology and link with Epstein-Barr virus, 666-667 histopathology, 667 radiographic features, 667-671, 693-694 CT and MRI assessment, 667-670 metabolic imaging, 670-671 treatment options, 674-681 antiretroviral therapy, 674 - 675chemotherapy and combined modality therapy, 675 - 678Epstein-Barr virus-targeted therapies, 679-681 radiation therapy, 674

Age, as prognostic factor in primary central nervous system lymphoma, 630–631 at diagnosis of primary central nervous system lymphoma, 689

AIDS. See Acquired immunodeficiency syndrome.

Algorithm, for treatment of primary intraocular lymphoma, 744–746

Anatomic distribution, of lesions in primary central nervous system lymphoma, 690–91

Angiography, fluorescein, in diagnosis of primary intraocular lymphoma, 741

Antiepileptic medications, for seizures in primary central nervous system lymphoma, 623 Antiretroviral therapy, for AIDS-related primary central nervous system lymphoma, 674–675

Autologous stem cell transplantation, for central nervous system involvement in non-Hodgkin's lymphoma, 605–606 with intensive chemotherapy for primary central nervous system lymphoma, 722–725

В

B-cell primary central nervous system lymphoma, indolent, 661–662

Biopsy, in diagnosis of primary central nervous system lymphoma, 698–699 in AIDS patients, 673–674, 698–699

> in immunocompetent patients, 698 vitreous, in diagnosis of primary intraocular lymphoma, 742

Blood-brain barrier disruption, enhancement of chemotherapy by, in primary central nervous system lymphoma, 725

Bone marrow transplantation, for primary central nervous system lymphoma, 622–623

Brain delivery, of chemotherapy, enhancement of by blood-brain barrier disruption in primary central nervous system lymphoma, 725

C

Central nervous system, involvement of, in non-Hodgkin's lymphoma, 597–609 diagnosis, 600–602 cerebrospinal fluid analysis, 601–602 neuroimaging, 600–601 prophylaxis, 602 risk factors, 599 treatment, 602–606 transplantation, 605–606 relapse to, in lymphoma patients, 751–763

Note: Page numbers of article titles are in **boldface** type.

INDEX

Central nervous system lymphoma, primary,	T-cell, 658-661
611–749	treatment, 611-627
AIDS-related, 665–687	bone marrow transplanta-
diagnostic considerations,	tion, 622–623
671-674	chemotherapy, 614–622
epidemiology and link with	for recurrent, 623
Epstein-Barr virus,	radiotherapy alone, 613–614
666–667	steroids, 612–613
histopathology, 667 radiographic features,	symptomatic treatment, 623 treatment-related neurotoxicity,
667–671	729–738
treatment options, 674–681	mechanisms of, 729–730
clinical features, 689–692	neuroimaging findings,
age and gender, 689	730–731
anatomic distribution of	neurologic and neurocogni-
lesions, 690-691	tive sequelae, 732–735
predisposing conditions, 690	white matter changes and
symptoms and signs,	cognitive outcome,
691-692	731–732
time from symptom onset to	unusual variants of, 651-664
diagnosis, 692	neurolymphomatosis,
diagnosis, 692–700	652-653
biopsy, 698–699	primary leptomeningeal lym-
cerebrospinal fluid analysis,	phoma, 653–657
694–697	rare pathologic variants,
differentials, 692–693	657-662
neuroimaging, 693–694	secondary, 751–763
ocular evaluation, 697–698	clinical features and diagnosis,
staging, 699–700	754–755
indolent B-cell, 661–662	incidence, 751–754
new treatment approaches, 719–728	outcome after relapse, 758–760 prophylaxis for, 760–761
blood-brain barrier disrup-	risk factors for relapse, 755–758
tion and intra-arterial	•
chemotherapy, 725	Cerebrospinal fluid analysis, in primary cen-
intensive chemotherapy and	tral nervous system lymphoma, 694–697
autologous stem cell	AIDS-related, 672–673, 697
transplantation,	in immunocompetent patients,
722–725	695–697
Phase II trials of single-agent	in suspected central nervous system involvement in non-Hodgkin's
chemotherapyregimens,	lymphoma, 601–602
719–721	
rituximab, 721–722	Chemotherapy, for central nervous system
pathology and genetics, 705-717	involvement in non-Hodgkin's lym-
gross pathology, 706	phoma, 602–604
incidence, 705–706	for primary central nervous system lymphoma, 614–622
microscopic pathology and	AIDS-related, 675–678
immunophenotype, 706–707	enhancement of, by blood-brain
molecular pathology, 712–	barrier disruption, 725
714	high-dose methotrexate, 616–622
pathogenesis, 711–712	intensive, with autologous stem
prognostic factors, 612, 629–649	cell transplantation, 722–725
age and performance status,	intra-arterial, 725
630-631	phase II trials of single-agent regi-
new variables with potential	mens, 719–721
value as, 636–643	systemic lymphoma regimens,
predictors of treatment-re-	615–616
lated toxicity, 643–645	for primary intraocular lymphoma,
scoring systems for, 631–636	744–746

Download English Version:

https://daneshyari.com/en/article/10026869

Download Persian Version:

https://daneshyari.com/article/10026869

<u>Daneshyari.com</u>