

# Intravenous Neck Injections in a Drug Abuser Resulting in Infection of a Laryngocele

Jeevanan Jahendran, Abdullah Sani,<sup>1</sup> Philip Rajan,<sup>1</sup> Gurdeep Singh Mann and Balachandran Appoo,

Department of Otorhinolaryngology and Head and Neck Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur, and <sup>1</sup>Department of Otorhinolaryngology and Head and Neck Surgery, Hospital Ipoh, Ipoh, Malaysia.

A pyolaryngocele is an uncommon complication of a laryngocele that has become infected. We present a case of a pyolaryngocele that was probably due to repeated injections in the neck veins. The pathogenesis, clinical features and management are discussed in detail. [*Asian J Surg* 2005;28(1):41–4]

**Key Words:** pyolaryngocele, laryngocele, laryngeal carcinoma, neck mass, surgical management

## Introduction

Pyolaryngoceles are considered a clinical rarity and only 34 cases have been published up to the time of writing. Abnormal dilatation of the saccule forming an air sac in contact with the laryngeal opening is called a laryngocele. A laryngomucocele is a laryngocele filled with mucus. An infective process occurring within a laryngomucocele is called a pyolaryngocele. We present a case of pyolaryngocele in an intravenous drug abuser who injected into neck veins. He presented with a neck swelling and subacute airway obstruction.

## Case report

A 40-year-old Sikh male lorry driver presented with a progressively enlarging, painless left neck mass over a 2-month period. Subsequently, he developed hoarseness followed by worsening stridor over the next month. There was associated low-grade fever for which he was prescribed oral antibiotics. However, this did not resolve his symptoms.

In the previous year, he had noticed a similar swelling, but that had resolved spontaneously after a course of antibiotics without any complications. A point of interest was that he had

been an intravenous drug abuser for more than 10 years and had injected drugs directly into the major vessels of the neck (jugular veins) and lower limbs (femoral veins), as all other peripheral veins were already difficult to access.

A soft, cystic swelling measuring about 5 × 4 cm was palpable on the left lateral aspect of the neck anterior to the sternomastoid but displacing the muscle posteriorly. The mass and the skin overlying it did not show any signs of inflammation, but there was tenderness on deep palpation.

Laryngeal endoscopy revealed a smooth bulging mass in the left supraglottic region, which probably arose from the ventricle, extending superiorly. This caused the epiglottis and the laryngeal inlet to be pushed to the right and resulted in partial obstruction of the airway. The appearance of the true cords as well as the subglottic region was normal except that movement of the left vocal cord was hindered due to the presence of the mass.

Routine blood investigations were essentially normal except for leucocytosis; hepatitis and retroviral screening were also negative. Computed tomography (CT) showed a fluid-filled mass arising from the larynx and extending through the thyrohyoid membrane into the external soft tissue plane (Figures 1 and 2).

Address correspondence and reprint requests to Dr. Jeevanan Jahendran, Department of Otorhinolaryngology and Head and Neck Surgery, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, 56000 Cheras, Kuala Lumpur, Malaysia.

E-mail: drjj70@yahoo.com • Date of acceptance: 10 September 2003


He was started on intravenous antibiotics and surgery was performed after 48 hours as there was no clinical improvement. During the watchful waiting period, there was no further deterioration in airway obstruction. Diagnosis of a pyolaryngocele was made and the mass was excised using a transcervical approach. The mass was separated from the surrounding structures up to the thyroid cartilage. The upper half of the ala of the thyroid cartilage was excised to gain access to the internal component. The entire sac was removed without breaching the laryngeal mucosa or damaging the superior laryngeal neurovascular bundle. Direct laryngoscopy after closure of the external wound showed that the larynx and the surrounding structures had almost retained their normal

anatomy except for minimal oedema over the left supraglottic area.


On opening the sac, we found it filled with thick yellowish material (Figure 3). Histological examination showed a fibrous-walled cyst filled with leucocytic material. There was no evidence of malignancy. Material sent for cultures and test for acid-fast bacilli was negative. The postoperative period was uneventful and the patient was well on subsequent follow-up, with no evidence of recurrence.

### Discussion


The laryngeal ventricle has been described as a horizontally arranged, elliptical orifice extending from the notch in the thyroid cartilage to the anterior edge of the arytenoid cartilage. The extension from its base is called the saccule or appendix of the ventricle.<sup>1</sup> It lies between the medial surface of the thyroid lamina and the ventricular fold. When the saccule dilates excessively, rising higher than the upper border of the thyroid cartilage, it forms a laryngocele.<sup>2</sup> Virchow coined the term laryngocele in 1867, but Dominique Larrey reported this condition as early as 1829.<sup>3</sup> Laryngoceles can be internal, external or mixed. An internal laryngocele is confined to the interior of the larynx and extends posterosuperiorly into the false vocal cord and aryepiglottic fold. An external laryngocele extends superiorly to appear as a lateral neck mass through the opening in the thyrohyoid membrane for the superior laryngeal neurovascular bundle. It is called a mixed laryngocele when both features are present. The incidence of laryngoceles is estimated at 1 in 2.5 million persons per year, with internal laryngoceles accounting for 30%, external 26% and mixed 44%.<sup>1,3-5</sup> Of these lesions, 15% are bilateral and approximately 10% become infected to form a pyolaryngocele.<sup>6</sup>



**Figure 1.** Axial computed tomography scan of the neck showing a fluid-filled mass arising from within the larynx extending to the laterocervical region of the neck and pushing the sternocleidomastoid posteriorly and shifting the larynx to the contralateral side.



**Figure 2.** Coronal computed tomography scan showing the mass arising from the left ventricle of the larynx passing through the thyrohyoid membrane to reach the lateral aspect of the neck.



**Figure 3.** The sac removed in its entirety. Thick yellowish discharge can be seen oozing from the sac (arrow).

Download English Version:

<https://daneshyari.com/en/article/10103050>

Download Persian Version:

<https://daneshyari.com/article/10103050>

[Daneshyari.com](https://daneshyari.com)