



Magnetic seed and cosmology as quantum hall effect

H. Falomir^a, J. Gamboa^{b,*}, P. Gondolo^c, F. Méndez^b

^a Departamento de Física, Universidad Nacional de La Plata, La Plata, Argentina

^b Departamento de Física, Universidad de Santiago de Chile, Casilla 307, Santiago, Chile

^c Department of Physics, University of Utah, Salt Lake City, Utah, USA

ARTICLE INFO

Article history:

Received 3 July 2018

Received in revised form 27 July 2018

Accepted 4 August 2018

Available online 30 August 2018

Editor: M. Trodden

ABSTRACT

In the framework of a bimetric model, we discuss a relation between the (modified) Friedmann equations and a mechanical system similar to the quantum Hall effect problem. Firstly, we show how these modified Friedmann equations are mapped to an anisotropic two-dimensional charged harmonic oscillator in the presence of a constant magnetic field, with the frequencies of the oscillator playing the role of the cosmological constants. This problem has two energy scales leading to the identification of two different regimes, namely, one dominated by the cosmological constants, with exponential expansions for the scale factors, and the other dominated by a magnetic seed, which would be responsible for both a component of dark energy and a primordial magnetic field. The latter regime would be described by a (nonperturbative) mapping between the cosmological evolution and the quantum Hall effect.

© 2018 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

The standard description of the Universe rests on the cosmological principle, which states that, on large scales, space-time is homogeneous and isotropic. The observations are consistent with this hypothesis for distances above 100 Mpc [1,2]. But this mathematical idealization, which greatly simplifies the physical interpretation of the model, has limitations for lower scales. In particular, the formation of structures can only be understood after the occurrence of some gravitational instability due to tiny deviations from a homogeneous distribution [3,4].

These departures from the cosmological principle can be observed, for example, in the spectrum of the cosmic microwave background (CMB), which presents temperature fluctuations of the order of 10^{-5} , showing that corrections to classical cosmology can be incorporated via perturbations [5].

However, one might wonder if there are other phenomena of cosmological interest that might require a non-perturbative analysis. This possibility is particularly relevant since, in many fields of physics, there are problems that are perturbative or non-perturbative depending on the range of parameters one is considering. As an example, one can consider a gas of charged particles subject to a magnetic field perpendicular to the plane. If the magnetic field is strong enough, the system presents the quantum Hall

effect, with a Hamiltonian spectrum that can not be perturbatively obtained from that of the free case.

This simple example could also be translated into the cosmological regime by noting that in the center of galaxies there are strong magnetic fields which are observed through the Zeeman's splitting they produce. Although the origin of these magnetic fields is at present unknown, the idea that a very small magnetic seed was formed in an early epoch of the universe evolution and that, after a dynamo mechanism, the field grew up to what is observed today in galaxies is widely accepted [6–10,25]. Our present knowledge does not allow us to determine when these magnetic seeds were created, but one can speculate that they might have been originated in the small inhomogeneities existing before the recombination epoch.

Very probably the primordial magnetic fields did not produce any relevant effect after the recombination, but these could be important in the first 100.000-years and eventually to affect the big-bang nucleosynthesis, the dynamics of the phase transitions and even baryogenesis and leptogenesis [11].

The magnetic seed must satisfy two consistency requirements. The first one is that the coherence length is not larger than about 10 kpc, and the second one is that the field in the magnetic seed must be between 10^{-19} and 10^{-22} G. In the analogous Hall system we discuss below, the coherence length corresponds to the magnetic depth ℓ_B ¹ [12], that is,

* Corresponding author.

E-mail addresses: falomir@fisica.unlp.edu.ar (H. Falomir), jorge.gamboa@usach.cl (J. Gamboa), paolo.gondolo@utah.edu (P. Gondolo), fernando.mendez@usach.cl (F. Méndez).

¹ Here we use natural units and $e = 1$ [13].

$$\ell_B = \frac{1}{\sqrt{B}} < 10 \text{ kpc.} \quad (1)$$

The second condition is necessary for the stability of the dynamo mechanism [8–10].

A central issue not solved so far is how to provide the cosmological standard model with a mechanism that incorporates a magnetic seed as a fundamental element [14]. Any possible answer to this question requires extra new ideas in a model that satisfies all constraints known so far and that incorporates the magnetic field as a central element.

In this direction and using arguments coming from the formation of primordial magnetic fields [8–10] (we say for $t \sim 10^6$ years), the mechanism proposed here should work.

The purpose of this paper consists in investigating the possible emergence of magnetic seeds in a model with two metrics with an effective interaction between them. This interaction can be considered as a relic of a causal primordial connection between sectors in a very early epoch of the Universe. This problem is considered in the context of a simple mechanical system that nevertheless reproduces the Friedmann's equations of the two interacting sectors. We emphasize that the important issue is not only the existence of a mapping between these apparently unrelated systems but also that the same mechanism contributes to the production of dark energy.

Interestingly, no matter how different the dark energy and magnetic seed scales might be since in the present approach both are linked through a dynamical mechanism which (see Eq. (11)) allows to fix them in a rather independent way.

In order to develop this idea let us consider the Lagrangian²

$$L = \frac{1}{2N} (\dot{x}_1^2 + \dot{x}_2^2) - \frac{N}{2} (\omega_1^2 x_1^2 + \omega_2^2 x_2^2) - \frac{\theta}{2} (x_1 \dot{x}_2 - \dot{x}_1 x_2). \quad (2)$$

Here x_1 and x_2 are the dynamical variables, the coefficients ω_1, ω_2 and θ are constants, and $N = N(t)$ is an auxiliary variable that transforms as $N(t) \rightarrow t'(s)N(t(s))$ when $t \rightarrow t(s)$, thus ensuring the invariance of the action under time reparametrizations. This Lagrangian yields the following Hamiltonian

$$H = \frac{N}{2} \left[p_1^2 + p_2^2 + \left(\omega_1^2 + \frac{\theta^2}{4} \right) x_1^2 + \left(\omega_2^2 + \frac{\theta^2}{4} \right) x_2^2 + \theta (x_1 p_2 - x_2 p_1) \right]. \quad (3)$$

This Hamiltonian describes an anisotropic two-dimensional charged harmonic oscillator with frequencies ω_1 and ω_2 , interacting with a constant magnetic field.

The Hamiltonian equations of motion for (3) are

$$\dot{x}_i = [x_i, H], \quad \dot{p}_i = [p_i, H],$$

where $[,]$ is the Poisson bracket, with the standard structure for the canonical variables, that is $[x_i, p_j] = \delta_{ij}$ and zero for the remaining brackets. Alternatively, one can define the new variables $\pi_i = p_i - \frac{\theta}{2} \epsilon_{ij} x_j$ and rewrite the Hamiltonian $H = H(x_i, \pi_j)$ in order to obtain the equations of motion

$$\dot{x}_i = [x_i, H], \quad \dot{\pi}_i = [\pi_i, H], \quad (4)$$

but with the following Poisson brackets

$$[x_i, x_j] = 0, \quad [x_i, \pi_j] = \delta_{ij}, \quad [\pi_i, \pi_j] = \epsilon_{ij} \theta. \quad (5)$$

The equations of motion, once the momenta are eliminated, reduce to

$$\ddot{x}_1 + \omega_1^2 x_1 + \theta \dot{x}_2 = 0, \quad (6)$$

$$\ddot{x}_2 + \omega_2^2 x_2 - \theta \dot{x}_1 = 0, \quad (7)$$

$$\dot{x}_1^2 + \dot{x}_2^2 + \omega_1^2 x_1^2 + \omega_2^2 x_2^2 = 0. \quad (8)$$

The constraint (8) is a consequence of time reparametrization invariance and, at the end of the derivation, the gauge $N \equiv 1$ has been chosen.

Notice that this constraint – from the point of view of the second order differential equations (6) and (7) – is in fact a relation between initial conditions since the left hand side is a constant of the motion. Indeed, multiplying (6) by \dot{x}_1 and (7) by \dot{x}_2 , and adding both equations, we immediately find that

$$\frac{d}{dt} [\dot{x}_1^2 + \dot{x}_2^2 + \omega_1^2 x_1^2 + \omega_2^2 x_2^2] = 0. \quad (9)$$

The physical solutions correspond to those for which the constant $\dot{x}_1^2 + \dot{x}_2^2 + \omega_1^2 x_1^2 + \omega_2^2 x_2^2$ vanishes.

One of the goals of this paper is to point out the following remarkable mapping. If we redefine the variables x_1, x_2 as follows,

$$x_1 = \frac{2}{3} a^{3/2}(t), \quad x_2 = \frac{2}{3} b^{3/2}(t), \quad (10)$$

and replace them in (6)–(7), the resulting equations turn out to be

$$2 \frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 + \frac{4}{3} \omega_1^2 = -2\theta \sqrt{ab} \frac{\dot{b}}{a^2}, \quad (11)$$

$$2 \frac{\ddot{b}}{b} + \left(\frac{\dot{b}}{b} \right)^2 + \frac{4}{3} \omega_2^2 = 2\theta \sqrt{ab} \frac{\dot{a}}{b^2}, \quad (12)$$

$$a^3 \left[\left(\frac{\dot{a}}{a} \right)^2 + \left(\frac{2}{3} \omega_1 \right)^2 \right] = -b^3 \left[\left(\frac{\dot{b}}{b} \right)^2 + \left(\frac{2}{3} \omega_2 \right)^2 \right]. \quad (13)$$

These equations are identical to the Friedmann equations for a cosmology with two metrics³ if we identify their respective cosmological constants Λ_1 and Λ_2 as

$$-\omega_1^2 \longleftrightarrow \frac{3}{4} \Lambda_1, \quad -\omega_2^2 \longleftrightarrow \frac{3}{4} \Lambda_2. \quad (14)$$

In fact, Eqs. (11)–(13) form a coupled system of nonlinear second order differential equations for the scale factors $a(t)$ and $b(t)$, where the right hand sides of (11)–(12) can be considered as sources of dark energy (see [17] for a discussion on a similar system and for string theory see [18]). Moreover, from these equations one can read off the effective pressure and density contributions induced by the coupling between scale factors. Indeed, expressing the Friedmann equations for the scale factor $a(t)$ in terms of the pressure p_b and energy density ρ_b of an additional component of “dark energy”, from Eqs. (11) and (13) one obtains the equivalence

$$8\pi G p_b = -2\theta \sqrt{ab} \frac{\dot{b}}{a^2},$$

$$\frac{8\pi G}{3} \rho_b = -\frac{1}{a^3} \left(\frac{4}{9} \omega_2^2 b^3 + \dot{b}^2 b \right).$$

This leads to the following equation of state for the effective component of dark energy,

² The approach proposed here is valid for any number of patches, however for simplicity in the presentation we will restrict ourselves to two of them.

³ The literature of cosmology with two metrics is very extensive, see for example [15] and [16].

Download English Version:

<https://daneshyari.com/en/article/10136801>

Download Persian Version:

<https://daneshyari.com/article/10136801>

[Daneshyari.com](https://daneshyari.com)