Accepted Manuscript



Structure and function of the histone chaperone FACT – Resolving FACTual issues

Katerina Gurova, Han-Wen Chang, Maria E. Valieva, Poorva Sandlesh, Vasily M. Studitsky

PII:	S1874-9399(18)30159-7
DOI:	doi:10.1016/j.bbagrm.2018.07.008
Reference:	BBAGRM 94291
To appear in:	*BBA - Gene Regulatory Mechanisms*
Received date:	16 April 2018
Revised date:	17 July 2018
Accepted date:	19 July 2018

Please cite this article as: Katerina Gurova, Han-Wen Chang, Maria E. Valieva, Poorva Sandlesh, Vasily M. Studitsky , Structure and function of the histone chaperone FACT – Resolving FACTual issues. Bbagrm (2018), doi:10.1016/j.bbagrm.2018.07.008

Structure and function of the histone chaperone FACT – resolving FACTual issues

Katerina Gurova[a,*], Han-Wen Chang[b], Maria E. Valieva[c], Poorva Sandlesh[a], Vasily M. Studitsky[b,c,*]

[a] Department of Cell Stress Biology, Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, NY, 14263, USA
[b] Fox Chase Cancer Center, Philadelphia, PA 19111, USA
[c] Biology Faculty, Lomonosov Moscow State University; Moscow, Russia

* Corresponding authors: Katerina Gurova, katerina.gurova@roswellpark.org, Vasily Studitsky, Vasily.Studitsky@fccc.edu

Highlights

The best-established FACT function is nucleosome maintenance during transcription.

The role of in FACT in facilitating transcription through chromatin in vivo requires further evaluation.

Aggressive tumors are more sensitive to FACT depletion than normal tissues.

The elevated requirement for FACT in tumor cells is associated with increased sensitivity to chromatin destabilization.