

Accepted Manuscript

Title: Methadone maintenance treatment among patients exposed to illicit fentanyl in Rhode Island; safety, dose, retention, and relapse at 6 Months

Authors: Andrew C. Stone, Jennifer J. Carroll, Josiah D. Rich, Traci C. Green



PII: S0376-8716(18)30472-1
DOI: <https://doi.org/10.1016/j.drugalcdep.2018.07.019>
Reference: DAD 7090

To appear in: *Drug and Alcohol Dependence*

Received date: 11-6-2018
Revised date: 14-7-2018
Accepted date: 22-7-2018

Please cite this article as: Stone AC, Carroll JJ, Rich JD, Green TC, Methadone maintenance treatment among patients exposed to illicit fentanyl in Rhode Island; safety, dose, retention, and relapse at 6 Months, *Drug and Alcohol Dependence* (2018), <https://doi.org/10.1016/j.drugalcdep.2018.07.019>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Methadone maintenance treatment among patients exposed to illicit fentanyl in Rhode Island; safety, dose, retention, and relapse at 6 Months

Andrew C. Stone^a, Jennifer J. Carroll^b, Josiah D. Rich^b, Traci C. Green^b

^a Discovery House Comprehensive Treatment Center, 1625 Diamond Hill Road, Woonsocket, RI 02895, USA.

^b The Warren Alpert School of Medicine of Brown University, Rhode Island Hospital, 55 Claverick Street, Providence, RI 02903, USA.

Correspondence

Andrew C. Stone

Discovery House Comprehensive Treatment Center

1625 Diamond Hill Road

Woonsocket, RI 02895, USA

ACStoneMD@gmail.com

Highlights

- First study to investigate outcomes in methadone maintenance treatment (MMT) patients exposed to fentanyl.
- Abstinence achieved by the majority (89%) of fentanyl-exposed MMT patients.
- Repeated exposure to fentanyl was seen frequently (71%) while in MMT.
- There were no deaths for those who remained in MMT.

Download English Version:

<https://daneshyari.com/en/article/10147217>

Download Persian Version:

<https://daneshyari.com/article/10147217>

[Daneshyari.com](https://daneshyari.com)